arXiv:1410.8636v1 [math.AP] 31 Oct 2014

ON THE DECAY AND STABILITY OF GLOBAL SOLUTIONS TO THE 3D INHOMOGENEOUS MHD SYSTEM

JUNXIONG JIA, JIGEN PENG, AND KEXUE LI

Abstract. In this paper, we investigative the large time decay and stability to any given global smooth solutions of the 3D incompressible inhomogeneous MHD systems. We proved that given a solution (a, u, B) of (1.2), the velocity field and magnetic field decay to 0 with an explicit rate, for u which coincide with incompressible inhomogeneous Navier-Stokes equations [21]. In particular, we give the decay rate of higher order derivatives of u and B which is useful to prove our main stability result. For a large solutions of (1.2) denoted by (a, u, B), we proved that a small perturbation to the initial data still generates a unique global smooth solution and the smooth solution keeps close to the reference solution (a, u, B). Due to the coupling between u and B, we used elliptic estimates to get $\|(u, B)\|_{L^1(\mathbb{R}^+;\dot{B}^{5/2}_{2,1})} < C$, which is different to Navier-Stokes equations.

1. Introduction and main results

Magnetic fields influence many fluids. Magnetohydrodynamics (MHD) is concerned with the interaction between fluid flow and magnetic field. The governing equations of nonhomogeneous MHD can be stated as follows [1],

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \quad (t, x) \in \mathbb{R}^+ \times \mathbb{R}^3, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) - \operatorname{div}(\mu \mathcal{M}) - (B \cdot \nabla)B + \nabla \Pi = 0, \\ \partial_t B - \lambda \Delta B - \operatorname{curl}(u \times B) = 0, \\ \operatorname{div} u = \operatorname{div} B = 0, \\ \rho|_{t=0} = \rho_0, \quad u|_{t=0} = u_0, \quad B|_{t=0} = B_0, \end{cases} \tag{1.1}$$

where ρ, $u = (u_1, u_2, u_3)$ stand for the density and velocity of the fluid, respectively, $\mathcal{M} = \frac{1}{2}(\partial_i u_j + \partial_j u_i)$, Π is a scalar pressure function. B is the magnetic field. $\mu(\rho) \geq 0$ denotes the viscosity of fluid, which we assume in this paper is a positive constant. $\lambda > 0$ is also a constant, which describes the relative strength of advection and diffusion of B.

If there is no magnetic field, i.e., $B = 0$, MHD system turns to be nonhomogeneous Navier-Stokes system. Since the second equation and third equation of (1.1) are similar, the study about MHD system has been along with that for Navier-Stokes one. Let us first recall some results about Navier stokes equations. When ρ_0 is bounded away from zero, the global existence of weak solutions was established by Kazhikov [4]. Moreover, Antontsev, Kazhikov and Monakhov [5] gave the first

2010 *Mathematics Subject Classification.* 35Q35, 76D03, 76W05.

Key words and phrases. Inhomogeneous MHD system, Stability of large solution, Decay rate, Besov space.

result on local existence and uniqueness of strong solutions. For the two dimensional case, they even proved that the strong solution is global. But the global existence of strong or smooth solutions in 3D is still an open problem.

Recently, R.Danchin proved the global existence in the Besov space framework [6]. His result states the global in time existence of regular solutions to the inhomogeneous Navier Stokes equations in $\mathbb{R}^n$ in the optimal Besov setting, under suitable smallness of the data. In particular his results allows the initial densities have a jump at the interface. At the same time, H. Abidi, G. Gui, P. Zhang [7] proved the local well-posedness of three-dimensional incompressible inhomogeneous Navier-Stokes equations with initial data in the critical Besov spaces, without assumptions of small density variation. And they also proved the global well-posedness when the initial velocity is small in $\dot{B}^{1/2}_{2,1}(\mathbb{R}^3)$. For more results in this direction, see [8, 9, 10]and reference therein.

Now, let us go back to the MHD system (1.1). As said before, the research for MHD goes along with that for Naiver-Stokes equations. The results are similar. When we assume ρ is a constant, which means the fluid is homogeneous, the MHD system has been extensively studied. Duraut and Lions [11] constructed a class of weak solutions with finite energy and a class of local strong solutions. Recently, C. Cao, J. Wu [12] proved global regularity of classical solutions for the MHD equations with mixed partial dissipation and magnetic diffusion. And they also give the global existence, conditional regularity and uniqueness of a weak solution for 2D MHD equations with only magnetic diffusion. For more results in this direction, see [13, 14]and reference therein.

When the fluid is nonhomogeneous. H. Abidi and M. Paicu [3] proved that the magneto-hydrodynamic system in $\mathbb{R}^N$ with variable density, variable viscosity and variable conductivity has a local weak solution in suitable Besov space if the initial density approaches a constant. And they also proved that the constructed solution exist globally in time if the initial data are small enough. X. Huang and Y. Wang [2] proved the global existence of strong solution with vacuum to the 2D nonhomogeneous incompressible MHD system, as long as the initial data satisfies some compatibility condition. In this paper, we only consider non-vacuum case.

Let $a := \frac{1}{\rho} - 1$ and take $\mu = \lambda = 1$, then MHD system becomes

$$
\begin{cases}
\partial_t a + u \cdot \nabla a = 0, \qquad (t,x) \in \mathbb{R}^+ \times \mathbb{R}^3, \\
\partial_t u + u \cdot \nabla u - (1+a)(\Delta u - \nabla \Pi) - (1+a)(B \cdot \nabla) B = 0, \\
\partial_t B - \Delta B + (u \cdot \nabla) B - (B \cdot \nabla) u = 0, \\
\mathrm{div} u = \mathrm{div} B = 0, \\
(a, u, B)|_{t=0} = (a_0, u_0, B_0).
\end{cases} \tag{1.2}
$$

or let $\rho := \frac{1}{1+a}$ and (ρ, u, B) solves

$$
\begin{cases}
\partial_t \rho + \mathrm{div}(\rho u) = 0, \qquad (t,x) \in \mathbb{R}^+ \times \mathbb{R}^3, \\
\rho \partial_t u + \rho u \cdot \nabla u - \Delta u - (B \cdot \nabla) B + \nabla \Pi = 0, \\
\partial_t B - \Delta B + (u \cdot \nabla) B - (B \cdot \nabla) u = 0, \\
\mathrm{div} u = \mathrm{div} B = 0, \\
(\rho, u, B)|_{t=0} = (\rho_0, u_0, B_0).
\end{cases} \tag{1.3}
$$

In what follows, we shall investigate the large time decay and stability for the above MHD system. Compared to the classical incompressible Navier-Stokes system (NS), that is, in the case when $a = 0$ and $B = 0$ in (1.2), the MHD system is more complex. Given a global large solution $u \in L^\infty_{\rm loc}([0,\infty); H^1(\mathbb{R}^3)) \cap L^2_{\rm loc}([0,\infty); H^2(\mathbb{R}^2(\mathbb{R}^3)))$, Ponce, Racke, Sideris and Titi [15] proved (NS) global stability under the additional assumption $\int_0^\infty \|\nabla u(t)\|_{L^2}^4\, dt < \infty$. Then Gallagher, Iftimie and Planchon [16] removed the additional assumption. For the inhomogeneous Navier-Stokes equations (INS), H. Abidi, G. Gui and P. Zhang [21] proved its decay and stability for large solutions.

Our first result concerns the global stability of the given solution of (1.2) when the initial density ρ_0 is close to a positive constant. This is a simple generalization of Theorem 1.1 in [21].

Theorem 1.1. *Let $\bar a_0 \in B^{5/2}_{2,1}(\mathbb{R}^3)$, $\bar u_0 \in B^{3/2}_{2,1}(\mathbb{R}^3)$ and $\bar B_0 \in B^{3/2}_{2,1}(\mathbb{R}^3)$ with $\mathrm{div}\bar u_0 = \mathrm{div}\bar B_0 = 0$, and let there exist two positive constants m and M so that*

$$m \le 1 + \bar a_0 \le M. \tag{1.4}$$

We assume that $\bar a \in C([0,\infty); B^{5/2}_{2,1}(\mathbb{R}^3))$ and

$$\bar u, \bar B \in C([0,\infty); B^{3/2}_{2,1}(\mathbb{R}^3)) \cap L^1_{loc}(\mathbb{R}^+; \dot B^{7/2}_{2,1}(\mathbb{R}^3))$$

is a given solution of MHD with initial data $(\bar a_0, \bar u_0)$. Then there exist positive constants c_1, C_1 and a large enough time $T_0 := T_0(\bar a_0, \bar u_0, \bar B_0)$ so that if

$$\|\bar a_0\|_{\dot B^{3/2}_{2,1}} \exp\left\{ C_1 \int_0^{T_0} \|\nabla \bar u(\tau)\|_{\dot B^{3/2}_{2,1}}\, d\tau \right\} \le c_1, \tag{1.5}$$

a constant c_2 exists so that $(a_0, u_0, B_0) := (\bar a_0 + \tilde a_0, \bar u_0 + \tilde u_0, \bar B_0 + \tilde B_0)$ generates a unique global solution with

$$\begin{aligned} &a \in C_b([0,\infty); B^{5/2}_{2,1}(\mathbb{R}^3)),\\ &u, B \in C_b([0,\infty); B^{3/2}_{2,1}(\mathbb{R}^3)) \cap L^1([0,\infty); \dot B^{7/2}_{2,1}(\mathbb{R}^3)), \end{aligned} \tag{1.6}$$

provided that $(\tilde a_0, \tilde u_0, \tilde B_0)$ satisfies

$$\|\tilde a_0\|_{B^{3/2}_{2,1}} + \|\tilde u_0\|_{B^{1/2}_{2,1}} + \|\tilde B_0\|_{B^{1/2}_{2,1}} \le c_2. \tag{1.7}$$

In order to get the stability of large solutions of system (1.2), here, we need to investigate the decay properties of the velocity field u and magnetic field B. Compared to the (INS) case, our case is more complex and we need to use the coupling between the equations of u and B. Due to the coupling between u and B, we can not get the estimate of $\|(u,B)\|_{L^1(\mathbb{R}^+; \dot B^{5/2}_{2,1})} < C$ by using propositions like Proposition 3.6 in [21], which is the methods used in (INS) case. In order to overcome this difficulty, we need to give the estimate of $\|\nabla a(t)\|_{L^\infty}$, so that in addition to get the higher order decay properties of u and B. The rigorous statement is the following Theorem.

Theorem 1.2. *For $p \in (1, \frac{6}{5})$, let $a_0 \in B^{5/2}_{2,1}$ and $u_0 \in L^p(\mathbb{R}^3) \cap B^2_{2,1}(\mathbb{R}^3)$ satisfying (1.4) and $\mathrm{div}\, u_0 = 0$. We assume that $a \in C([0,\infty); B^{5/2}_{2,1})$, $u \in C([0,\infty); B^2_{2,1}(\mathbb{R}^3)) \cap L^1_{\rm loc}(\mathbb{R}^+; \dot B^4_{2,1}(\mathbb{R}^3))$ and $B \in C([0,\infty); B^2_{2,1}(\mathbb{R}^3)) \cap L^1_{\rm loc}(\mathbb{R}^+; \dot B^4_{2,1}(\mathbb{R}^3))$ is a given*

global solution of (1.2) with initial data (a_0, u_0, B_0). *Then there exists a positive time* t_0 *such that there hold*

$$\begin{aligned}
&\|u(t)\|_{L^2} + \|B(t)\|_{L^2} \le C\,(1+t)^{-\beta(p)},\\
&\|\nabla u(t)\|_{L^2} + \|\nabla B(t)\|_{L^2} \le C\,(1+t)^{-\frac{1}{2}-\beta(p)}, \quad \textit{for } t \ge t_0,\\
&\int_{t_0}^{\infty} (1+t)^{(1+2\,\beta(p))^-} \Big(\|(\partial_t u, \partial_t B)\|_{L^2}^2 + \|(\Delta u, \Delta B)\|_{L^2}^2 + \|\nabla \Pi\|_{L^2}^2 \Big)\, dt \le C,\\
&\int_{t_0}^{\infty} \Big(\|u(t)\|_{L^\infty} + \|B(t)\|_{L^\infty} + \|\nabla u(t)\|_{L^\infty} + \|\nabla B(t)\|_{L^\infty} \Big)\, dt \le C
\end{aligned} \tag{1.8}$$

and

$$\begin{aligned}
&\sup_{t\ge t_0} \Big(\|\nabla^2 u(t)\|_{L^2}^2 + \|\nabla^2 B(t)\|_{L^2}^2 \Big) + \int_{t_0}^{\infty} \|\partial_t \nabla u(t)\|_{L^2}^2 + \|\partial_t \nabla B(t)\|_{L^2}^2\, dt\\
&\qquad + \int_{t_0}^{\infty} \|\nabla^3 u(t)\|_{L^2}^2 + \|\nabla^3 B(t)\|_{L^2}^2\, dt \le\ C,
\end{aligned} \tag{1.9}$$

where $\beta(p) = \frac{3}{4}(\frac{2}{p} - 1)$. $(1 + 2\beta(p))^-$ *denotes any positive number smaller than* $1 + 2\beta(p)$, *and the constant* C *depends on the initial data.*

At the above theorem in hand, we can use the elliptic estimates and various interpolation theorems in Besov space to get the estimate of $\|(u, B)\|_{L^1(\mathbb{R}^+;\dot{B}^{5/2}_{2,1})} < C$ which is completely different to (INS) case. Then after complex calculations, we can get the global estimates of the reference solution $(\bar{a}, \bar{u}, \bar{B})$. At last, similar to Theorem 1.2 but need more complex calculations, we obtain the decay properties of the perturbed solution $(a - \bar{a}, u - \bar{u}, B - \bar{B})$. Using the decay properties of the reference solution and perturbed solution, we finally got the following theorem.

Theorem 1.3. *For* $p \in (1, \frac{6}{5})$, *let* $\bar{a}_0 \in B^{7/2}_{2,1}(\mathbb{R}^3)$, $\bar{u}_0 \in L^p(\mathbb{R}^3) \cap B^2_{2,1}(\mathbb{R}^3)$, $\bar{B}_0 \in L^p(\mathbb{R}^3) \cap B^2_{2,1}(\mathbb{R}^3)$ *satisfy* $\operatorname{div} \bar{u}_0 = \operatorname{div} \bar{B}_0 = 0$ *and (1.4). We assume that* $\bar{a} \in C([0,\infty); B^{7/2}_{2,1}(\mathbb{R}^3))$, $\bar{u} \in C([0,\infty); B^2_{2,1}(\mathbb{R}^3)) \cap L^1_{\rm loc}(\mathbb{R}^+; \dot{B}^4_{2,1})$, $\bar{B} \in C([0,\infty); B^2_{2,1}(\mathbb{R}^3)) \cap L^1_{\rm loc}(\mathbb{R}^+; \dot{B}^4_{2,1})$ *is a given global solution of (1.2) with initial data* $(\bar{a}_0, \bar{u}_0, \bar{B}_0)$. *Then there exists a constant* c *so that if*

$$(\tilde{a}_0, \tilde{u}_0, \tilde{B}_0) \in B^{7/2}_{2,1}(\mathbb{R}^3) \times \Big(L^p(\mathbb{R}^3) \cap B^2_{2,1}(\mathbb{R}^3) \Big) \times \Big(L^p(\mathbb{R}^3) \cap B^2_{2,1}(\mathbb{R}^3) \Big)$$

with

$$A_0 := \|(\tilde{u}_0, \tilde{B}_0)\|_{H^1} + \|(\tilde{u}_0, \tilde{B}_0)\|_{L^p} + \|\tilde{a}_0\|_{B^{3/2}_{2,1}} \le c,$$

$(a_0, u_0, B_0) := (\bar{a}_0 + \tilde{a}_0, \bar{u}_0 + \tilde{u}_0, \bar{B}_0 + \tilde{B}_0)$ *generates a unique global smooth solution* (a, u, B) *to (1.2) that satisfies*

$$\begin{aligned}
&a \in C_b([0,\infty); B^{7/2}_{2,1}(\mathbb{R}^3)),\\
&u \in C_b([0,\infty); L^p(\mathbb{R}^3 \cap B^2_{2,1}(\mathbb{R}^3))) \cap L^1(\mathbb{R}^+; \dot{B}^4_{2,1}(\mathbb{R}^4)),\\
&B \in C_b([0,\infty); L^p(\mathbb{R}^3 \cap B^2_{2,1}(\mathbb{R}^3))) \cap L^1(\mathbb{R}^+; \dot{B}^4_{2,1}(\mathbb{R}^4)).
\end{aligned}$$

Moreover, there holds

$$\|a - \tilde{a}\|_{\tilde{L}^\infty(\mathbb{R}^+; B^{s+1}_{2,1})} \le C A_0^{\frac{5}{4} - \frac{1}{2}s} \tag{1.10}$$

for any $s \in [\frac{1}{2}, \frac{5}{2}]$ *and*

$$\begin{aligned}\|(u-\bar{u}, B-\bar{B})\|_{\tilde{L}^\infty(\mathbb{R}^+;B^s_{2,1})} + \|(u-\bar{u}, B-\bar{B})\|_{L^\infty(\mathbb{R}^+;L^p)} \\ + \|(u-\bar{u}, B-\bar{B})\|_{L^1(\mathbb{R}^+;\dot{B}^{s+2}_{2,1})} \le CA_0^{\frac{4}{3}-\frac{2}{3}s}\end{aligned} \tag{1.11}$$

for any $s \in [\frac{1}{2}, 2]$.

Remark 1.4. The above Theorems may not be obtained by regarding the term $B \cdot \nabla B$ as a source term in the velocity equation. The reason is that if we regard this term as a source term, we will encounter terms like $\bar{B} \cdot \nabla \tilde{B}$ and $\bar{B} \cdot \nabla \tilde{u}$ in (3.16) and (6.43). For the appearance of these terms, the Bootstrap argument will not work. So we consider the linear system (2.12) is necessary and the higher order decay estimates is also necessary to get the results for MHD system.

The paper is organized as follows. In section 2, we will give some notations, a brief introduction to the Besov space and some useful Lemmas. In section 3, as a warm up, we give the proof of Theorem 1.1. Then, in section 4, we proved Theorem 1.2 in a series of propositions. Using Theorem 1.2, we got the global estimates of the reference solutions in section 5. At last, we proved the decay properties of the perturbed solutions and Theorem 1.3 in section 6. In the appendix, we proved some simple Lemmas which will be used in the above sections.

2. Preliminaries

Throughout this paper we will use the following notations.

- For any tempered distribution u both $\widehat{u}$ and $\mathcal{F}u$ denote the Fourier transform of u.
- The norm in the mixed space-time Lebesgue space $L^p([0,T];L^r(\mathbb{R}^d))$ is denoted by $\|\cdot\|_{L^p_T L^r}$ (with the obvious generalization to $\|\cdot\|_{L^p_T X}$ for any normed space X).
- For X a Banach space and I an interval of $\mathbb{R}$, we denote by $C(I;X)$ the set of continuous functions on I with values in X, and by $C_b(I;X)$ the subset of bounded functions of $C(I;X)$.
- For any pair of operators P and Q on some Banach space X, the commutator $[P,Q]$ is given by $PQ-QP$.
- C stands for a "harmless" constant, and we sometimes use the notation $A \lesssim B$ as an equivalent of $A \le CB$. The notation $A \approx B$ means that $A \lesssim B$ and $B \lesssim A$.
- $\{c_{j,r}\}_{j\in\mathbb{Z}}$ a generic element of the sphere of $\ell^r(\mathbb{Z})$, and $(c_k)_{k\in\mathbb{Z}}$ (respectively, $(d_j)_{j\in\mathbb{Z}}$) a generic element of the sphere of $\ell^2(\mathbb{Z})$ (respectively, $\ell^1(\mathbb{Z})$).
- Denote γ^- be any number smaller than γ.

Then, we give a short introduction to the Besov type space. Details about Besov type space can be found in [17] or [18]. There exist two radial positive functions $\chi \in \mathcal{D}(\mathbb{R}^d)$ and $\varphi \in \mathcal{D}(\mathbb{R}^d \backslash \{0\})$ such that

- $\chi(\xi) + \sum_{q\ge 0} \varphi(2^{-q}\xi) = 1$; $\forall q \ge 1$, $\mathrm{supp}\chi \cap \mathrm{supp}\varphi(2^{-q}\cdot) = \phi$,
- $\mathrm{supp}\varphi(2^{-j}\cdot) \cap \mathrm{supp}\varphi(2^{-k}\cdot) = \phi$, if $|j-k| \ge 2$,

For every $v \in S'(\mathbb{R}^d)$ we set

$$\Delta_{-1}v = \chi(D)v, \quad \forall q \in \mathbb{N}, \quad \Delta_j v = \varphi(2^{-q}D)v \quad \text{and} \quad S_j = \sum_{-1\le m\le j-1} \Delta_m.$$

The homogeneous operators are defined by

$$\dot{\Delta}_q v = \varphi(2^{-q}D)v, \quad \dot{S}_j = \sum_{m\leq j-1} \dot{\Delta}_j v, \quad \forall q \in \mathbb{Z}.$$

One can easily verifies that with our choice of φ,

$$\Delta_j \Delta_k f = 0 \quad \text{if} \quad |j-k| \geq 2 \tag{2.1}$$

$$\Delta_j (S_{k-1} f \Delta_k f) = 0 \quad \text{if} \quad |j-k| \geq 5. \tag{2.2}$$

As in Bony's decomposition, we split the product uv into three parts

$$uv = T_u v + T_v u + R(u,v),$$

with

$$T_u v = \sum_j S_{j-1} u \Delta_j v,$$

$$R(u,v) = \sum_j \Delta_j u \widetilde{\Delta}_j v$$

where $\widetilde{\Delta}_j = \Delta_{j-1} + \Delta_j + \Delta_{j+1}$.

Let us now define inhomogeneous Besov spaces. For $(p,r) \in [1,+\infty]^2$ and $s \in \mathbb{R}$ we define the inhomogeneous Besov space $B^s_{p,r}$ as the set of tempered distributions u such that

$$\|u\|_{B^s_{p,r}} := (2^{js}\|\Delta_j u\|_{L^p})_{\ell^r} < +\infty.$$

The homogeneous Besov space $\dot{B}^s_{p,r}$ is defined as the set of $u \in S'(\mathbb{R}^d)$ up to polynomials such that

$$\|u\|_{\dot{B}^s_{p,r}} := (2^{js}\|\dot{\Delta}_j u\|_{L^p})_{\ell^r} < +\infty.$$

Notice that the usual Sobolev spaces H^s coincide with $B^s_{2,2}$ for every $s \in \mathbb{R}$ and that the homogeneous spaces $\dot{H}^s$ coincide with $\dot{B}^s_{2,2}$.

We shall need some mixed space-time spaces. Let $T > 0$ and $\rho \geq 1$, we denote by $L^\rho_T B^s_{p,r}$ the space of distribution u such that

$$\|u\|_{L^\rho_T \dot{B}^s_{p,r}} := \|(2^{js}\|\dot{\Delta}_j u\|_{L^p})_{\ell^r}\|_{L^\rho_T} < +\infty.$$

We say that u belongs to the space $\widetilde{L}^\rho_T B^s_{p,r}$ if

$$\|u\|_{\widetilde{L}^\rho_T \dot{B}^s_{p,r}} := (2^{js}\|\dot{\Delta}_j u\|_{L^\rho_T L^p})_{\ell^r} < +\infty,$$

which appeared firstly in [19]. Through a direct application of the Minkowski inequality, the following links between these spaces is true [25]. Let $\varepsilon > 0$, then

$$L^\rho_T B^s_{p,r} \hookrightarrow \widetilde{L}^\rho_T B^s_{p,r} \hookrightarrow L^\rho_T B^{s-\varepsilon}_{p,r}, \quad \text{if } r \geq \rho,$$

$$L^\rho_T B^{s+\varepsilon}_{p,r} \hookrightarrow \widetilde{L}^\rho_T B^s_{p,r} \hookrightarrow L^\rho_T B^s_{p,r}, \quad \text{if } \rho \geq r.$$

Lemma 2.1. [18] *Let $\mathcal{B}$ be a ball and $\mathcal{C}$ be a ring of $\mathbb{R}^3$. A constant C exists so that for any positive real number λ, any nonnegative integer k, any smooth homogeneous function σ of degree m, and any couple of read number (a,b) with $b \geq a \geq 1$, there hold*

$$\begin{aligned}
&\operatorname{supp}\hat{u} \subset \lambda\mathcal{B} \Rightarrow \sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^b} \leq C^{k+1}\lambda^{k+3(\frac{1}{a}-\frac{1}{b})}\|u\|_{L^a},\\
&\operatorname{supp}\hat{u} \subset \lambda\mathcal{C} \Rightarrow C^{-1-k}\lambda^k\|u\|_{L^a} \leq \sup_{|\alpha|=k} \|\partial^\alpha u\|_{L^a} \leq C^{1+k}\lambda^k\|u\|_{L^a},\\
&\operatorname{supp}\hat{u} \subset \lambda\mathcal{C} \Rightarrow \|\sigma(D)u\|_{L^b} \leq C_{\sigma,m}\lambda^{m+3(\frac{1}{a}-\frac{1}{b})}\|u\|_{L^a}.
\end{aligned}$$

Lemma 2.2. [21] *Let $r \in [1,\infty]$, $u \in \dot{B}^s_{2,r}(\mathbb{R}^3)$, and $v \in \dot{B}^{5/2}_{2,1}(\mathbb{R}^3)$ with $\operatorname{div} v = 0$. Then there hold the following:*

(i) If $-\frac{5}{2} < s < \frac{5}{2}$ (or $s = \frac{5}{2}$ with $r = 1$),

$$\|[\Delta_q, v\cdot\nabla]u\|_{L^2} \lesssim c_{q,r}2^{-sq}\|v\|_{\dot{B}^{5/2}_{2,1}}\|u\|_{\dot{B}^s_{2,r}}. \tag{2.3}$$

(ii) If $s > -\frac{5}{2}$ and $v \in \dot{B}^s_{2,r}(\mathbb{R}^3)$, $\nabla u \in L^\infty(\mathbb{R}^3)$

$$\|[\Delta_q, v\cdot\nabla]u\|_{L^2} \lesssim c_{q,r}2^{-sq}\left(\|v\|_{\dot{B}^{5/2}_{2,1}}\|u\|_{\dot{B}^s_{2,r}} + \|v\|_{\dot{B}^s_{2,r}}\|\nabla u\|_{L^\infty}\right). \tag{2.4}$$

(iii) If $s > -1$,

$$\|[\Delta_q, v\cdot\nabla]v\|_{L^2} \lesssim c_{q,r}2^{-sq}\|\nabla v\|_{L^\infty}\|v\|_{\dot{B}^s_{2,r}}. \tag{2.5}$$

Lemma 2.3. [21] *Let v be a divergence-free vector field with $\nabla v \in L^1([0,T];\dot{B}^{3/2}_{2,1})$. For $s \in (-\frac{5}{2}, \frac{5}{2}]$, given $f_0 \in \dot{B}^s_{2,1}$, $F \in L^1([0,T];\dot{B}^s_{2,1})$, the transport equation*

$$\begin{cases}\partial_t f + v\cdot\nabla f = F \quad \text{in } \mathbb{R}^+\times\mathbb{R}^3,\\ f|_{t=0} = f_0,\end{cases} \tag{2.6}$$

has a unique solution $f \in C([0,T];\dot{B}^s_{2,1})$. Moreover, there holds for all $t \in [0,T]$

$$\begin{aligned}\|f\|_{\tilde{L}^\infty_T(\dot{B}^s_{2,1})} \leq \|f_0\|_{\dot{B}^s_{2,1}} &+ C\int_0^t \|f(\tau)\|_{\dot{B}^s_{2,1}}\|\nabla v(\tau)\|_{\dot{B}^{3/2}_{2,1}}\,d\tau\\ &+ C\|F\|_{L^1_t(\dot{B}^s_{2,1})}.\end{aligned} \tag{2.7}$$

If $s \in (0, \frac{5}{2}]$, there also holds

$$\begin{aligned}\|f\|_{\tilde{L}^\infty_t(B^s_{2,1})} \leq \|f_0\|_{B^s_{2,1}} &+ C\int_0^t \|f(t')\|_{B^s_{2,1}}\|\nabla v(t')\|_{\dot{B}^{3/2}_{2,1}}\,dt'\\ &+ C\|F\|_{L^1_t(B^s_{2,1})}.\end{aligned} \tag{2.8}$$

Lemma 2.4. [17] *Let $1 \leq p \leq p_1 \leq \infty$, $1 \leq r \leq \infty$ and $p' := (1-1/p)^{-1}$. Assume that*

$$\sigma > -N\min\Big(\frac{1}{p_1}, \frac{1}{p'}\Big) \quad \text{or} \quad \sigma > -1 - N\min\Big(\frac{1}{p_1}, \frac{1}{p'}\Big) \quad \text{if} \quad \operatorname{div} v = 0.$$

Then for the transport equation:

$$\begin{cases}\partial_t f + v\cdot\nabla f = g,\\ f|_{t=0} = f_0,\end{cases}$$

there exists a constant C depending only on N, p, p_1, r and σ, such that the following estimates hold true:

$$\|f\|_{\tilde{L}^\infty_t(B^\sigma_{p,r})} \le \Big(\|f_0\|_{B^\sigma_{p,r}} + \int_0^t e^{-C\int_0^\tau Z(\tau')\,d\tau'}\|g(\tau)\|_{B^\sigma_{p,r}}\,d\tau\Big)e^{C\int_0^t Z(\tau)\,d\tau}, \tag{2.9}$$

with

$$\begin{cases} Z(t) = \|\nabla v(t)\|_{B^{\frac{N}{p_1}}_{p_1,\infty}\cap L^\infty} & \text{if} \quad \sigma < 1+\frac{N}{p_1}, \\ Z(t) = \|\nabla v(t)\|_{B^{\sigma-1}_{p_1,1}} & \text{if} \quad \sigma > 1+\frac{N}{p_1} \quad \text{or} \quad \Big\{\sigma = 1+\frac{N}{p_1} \quad \text{and} \quad r=1\Big\}. \end{cases}$$

If $f = v$ then for all $\sigma > 0$ ($\sigma > -1$ if $\operatorname{div} v = 0$) estimates (2.7) hold with

$$Z(t) = \|\nabla v(t)\|_{L^\infty}.$$

Lemma 2.5. [21] *Let $s \in (-\frac{3}{2}, 2)$, $\vec{F} = (F_1, F_2, F_3) \in L^1_T(\dot{B}^s_{2,1})$, $a \in \tilde{L}^\infty_T(\dot{H}^2)$ with $\underline{a} := \inf_{(t,x)\in[0,T]\times\mathbb{R}^3}(1+a(t,x)) > 0$, and $\Pi \in \tilde{L}^1_T(\dot{H}^{s+1/2})$, which solves*

$$\operatorname{div}\big((1+a)\nabla\Pi\big) = \operatorname{div}\vec{F}. \tag{2.10}$$

Then there holds

$$\underline{a}\|\nabla\Pi\|_{L^1_T(\dot{B}^s_{2,1})} \lesssim \|\vec{F}\|_{L^1_T(\dot{B}^s_{2,1})} + \|a\|_{\tilde{L}^\infty_T(\dot{H}^2)}\|\nabla\Pi\|_{\tilde{L}^1_T(\dot{H}^{s-1/2})}. \tag{2.11}$$

Lemma 2.6. *For $s \in (-\frac{3}{2}, 1)$, $r = 1$ or 2. Let $u_0 \in \dot{B}^s_{2,r}$, $B_0 \in \dot{B}^s_{2,r}$ and $v \in L^1_T(\dot{B}^{5/2}_{2,1})$, $w \in L^1_T(\dot{B}^{5/2}_{2,1})$ be two divergence-free vector field. Letting $f \in \tilde{L}^1_T(\dot{B}^s_{2,r})$ and $a \in L^\infty_T(\dot{H}^2)\cap L^\infty_T(\dot{H}^{s+3/2})$ with $1+a \ge \underline{c} > 0$, we assume that $u \in L^\infty_T(\dot{B}^s_{2,r})\cap L^1_T(\dot{B}^{s+\frac{3}{2}}_{2,r})$, $B \in L^\infty_T(\dot{B}^s_{2,r}) \cap L^1_T(\dot{B}^{s+\frac{3}{2}}_{2,r})$ and $\Pi \in L^1_T(\dot{H}^1)$ solve*

$$\begin{cases} \partial_t u - w\cdot\nabla B + v\cdot\nabla u - \Delta u + \nabla\Pi = f + a(\Delta u - \nabla\Pi), \quad \mathbb{R}^+\times\mathbb{R}^3, \\ \partial_t B + v\cdot\nabla B - w\cdot\nabla u - \Delta B = g, \\ \operatorname{div} u = \operatorname{div} B = 0, \\ (u,B)|_{t=0} = (u_0, B_0). \end{cases} \tag{2.12}$$

Then there holds

$$\begin{aligned} &\|(u,B)\|_{\tilde{L}^\infty_T(\dot{B}^s_{2,r})} + \|(u,B)\|_{\tilde{L}^1_T(\dot{B}^{s+2}_{2,r})} \\ &\lesssim \exp\left(C\int_0^T \|v(t)\|_{\dot{B}^{5/2}_{2,1}} + \|w(t)\|_{\dot{B}^{5/2}_{2,1}}\,dt\right)\Big\{\|(u_0,B_0)\|_{\dot{B}^s_{2,r}} \\ &\qquad + \|(f,g)\|_{\tilde{L}^1_T(\dot{B}^s_{2,r})} + \|a\|_{L^\infty_T(\dot{H}^{s+3/2})}\|\nabla\Pi\|_{L^1_T(L^2)} \\ &\qquad + \|a\|_{L^\infty_T(\dot{H}^2)}\|u\|_{L^1_T(\dot{B}^{s+3/2}_{2,r})}\Big\} \end{aligned} \tag{2.13}$$

Proof. We apply the operator Δ_q to (2.12); then a standard commutator process gives

$$\begin{aligned} &\partial_t\Delta_q u - (w\cdot\nabla)\Delta_q B + (v\cdot\nabla)\Delta_q u - \operatorname{div}((1+a)\nabla\Delta_q u) \\ &\qquad + \Delta_q\nabla((1+a)\Pi) = -[w,\Delta_q]\cdot\nabla B + [v,\Delta_q]\cdot\nabla u - (\nabla a\cdot\nabla)\Delta_q u \\ &\qquad\qquad + \Delta_q(\nabla a\cdot\Pi) + R_q + \Delta_q f, \end{aligned} \tag{2.14}$$

$$\begin{aligned} &\partial_t\Delta_q B + v\cdot\nabla\Delta_q B - w\cdot\nabla\Delta_q u - \Delta\Delta_q B \\ &\qquad = \Delta_q g + [v,\Delta_q]\nabla B - [w,\Delta_q]\nabla u, \end{aligned} \tag{2.15}$$

with

$$R_q = \mathrm{div}[\Delta_q, a]\nabla u + [\nabla a\cdot\nabla, \Delta_q]u.$$

Thanks to the fact that $\mathrm{div}u = \mathrm{div}v = \mathrm{div}w = \mathrm{div}B = 0$ and $1 + a \geq \underline{c}$, we get by taking the L^2 inner product of (2.14), (2.15) with $\Delta_q u$ and $\Delta_q B$ separately that

$$\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\big(\|\Delta_q u(t)\|_{L^2}^2 + \|\Delta_q B\|_{L^2}^2\big) + c\,2^{2q}\Big(\|\Delta_q u\|_{L^2}^2 + \|\Delta_q B\|_{L^2}^2\Big)\\
&\lesssim \big(\|\Delta_q u\|_{L^2} + \|\Delta_q B\|_{L^2}\big)\big(\|[w,\Delta_q]\nabla B\|_{L^2} + \|[v,\Delta_q]\nabla u\|_{L^2}\\
&\quad + \|[v,\Delta_q]\nabla B\|_{L^2} + \|[w,\Delta_q]\nabla u\|_{L^2} + \|R_q\|_{L^2} + \|\Delta_q(\Pi\cdot\nabla a)\|_{L^2}\\
&\quad + \|\Delta_q f\|_{L^2} + \|\Delta_q g\|_{L^2}\big) + \|\Delta a\|_{L^2}\|\Delta_q u\|_{L^4}^2.
\end{aligned}$$

Thanks to Lemma 2.1, we have

$$\|\Delta a\|_{L^2}\|\Delta_q u\|_{L^4}^2 \lesssim 2^{(3/2)q}\|\Delta a\|_{L^2}\|\Delta_q u\|_2^2.$$

It follows from the product law that for all $s \in (-\frac{3}{2}, 1)$ that

$$\|\Pi\nabla a(t)\|_{\dot B^s_{2,1}} \lesssim \|\nabla a(t)\|_{\dot H^{s+\frac12}}\|\Pi(t)\|_{\dot H^1}.$$

Similar to the proof in Proposition 3.6 in [21], we have

$$\|R_q\|_{L^2} \lesssim d_q(t)2^{-qs}\|a(t)\|_{\dot H^2}\|u(t)\|_{\dot H^{s+3/2}}.$$

Using Lemma 2.2, we have the following four inequalities:

$$\begin{aligned}
\|[w,\Delta_q]\nabla B\|_{L^2} &\lesssim c_{q,r}2^{-qs}\|w\|_{\dot B^{5/2}_{2,1}}\|B\|_{\dot B^s_{2,r}},\\
\|[v,\Delta_q]\nabla u\|_{L^2} &\lesssim c_{q,r}2^{-qs}\|v\|_{\dot B^{5/2}_{2,1}}\|u\|_{\dot B^s_{2,r}},\\
\|[v,\Delta_q]\nabla B\|_{L^2} &\lesssim c_{q,r}2^{-qs}\|v\|_{\dot B^{5/2}_{2,1}}\|B\|_{\dot B^s_{2,r}},\\
\|[w,\Delta_q]\nabla u\|_{L^2} &\lesssim c_{q,r}2^{-qs}\|w\|_{\dot B^{5/2}_{2,1}}\|u\|_{\dot B^s_{2,r}}.
\end{aligned}$$

Combining the above four estimates, we arrive at

$$\begin{aligned}
&\|(u,B)\|_{\tilde L^\infty_T(\dot B^s_{2,r})} + \|(u,B)\|_{\tilde L^1_T(\dot B^{s+2}_{2,r})}\\
&\lesssim \|(u_0,B_0)\|_{\dot B^s_{2,r}} + \|(f,g)\|_{\tilde L^1_T(\dot B^s_{2,r})} + \|a\|_{L^\infty_T(\dot H^{s+3/2})}\|\nabla\Pi\|_{L^1_T(L^2)}\\
&\quad + \|a\|_{L^\infty_T(\dot H^2)}\|u\|_{L^1_T(\dot B^{s+3/2}_{2,r})}\\
&\quad + \int_0^T \big(\|w\|_{\dot B^{5/2}_{2,1}} + \|v\|_{\dot B^{5/2}_{2,1}}\big)\big(\|u\|_{\dot B^s_{2,r}} + \|B\|_{\dot B^s_{2,r}}\big)\,d\tau.
\end{aligned}$$

Through Gronwall's inequality, we complete the proof. □

Remark 2.7. It is easy to observe from the following

$$\|\Pi\nabla a\|_{\tilde L^1_T(\dot B^s_{2,r})} \lesssim \|\nabla a\|_{\tilde L^\infty_T\dot H^1}\|\Pi\|_{\tilde L^1_T(\dot H^{s+1/2})} \quad \text{for all } s\in(-\frac{3}{2},1),$$

that

$$\begin{aligned}
&\|(u,B)\|_{\tilde L^\infty_T(\dot B^s_{2,r})} + \|(u,B)\|_{\tilde L^1_T(\dot B^{s+2}_{2,r})}\\
&\lesssim \exp\left(C\int_0^T \|(v(t),B(t))\|_{\dot B^{5/2}_{2,1}}\,dt\right)\Big\{\|(u_0,B_0)\|_{\dot B^s_{2,r}}\\
&\quad + \|(f,g)\|_{\tilde L^1_T(\dot B^s_{2,r})} + \|a\|_{\tilde L^\infty_T(\dot H^2)}\big(\|\Pi\|_{\tilde L^1_T(\dot H^{s+1/2})} + \|u\|_{L^1_T(\dot B^{s+3/2}_{2,r})}\big)\Big\}.
\end{aligned} \tag{2.16}$$

Remark 2.8. Note that div$u = 0$, taking div to the first equation of (2.12), we obtain

$$\operatorname{div}\big((1+a)\nabla\Pi\big) = \operatorname{div}\big(f + a\Delta u + w\cdot\nabla B - v\cdot\nabla u\big),$$

then is follows from Lemma 2.5 that for $s \in (-\frac{3}{2}, \frac{3}{2})$,

$$\begin{aligned}\|\nabla\Pi\|_{L^1_T(\dot B^s_{2,1})} \lesssim{}& \|v\|_{L^\infty_T(\dot H^s)}\|u\|_{L^1_T(\dot B^{5/2}_{2,1})} + \|w\|_{L^\infty_T(\dot H^s)}\|B\|_{L^1_T(\dot B^{5/2}_{2,1})}\\ &+ \|a\|_{\tilde L^\infty_T(\dot B^{3/2}_{2,1})}\|u\|_{\tilde L^1_T\dot H^{s+2}} + \|a\|_{\tilde L^\infty_T(\dot H^2)}\|\nabla\Pi\|_{\tilde L^1_T(\dot H^{s-1/2})}\\ &+ \|f\|_{\tilde L^1_T(\dot B^s_{2,1})}.\end{aligned} \tag{2.17}$$

Remark 2.9. If the parameter $s = 1$, then we have the following estimation

$$\begin{aligned}&\|u\|_{\tilde L^\infty_T(\dot B^1_{2,1})} + \|u\|_{\tilde L^1_T(\dot B^3_{2,1})} + \|B\|_{\tilde L^\infty_T(\dot B^1_{2,1})} + \|B\|_{\tilde L^1_T(\dot B^3_{2,1})}\\ &\lesssim \exp\left(C\int_0^T \|v\|_{\dot B^{5/2}_{2,1}} + \|w\|_{\dot B^{5/2}_{2,1}}\,dt\right)\Big(\|u_0\|_{\dot B^1_{2,1}} + \|B_0\|_{\dot B^1_{2,1}}\\ &\quad + \|f\|_{\tilde L^1_T(\dot B^1_{2,1})} + \|g\|_{\tilde L^1_T(\dot B^1_{2,1})} + \|a\|_{L^\infty_T(\dot B^2_{2,1})}\|\Pi\|_{L^1_T(\dot B^{3/2}_{2,1})}\\ &\quad + \|a\|_{L^\infty_T(\dot B^2_{2,1})}\|u\|_{L^1_T(\dot B^{5/2}_{2,1})}\Big).\end{aligned} \tag{2.18}$$

The proof is similar, so we only give the main estimations in the appendix.

3. Stability of Global Solutions with Densities Close to 1

The aim of this section is to investigate the global stability of the given solution of (1.2) with the initial density of which is close to 1, namely Theorem 1.1. Next, we give the detailed statement.

Proof. To deal with the global well-posedness of (1.2) with initial data (a_0, u_0, B_0) given by the theorem, we need some global-in-time control of the reference solution $(\bar a, \bar u, \bar B)$. In what follows, we shall always denote $\bar\rho := \frac{1}{1+\bar a}$. Then we get by a standard energy estimate to (1.3) that

$$\begin{aligned}&\frac{1}{2}\frac{d}{dt}\|\sqrt{\bar\rho}\bar u(t)\|^2_{L^2} + \|\nabla\bar u(t)\|^2_{L^2} - \int_{\mathbb{R}^3}(\bar B\cdot\nabla)\bar B\,\bar u\,dx = 0,\\ &\frac{1}{2}\frac{d}{dt}\|\bar B(t)\|^2_{L^2} + \|\nabla\bar B(t)\|^2_{L^2} - \int_{\mathbb{R}^3}(\bar B\cdot\nabla)\bar u\,\bar B\,dx = 0\end{aligned}$$

Combining the above two energy equality, we have

$$\frac{1}{2}\frac{d}{dt}\left(\|\sqrt{\bar\rho}\bar u(t)\|^2_{L^2} + \|\bar B(t)\|^2_{L^2}\right) + \|\nabla\bar u(t)\|^2_{L^2} + \|\nabla\bar B(t)\|^2_{L^2} = 0, \quad \text{for } t>0. \tag{3.1}$$

After integration, we have

$$\frac{1}{2}\|(\sqrt{\bar\rho}\bar u(t), \bar B(t))\|^2_{L^2} + \int_0^t \|(\nabla\bar u(\tau), \nabla\bar B(\tau))\|^2_{L^2}\,d\tau = \frac{1}{2}\|(\sqrt{\bar\rho_0}\bar u_0, \bar B_0)\|^2_{L^2}. \tag{3.2}$$

The transport equation in (1.3) gives

$$\|\bar\rho(t) - 1\|_{L^p} = \|\bar\rho_0 - 1\|_{L^p}, \quad \text{for all } p\in[1,\infty]. \tag{3.3}$$

Using the interpolation inequality

$$\|\bar{u}\|_{\dot{B}^{1/2}_{2,1}} \lesssim \|\bar{u}\|^{1/2}_{L^2}\|\nabla\bar{u}\|^{1/2}_{L^2},$$
$$\|\bar{B}\|_{\dot{B}^{1/2}_{2,1}} \lesssim \|\bar{B}\|^{1/2}_{L^2}\|\nabla\bar{B}\|^{1/2}_{L^2},$$

and from (3.2), we deduce that

$$\begin{aligned}\int_0^t \|\bar{u}(\tau)\|^4_{\dot{B}^{1/2}_{2,1}}\,d\tau &\lesssim \int_0^t \|\bar{u}(\tau)\|^2_{L^2}\|\nabla\bar{u}(\tau)\|^2_{L^2}\,d\tau\\ &\lesssim \|\bar{u}_0\|^4_{L^2} + \|\bar{B}_0\|^4_{L^2}, \quad \text{for } t>0,\\ \int_0^t \|\bar{B}(\tau)\|^4_{\dot{B}^{1/2}_{2,1}}\,d\tau &\lesssim \int_0^t \|\bar{B}(\tau)\|^2_{L^2}\|\nabla\bar{B}(\tau)\|^2_{L^2}\,d\tau\\ &\lesssim \|\bar{u}_0\|^4_{L^2} + \|\bar{B}_0\|^4_{L^2}, \quad \text{for } t>0,\end{aligned}$$

Hence, for any $\epsilon>0$, there exists $T_0=T_0(\epsilon)>0$ such that

$$\begin{aligned}&\|\bar{u}(T_0)\|_{\dot{B}^{1/2}_{2,1}} < \epsilon,\\ &\|\bar{B}(T_0)\|_{\dot{B}^{1/2}_{2,1}} < \epsilon.\end{aligned} \tag{3.4}$$

On the other hand, applying Lemma 2.3 to the transport equation in (1.2) gives

$$\|\bar{a}(t)\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})} \le \|\bar{a}_0\|_{\dot{B}^{3/2}_{2,1}} + C\int_0^t \|\bar{a}(\tau)\|_{\dot{B}^{3/2}_{2,1}}\|\nabla\bar{u}(\tau)\|_{\dot{B}^{3/2}_{2,1}}\,d\tau,$$

for any $t\ge 0$. Then applying Gronwall's inequality yields

$$\|\bar{a}\|_{\tilde{L}^\infty_{T_0}(\dot{B}^{3/2}_{2,1})} \le \|\bar{a}_0\|_{\dot{B}^{3/2}_{2,1}}\exp\left\{C\int_0^{T_0}\|\nabla\bar{u}(\tau)\|_{\dot{B}^{3/2}_{2,1}}\,d\tau\right\}.$$

Following the same idea, it is easy to obtain that for $t\ge T_0$

$$\|\bar{a}\|_{\tilde{L}^\infty([T_0,t];\dot{B}^{3/2}_{2,1})} \le \|\bar{a}(T_0)\|_{\dot{B}^{3/2}_{2,1}} + C\|\bar{a}\|_{L^\infty([T_0,t];\dot{B}^{3/2}_{2,1})}\|\nabla\bar{u}\|_{L^1([T_0,t];\dot{B}^{3/2}_{2,1})}. \tag{3.5}$$

Note that for $\bar{a}$ small, we can rewrite the momentum equation and magnetic field equation in (1.2) as

$$\begin{aligned}&\partial_t\bar{u} + (\bar{u}\cdot\nabla)\bar{u} - \Delta\bar{u} + \nabla\bar{\Pi} = \bar{a}(\Delta\bar{u}-\nabla\bar{\Pi}) + (\bar{B}\cdot\nabla)\bar{B} + \bar{a}(\bar{B}\cdot\nabla)\bar{B},\\ &\partial_t\bar{B} + (\bar{u}\cdot\nabla)\bar{B} - \Delta\bar{B} = (\bar{B}\cdot\nabla)\bar{u},\end{aligned}$$

from which we get by using a standard energy estimate about heat equation that

$$\begin{aligned}&\|(\bar{u},\bar{B})\|_{\tilde{L}^\infty([T_0,t];\dot{B}^{1/2}_{2,1})} + \|(\bar{u},\bar{B})\|_{L^1([T_0,t];\dot{B}^{5/2}_{2,1})} + \|\nabla\bar{\Pi}\|_{L^1([T_0,t];\dot{B}^{1/2}_{2,1})}\\ \lesssim&\|(\bar{u}(T_0),\bar{B}(T_0))\|_{\dot{B}^{1/2}_{2,1}} + \|\bar{a}\bar{B}\nabla\bar{B}\|_{L^1([T_0,t];\dot{B}^{1/2}_{2,1})} + \|\bar{a}(\Delta\bar{u}-\nabla\bar{\Pi})\|_{L^1([T_0,t];\dot{B}^{1/2}_{2,1})}\\ &+\left\|\big((\bar{u}\cdot\nabla)\bar{u},(\bar{B}\cdot\nabla)\bar{B},(\bar{u}\cdot\nabla)\bar{B},(\bar{B}\cdot\nabla)\bar{u}\big)\right\|_{L^1([T_0,t];\dot{B}^{1/2}_{2,1})}.\end{aligned} \tag{3.6}$$

For any $t\ge T_0$, we denote

$$\begin{aligned}\bar{Z}(t) :=& \|\bar{a}\|_{\tilde{L}^\infty([T_0,t];\dot{B}^{3/2}_{2,1})} + \|(\bar{u},\bar{B})\|_{\tilde{L}^\infty([T_0,t];\dot{B}^{1/2}_{2,1})} + \|\nabla\Pi\|_{L^1([T_0,t];\dot{B}^{1/2}_{2,1})}\\ &+\|(\bar{u},\bar{B})\|_{L^1([T_0,t];\dot{B}^{5/2}_{2,1})}.\end{aligned}$$

From the product law, we get

$$
\begin{aligned}
&\|(\bar u\cdot\nabla)\bar u\|_{L^1([T_0,t];\dot B^{1/2}_{2,1})}\lesssim \|\bar u\|_{L^\infty([T_0,t];\dot B^{1/2}_{2,1})}\|\nabla\bar u\|_{L^1([T_0,t];\dot B^{3/2}_{2,1})},\\
&\|(\bar B\cdot\nabla)\bar B\|_{L^1([T_0,t];\dot B^{1/2}_{2,1})}\lesssim \|\bar B\|_{L^\infty([T_0,t];\dot B^{1/2}_{2,1})}\|\nabla\bar B\|_{L^1([T_0,t];\dot B^{3/2}_{2,1})},\\
&\|\bar a\,(\bar B\cdot\nabla)\bar B\|_{L^1([T_0,t];\dot B^{1/2}_{2,1})}\\
&\qquad\lesssim \|\bar a\|_{L^\infty([T_0,t];\dot B^{3/2}_{2,1})}\|\bar B\|_{L^\infty([T_0,t];\dot B^{1/2}_{2,1})}\|\nabla\bar B\|_{L^1([T_0,t];\dot B^{3/2}_{2,1})},\\
&\|\bar a(\Delta\bar u-\nabla\bar\Pi)\|_{L^1([T_0,t];\dot B^{1/2}_{2,1})}\\
&\qquad\lesssim \|\bar a\|_{L^\infty([T_0,t];\dot B^{3/2}_{2,1})}\Big(\|\bar u\|_{L^1([T_0,t];\dot B^{5/2}_{2,1})}+\|\nabla\bar\Pi\|_{L^1([T_0,t];\dot B^{1/2}_{2,1})}\Big)\\
&\|(\bar u\cdot\nabla)\bar B\|_{L^1([T_0,t];\dot B^{1/2}_{2,1})}\lesssim \|\bar u\|_{L^\infty([T_0,t];\dot B^{1/2}_{2,1})}\|\nabla\bar B\|_{L^1([T_0,t];\dot B^{3/2}_{2,1})},\\
&\|(\bar B\cdot\nabla)\bar u\|_{L^1([T_0,t];\dot B^{1/2}_{2,1})}\lesssim \|\bar B\|_{L^\infty([T_0,t];\dot B^{1/2}_{2,1})}\|\nabla\bar u\|_{L^1([T_0,t];\dot B^{3/2}_{2,1})},
\end{aligned}
$$

Then we get by summing up (3.5), (3.6) and the above estimates that

$$
\bar Z(t)\le \|\bar a(T_0)\|_{\dot B^{3/2}_{2,1}}+\|(\bar u(T_0),\bar B(T_0))\|_{\dot B^{1/2}_{2,1}}+C\,\bar Z(t)^2+C\,\bar Z(t)^3.
$$

Let

$$
\bar T:=\sup_{t>T_0}\Big\{t\,:\,\bar Z(t)\le 2\,\Big(\|\bar a(T_0)\|_{\dot B^{3/2}_{2,1}}+\|(\bar u(T_0),\bar B(T_0))\|_{\dot B^{1/2}_{2,1}}\Big)\Big\} \tag{3.7}
$$

With no loss of generality, we can assume that

$$
\|\bar a(T_0)\|_{\dot B^{3/2}_{2,1}}+\|\bar u(T_0)\|_{\dot B^{1/2}_{2,1}}+\|\bar B(T_0)\|_{\dot B^{1/2}_{2,1}}\le 1. \tag{3.8}
$$

Then, if $\bar T<\infty$, for $T_0<t<\bar T$, we have

$$
\bar Z(t)\le\Big(\|\bar a(T_0)\|_{\dot B^{3/2}_{2,1}}+\|(\bar u(T_0),\bar B(T_0))\|_{\dot B^{1/2}_{2,1}}\Big)\big(1+6C\bar Z(t)\big)
$$

Taking ϵ in (3.4) small enough so that $\epsilon\le\frac{1}{108C}$, then if c_1 in Theorem 1.1 is so small that $c_1\le\frac{1}{108C}$, we have

$$
\bar Z(t)\le\frac{3}{2}\Big(\|\bar a(T_0)\|_{\dot B^{3/2}_{2,1}}+\|(\bar u(T_0),\bar B(T_0))\|_{\dot B^{1/2}_{2,1}}\Big)\quad\text{for } T_0\le t\le\bar T. \tag{3.9}
$$

This contradicts with the definition of (3.7), and therefore $\bar T=\infty$. Moreover, there hold

$$
\begin{aligned}
&\|\bar a\|_{\tilde L^\infty(\mathbb{R}^+;\dot B^{3/2}_{2,1})}\le 2\,\|\bar a_0\|_{\dot B^{3/2}_{2,1}}\exp\left\{C\int_0^{T_0}\|\nabla\bar u(\tau)\|_{\dot B^{3/2}_{2,1}}\,d\tau\right\}+4\epsilon,\\
&\|(\bar u,\bar B)\|_{\tilde L^\infty(\mathbb{R}^+;\dot B^{1/2}_{2,1})}+\|(\bar u,\bar B)\|_{L^1(\mathbb{R}^+;\dot B^{5/2}_{2,1})}+\|\nabla\bar\Pi\|_{L^1(\mathbb{R}^+;\dot B^{1/2}_{2,1})}\le C.
\end{aligned} \tag{3.10}
$$

With (3.10), we can solve the global well-posedness with initial data $(a_0,u_0,B_0)=(\bar a_0+\tilde a_0,\bar u_0+\tilde u_0,\bar B_0+\tilde B_0)$ for $(\tilde a_0,\tilde u_0,\tilde B_0)$ sufficiently small. Let $\tilde u:=u-\bar u$,

$\tilde{B} := B - \bar{B}$, then $(a, \tilde{u}, \tilde{B})$ solves

$$
\begin{cases}
\partial_t a + (\bar{u}+\tilde{u})\nabla a = 0, \\
\partial_t \tilde{u} - \Delta\tilde{u} + \nabla\tilde{\Pi} = -(\bar{u}+\tilde{u})\nabla\tilde{u} - (\tilde{u}\cdot\nabla)\bar{u} + a\,(\Delta\tilde{u} - \nabla\tilde{\Pi}) \\
\qquad\qquad\qquad\qquad + (a-\bar{a})(\Delta\bar{u} - \nabla\bar{\Pi}) + (a-\bar{a})(\bar{B}\cdot\nabla\bar{B}) \\
\qquad\qquad\qquad\qquad + (1+a)(\bar{B}+\tilde{B})\nabla\tilde{B} + (1+a)(\tilde{B}\cdot\nabla)\bar{B}, \\
\partial_t \tilde{B} - \Delta\tilde{B} = -(\tilde{u}+\bar{u})\nabla\tilde{B} - (\tilde{u}\cdot\nabla)\bar{B} + (\tilde{B}+\bar{B})\nabla\tilde{u} + (\tilde{B}\cdot\nabla)\bar{u}, \\
\mathrm{div}\tilde{u} = \mathrm{div}\tilde{B} = 0, \\
(a, \tilde{u}, \tilde{B})|_{t=0} = (\bar{a}_0 + \tilde{a}_0, \tilde{u}_0, \tilde{B}_0).
\end{cases} \tag{3.11}
$$

Applying Lemma 2.3 to the first equation of (3.11) gives

$$
\begin{aligned}
\|a\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})} \le \|a_0\|_{\dot{B}^{3/2}_{2,1}} &+ C\int_0^t \|a(\tau)\|_{\dot{B}^{3/2}_{2,1}} \|\bar{u}(\tau)\|_{\dot{B}^{5/2}_{2,1}}\, d\tau \\
&+ C\,\|a\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})} \|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})},
\end{aligned} \tag{3.12}
$$

which along with (3.10) and the Gronwall's inequality ensures

$$
\|a\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})} \lesssim \left(\|a_0\|_{\dot{B}^{3/2}_{2,1}} + \|a\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})} \|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\right). \tag{3.13}
$$

Reformulate equation (3.11) as follows

$$
\begin{cases}
\partial_t \tilde{u} - (\bar{B}\cdot\nabla)\tilde{B} + (\bar{u}\cdot\nabla)\tilde{u} - \Delta\tilde{u} + \nabla\tilde{\Pi} = \tilde{f} + a\,(\Delta\tilde{u} - \nabla\tilde{\Pi}), \\
\partial_t \tilde{B} + (\bar{u}\cdot\nabla)\tilde{B} - (\bar{B}\cdot\nabla)\tilde{u} - \Delta\tilde{B} = \tilde{g},
\end{cases} \tag{3.14}
$$

with

$$
\begin{aligned}
\tilde{f} &= -(\tilde{u}\cdot\nabla)\tilde{u} - (\tilde{u}\cdot\nabla)\bar{u} + (a-\bar{a})(\Delta\bar{u} - \nabla\bar{\Pi}) + (a-\bar{a})(\bar{B}\cdot\nabla\bar{B}) \\
&\quad + (\tilde{B}\cdot\nabla)\tilde{B} + (\tilde{B}\cdot\nabla)\bar{B} + a\,\tilde{B}\cdot\nabla\tilde{B} + a\,\bar{B}\cdot\nabla\tilde{B} + a\,\tilde{B}\cdot\nabla\bar{B} \\
\tilde{g} &= -(\tilde{u}\cdot\nabla)\tilde{B} - (\tilde{u}\cdot\nabla)\bar{B} + (\tilde{B}\cdot\nabla)\tilde{u} + (\tilde{B}\cdot\nabla)\bar{u}.
\end{aligned}
$$

Using standard estimates, we easily obtain that

$$
\begin{aligned}
&\|(\tilde{u},\tilde{B})\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|(\tilde{u},\tilde{B})\|_{\tilde{L}^1_t(\dot{B}^{5/2}_{2,1})} + \|\nabla\tilde{\Pi}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \\
&\lesssim \|(\tilde{u}_0,\tilde{B}_0)\|_{\dot{B}^{1/2}_{2,1}} + \int_0^t \|(\tilde{u},\tilde{B})\|_{\dot{B}^{1/2}_{2,1}} \|(\tilde{u},\tilde{B})\|_{\dot{B}^{5/2}_{2,1}}\, d\tau \\
&\quad + \int_0^t \|(\tilde{u},\tilde{B})\|_{\dot{B}^{1/2}_{2,1}} \|(\bar{u},\bar{B})\|_{\dot{B}^{5/2}_{2,1}}\, d\tau + \int_0^t \|(a-\bar{a})(\Delta\bar{u} - \nabla\bar{\Pi})\|_{\dot{B}^{1/2}_{2,1}}\, d\tau \\
&\quad + \int_0^t \|a\,(\Delta\tilde{u} - \nabla\tilde{\Pi})\|_{\dot{B}^{1/2}_{2,1}}\, d\tau + \int_0^t \|(a-\bar{a})\bar{B}\cdot\nabla\bar{B}\|_{\dot{B}^{1/2}_{2,1}}\, d\tau \\
&\quad + \int_0^t \|a\|_{\dot{B}^{3/2}_{2,1}} \|\tilde{B}\|_{\dot{B}^{1/2}_{2,1}} \|\nabla\tilde{B}\|_{\dot{B}^{3/2}_{2,1}}\, d\tau + \int_0^t \|a\|_{\dot{B}^{3/2}_{2,1}} \|\bar{B}\|_{\dot{B}^{1/2}_{2,1}} \|\nabla\tilde{B}\|_{\dot{B}^{3/2}_{2,1}}\, d\tau \\
&\quad + \int_0^t \|a\|_{\dot{B}^{3/2}_{2,1}} \|\tilde{B}\|_{\dot{B}^{1/2}_{2,1}} \|\nabla\bar{B}\|_{\dot{B}^{3/2}_{2,1}}\, d\tau
\end{aligned}
$$

Then Gronwall's inequality ensures that

$$
\begin{aligned}
&\|(\tilde{u},\tilde{B})\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})}+\|(\tilde{u},\tilde{B})\|_{L^1_t(\dot{B}^{5/2}_{2,1})}+\|\nabla\tilde{\Pi}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}\\
\lesssim&\|(\tilde{u}_0,\tilde{B}_0)\|_{\dot{B}^{1/2}_{2,1}}+\|(\tilde{u},\tilde{B})\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})}\|(\tilde{u},\tilde{B})\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\\
&+\|a\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})}\Big(\|(\tilde{u},\tilde{B})\|_{L^1_t(\dot{B}^{5/2}_{2,1})}+\|\tilde{B}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})}+\|\nabla\tilde{\Pi}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}\Big)\\
&+\|\bar{a}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}+\|a\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})}\|\tilde{B}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})}\|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\\
&+\int_0^t\|a\|_{\dot{B}^{3/2}_{2,1}}\Big(\|\bar{u}\|_{\dot{B}^{5/2}_{2,1}}+\|\nabla\bar{\Pi}\|_{\dot{B}^{1/2}_{2,1}}+\|\bar{B}\|_{\dot{B}^{1/2}_{2,1}}\|\nabla\bar{B}\|_{\dot{B}^{3/2}_{2,1}}\Big)\,d\tau
\end{aligned}
\tag{3.15}
$$

Let

$$
\tilde{Z}(t):=\|a\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})}+\|(\tilde{u},\tilde{B})\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})}+\|(\tilde{u},\tilde{B})\|_{L^1_t(\dot{B}^{5/2}_{2,1})}+\|\nabla\tilde{\Pi}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}.
$$

Summing up (3.13) and (3.15), and then applying Gronwall's inequality, we obtain

$$
\tilde{Z}(t)\le C\,\Big(\|(\tilde{u}_0,\tilde{B}_0)\|_{\dot{B}^{1/2}_{2,1}}+\|\tilde{a}_0\|_{\dot{B}^{3/2}_{2,1}}+\|\bar{a}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}+\tilde{Z}(t)^2+\tilde{Z}(t)^3\Big).
\tag{3.16}
$$

Using similar arguments used in proving (3.9), we can show that if $\|(\tilde{u}_0,\tilde{B}_0)\|_{\dot{B}^{1/2}_{2,1}}+\|\tilde{a}_0\|_{\dot{B}^{3/2}_{2,1}}+2c_1+4\epsilon\le\frac{1}{12\,C^2}$, there holds

$$
\tilde{Z}(t)\le 2\,C\,\Big(\|(\tilde{u}_0,\tilde{B}_0)\|_{\dot{B}^{1/2}_{2,1}}+\|\tilde{a}_0\|_{\dot{B}^{3/2}_{2,1}}+\|\bar{a}\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})}\Big)\quad\text{for all } t>0.
\tag{3.17}
$$

With (3.17), we can prove the propagation of regularity for smoother initial data. Applying Lemma 2.3, we obtain for any $t>0$,

$$
\|a\|_{\tilde{L}^\infty_t(\dot{B}^{5/2}_{2,1})}\le\|a_0\|_{\dot{B}^{5/2}_{2,1}}+C\int_0^t\|a(\tau)\|_{\dot{B}^{5/2}_{2,1}}\|\nabla u(\tau)\|_{\dot{B}^{3/2}_{2,1}}\,d\tau.
$$

This along with (3.10) and (3.17) implies

$$
\|a\|_{\tilde{L}^\infty_t(\dot{B}^{5/2}_{2,1})}\le C\,\|a_0\|_{\dot{B}^{5/2}_{2,1}}.
\tag{3.18}
$$

Noticing that $\|a(t)\|_{L^2}=\|a_0\|_{L^2}$, this along with the above inequality shows that $a\in\tilde{L}^\infty(\mathbb{R}^+;B^{5/2}_{2,1}(\mathbb{R}^3))$. Applying standard energy estimate to the second and third equation of (1.2) gives

$$
\begin{aligned}
&\|(u,B)\|_{\tilde{L}^\infty_t(\dot{B}^s_{2,1})}+\|(u,B)\|_{L^1_t(\dot{B}^{s+2}_{2,1})}+\|\nabla\Pi\|_{L^1_t(\dot{B}^s_{2,1})}\\
\le&\|(u_0,B_0)\|_{\dot{B}^s_{2,1}}+\|a\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\Big(\|u\|_{L^1_t(\dot{B}^{s+2}_{2,1})}+\|\nabla\Pi\|_{L^1_t(\dot{B}^s_{2,1})}\Big)\\
&+C\,\Big(1+\|a\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\Big)\int_0^t\|(\nabla u(\tau),\nabla B(\tau))\|_{L^\infty}\|(u(\tau),B(\tau))\|_{\dot{B}^s_{2,1}}\,d\tau\\
&+\int_0^t\|(\nabla u,\nabla B)\|_{\dot{B}^{3/2}_{2,1}}\|B\|_{\dot{B}^s_{2,1}}\,d\tau,
\end{aligned}
$$

for $s\in[0,\frac{3}{2}]$ and $t>0$, this together with (3.10) and (3.17) gives rise to

$$
\|(u,B)\|_{\tilde{L}^\infty_t(\dot{B}^s_{2,1})}+\|(u,B)\|_{L^1_t(\dot{B}^{s+2}_{2,1})}+\|\nabla\Pi\|_{L^1_t(\dot{B}^s_{2,1})}\le C\quad\text{for all } t>0.
\tag{3.19}
$$

With (3.18) and (3.19), we can easily prove by a classical argument that

$$
\begin{aligned}
&a\in C_b([0,\infty);B^{5/2}_{2,1}),\\
&u,B\in C_b([0,\infty);B^{3/2}_{2,1}).
\end{aligned}
$$

Now, the proof of Theorem 1.1 is completed. □

4. Decay in Time Estimates for the Reference Solutions

In this section, we will give the decay estimates, namely Theorem 1.2. The main ingredient of the proof will be H. Abidi, G. Gui and P. Zhang's approach in [21]. The difference is that we need to give the decay estimates for more higher order derivatives of momentum and magnetic fields, which is required for the global in time estimates proved in the next section.

In what follows, we shall always denote $\rho(t,x) := \frac{1}{1+a(t,x)}$ so that we can use both (1.2) and (1.3) just according to our convenience. In order to make our presentation clearly, we divided the proof of Theorem 1.2 into the following propositions:

Proposition 4.1. Under the same assumptions of Theorem 1.2, there exists $t_0 > 0$ and two positive constants e_1 and e_2 such that there holds

$$\begin{aligned}\frac{d}{dt}\|(\nabla u(t), \nabla B(t))\|_{L^2}^2 &+ e_1\|(\partial_t u(t), \partial_t B(t))\|_{L^2}^2 \\ &+ e_2\|(\nabla^2 u(t), \nabla^2 B(t))\|_{L^2}^2 \le 0 \quad \text{for all } t \ge t_0,\end{aligned} \tag{4.1}$$

or consequently

$$\begin{aligned}\sup_{t\ge t_0}\|(\nabla u(t), \nabla B(t))\|_{L^2}^2 &+ \int_{t_0}^{\infty} e_1\|(\partial_t u(t), \partial_t B(t))\|_{L^2}^2\, d\tau \\ &+ \int_{t_0}^{\infty} e_2\|(\nabla^2 u(t), \nabla^2 B(t))\|_{L^2}^2\, d\tau \le \|(\nabla u(t_0), \nabla B(t_0))\|_{L^2}^2.\end{aligned} \tag{4.2}$$

Proof. Taking L^2 inner product of the momentum equation of (1.3) with $\frac{1}{\rho}\Delta u$ that

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\|\nabla u(t)\|_{L^2}^2 + \|\frac{1}{\sqrt{\rho}}\Delta u(t)\|_{L^2}^2 &= \int_{\mathbb{R}^3}(u\cdot\nabla)u\,\Delta u\,dx \\ &+ \int_{\mathbb{R}^3}\frac{1}{\rho}\nabla\Pi\,\Delta u\,dx - \int_{\mathbb{R}^3}\frac{1}{\rho}(B\cdot\nabla)B\,\Delta u\,dx.\end{aligned} \tag{4.3}$$

Taking L^2 inner product of the magnetic fields equation of (1.3) with ΔB that

$$\frac{1}{2}\frac{d}{dt}\|\nabla B(t)\|_{L^2}^2 + \|\Delta B(t)\|_{L^2}^2 = \int_{\mathbb{R}^3}(u\cdot\nabla)B\,\Delta B\,dx - \int_{\mathbb{R}^3}(B\cdot\nabla)u\,\Delta B\,dx. \tag{4.4}$$

Thanks to the equation of (1.3) and div$u = 0$, we have

$$\begin{aligned}&\|\Delta u(t)\|_{L^2} + \|\nabla\Pi\|_{L^2} \\ \le&\sqrt{2}\|\Delta u(t) - \nabla\Pi(t)\|_{L^2} \\ \le&\sqrt{2}\|\rho\partial_t u + \rho u\cdot\nabla u - B\cdot\nabla B\|_{L^2} \\ \le& C\,\|\sqrt{\rho}\partial_t u\|_{L^2} + C\,\|u(t)\|_{L^2}^{1/2}\|\nabla u(t)\|_{L^2}^{1/2}\|\Delta u(t)\|_{L^2} \\ &+ C\,\|B(t)\|_{L^2}^{1/2}\|\nabla B(t)\|_{L^2}^{1/2}\|\Delta B(t)\|_{L^2}.\end{aligned} \tag{4.5}$$

Combining (4.3), (4.4) and (4.5), we obtain

$$\begin{aligned}&\frac{d}{dt}\|(\nabla u(t), \nabla B(t))\|_{L^2}^2 + c\,\|(\nabla^2 u(t), \nabla^2 B(t))\|_{L^2}^2 \\ \le& C\,\|(u(t), B(t))\|_{L^2}^{1/2}\|(\nabla u(t), \nabla B(t))\|_{L^2}^{1/2}\|(\nabla^2 u(t), \nabla^2 B(t))\|_{L^2}^2 \\ &+ C\,\|\sqrt{\rho}\partial_t u(t)\|_{L^2}^2,\end{aligned} \tag{4.6}$$

for some positive constant c.

On the other hand, taking L^2 inner product of the momentum equations in (1.3) with $\partial_t u$, we obtain

$$\|\sqrt{\rho}\partial_t u(t)\|_{L^2}^2 + \frac{1}{2}\frac{d}{dt}\|\nabla u(t)\|_{L^2}^2 = -\int_{\mathbb{R}^3} \rho\, u\cdot\nabla u\,\partial_t u\,dx + \int_{\mathbb{R}^3} B\cdot\nabla B\,\partial_t u\,dx.$$

Taking L^2 inner product of the magnetic field equations in (1.3) with $\partial_t B$, we have

$$\|\partial_t B(t)\|_{L^2}^2 + \frac{1}{2}\frac{d}{dt}\|\nabla B(t)\|_{L^2}^2 = -\int_{\mathbb{R}^3} u\cdot\nabla B\,\partial_t B\,dx + \int_{\mathbb{R}^3} B\cdot\nabla u\,\partial_t B\,dx.$$

The above two equalities gives

$$\begin{aligned}\frac{d}{dt}\|(\nabla u(t),\nabla B(t))\|_{L^2}^2 &+ \|(\sqrt{\rho}\partial_t u(t),\partial_t B(t))\|_{L^2}^2 \\ &\le C\,\|(u,B)\|_{L^2}\|(\nabla u,\nabla B)\|_{L^2}\|(\nabla^2 u,\nabla^2 B)\|_{L^2}^2.\end{aligned} \tag{4.7}$$

The above inequalities along with (4.6) ensures a positive constant e_1 such that

$$\frac{d}{dt}\|(\nabla u,\nabla B)\|_{L^2}^2 + e_1\|(\partial_t u,\partial_t B)\|_{L^2}^2 + \mathcal{A}\|(\nabla^2 u,\nabla^2 B)\|_{L^2}^2 \le 0. \tag{4.8}$$

where

$$\mathcal{A} := \frac{c}{2\,C} - \frac{1}{2}\|(u,B)\|_{L^2}^{1/2}\|(\nabla u,\nabla B)\|_{L^2}^{1/2} - C\,\|(u,B)\|_{L^2}\|(\nabla u,\nabla B)\|_{L^2}$$

By (3.2), for any $\eta>0$, there exists $t_0 = t_0(\eta) > 0$ such that

$$\|\nabla u(t_0)\|_{L^2} + \|\nabla B(t_0)\|_{L^2} \le \eta.$$

Now choosing $\eta > 0$ small enough such that

$$\eta^{1/2}\|(u_0,B_0)\|_{L^2}^{1/2}\left(1 + \|(u_0,B_0)\|_{L^2}^{1/2}\eta^{1/2}\right) \le \frac{c}{16\,C^2}, \tag{4.9}$$

we define

$$\tau^* := \sup\left\{t \ge t_0 \,:\, \|(\nabla u(t),\nabla B(t))\|_{L^2} \le 2\eta\right\}. \tag{4.10}$$

We claim that $\tau^* = \infty$. Indeed, if $\tau^* < \infty$, (4.8) and (4.9) imply that

$$\begin{aligned}\frac{d}{dt}\|(\nabla u,\nabla B)\|_{L^2}^2 &+ e_1\|(\partial_t u,\partial_t B)\|_{L^2}^2 \\ &+ \frac{c}{4\,C}\|(\nabla^2 u,\nabla^2 B)\|_{L^2}^2 \le 0 \quad \text{for } t\in[t_0,\tau^*],\end{aligned}$$

which gives

$$\begin{aligned}\|(\nabla u,\nabla B)\|_{L^2}^2 + \int_{t_0}^{\tau^*} e_1\|(\partial_t u,\partial_t B)\|_{L^2}^2\,d\tau + \int_{t_0}^{\tau^*}\frac{c}{4\,C}\|(\nabla^2 u,\nabla^2 B)\|_{L^2}^2\,d\tau \\ \le \|(\nabla u(t_0),\nabla B(t_0))\|_{L^2}^2 \le \eta^2.\end{aligned}$$

This is a contradiction to the definition of τ^*, and thus $\tau^* = \infty$. Then the proof is completed. □

Proposition 4.2. Under the assumptions of Theorem 1.2, there holds $u, B \in C([0,\infty); L^p(\mathbb{R}^3))$ where $p \in (1, \frac{6}{5})$.

Proof. Multiplying the u^i equation in (1.3) by $|u^i|^{p-1}\mathrm{sgn}(u^i)$ for $i = 1, 2, 3$ and integrating the resulting equation over $\mathbb{R}^3$, we obtain that

$$\begin{aligned}\frac{1}{p}\frac{d}{dt}\int_{\mathbb{R}^3}\rho|u^i|^p\,dx + \frac{4\,(p-1)}{p^2}\int_{\mathbb{R}^3}|\nabla|u^i|^{p/2}|^2\,dx\\ \lesssim \int_{\mathbb{R}^3} B\cdot\nabla B^i\,|u^i|^{p-1}\mathrm{sgn}(u^i)\,dx + \|\nabla\Pi\|_{L^p}\|u^i\|_{L^p}^{p-1}.\end{aligned}$$

Using similar ideas, we have

$$\begin{aligned}&\frac{1}{p}\frac{d}{dt}\int_{\mathbb{R}^3}|B^i|^p\,dx + \frac{4\,(p-1)}{p^2}\int_{\mathbb{R}^3}|\nabla|B^i|^{p/2}|^2\,dx\\ &= -\int_{\mathbb{R}^3} u\cdot\nabla B^i\,|B^i|^{p-1}\mathrm{sgn}(B^i)\,dx + \int_{\mathbb{R}^3} B\cdot\nabla u^i\,|B^i|^{p-1}\mathrm{sgn}(B^i)\,dx\end{aligned}$$

Using Hólder's inequalities, we obtain

$$\begin{aligned}\frac{d}{dt}\|\rho^{1/p}u\|_{L^p}^p + \frac{4\,(p-1)}{p}\int_{\mathbb{R}^3}|\nabla|u|^{p/2}|^2\,dx\\ \lesssim \|B\|_{\frac{2p}{2-p}}\|\nabla B\|_{L^2}\|u\|_{L^p}^{p-1} + \|\nabla\Pi\|_{L^p}\|u\|_{L^p}^{p-1}.\end{aligned}$$

Considering $p\in(1,\frac{6}{5})$, we can easily obtain that

$$\begin{aligned}\frac{d}{dt}\|\rho^{1/p}u\|_{L^p}^p + \frac{4\,(p-1)}{p}\int_{\mathbb{R}^3}|\nabla|u|^{p/2}|^2\,dx\\ \lesssim \left(\|B\|_{L^2\cap L^3}\|\nabla B\|_{L^2} + \|\nabla\Pi\|_{L^p}\right)\|u\|_{L^p}^{p-1}.\end{aligned}\tag{4.11}$$

Also by simple calculation, we have

$$\begin{aligned}\frac{d}{dt}\|B\|_{L^p}^p + \frac{4\,(p-1)}{p}\int_{\mathbb{R}^3}|\nabla|B|^{p/2}|^2\,dx\\ \lesssim \|u\|_{L^2\cap L^3}\|\nabla B\|_{L^2}\|B\|_{L^p}^{p-1} + \|B\|_{L^2\cap L^3}\|\nabla u\|_{L^2}\|B\|_{L^p}^{p-1}.\end{aligned}\tag{4.12}$$

Summing up (4.11) and (4.12), we easily obtain that

$$\begin{aligned}\|(u,B)\|_{L_t^\infty(L^p)} \lesssim \|(u_0,B_0)\|_{L^p} + \|\nabla\Pi\|_{L_t^1(L^p)} + \int_0^t\|B\|_{L^2\cap L^3}\|\nabla u\|_{L^2}\,d\tau\\ + \int_0^t\left(\|B\|_{L^2\cap L^3} + \|u\|_{L^2\cap L^3}\right)\|\nabla B\|_{L^2}\,d\tau\end{aligned}\tag{4.13}$$

On the other hand, applying the operator div to the first equation in (1.2), we get

$$\Delta\Pi = \mathrm{div}\left(-u\cdot\nabla u + (1+a)B\cdot\nabla B + a\,(\Delta u - \nabla\Pi)\right),$$

which together with the classical elliptic estimates implies

$$\begin{aligned}\|\nabla\Pi\|_{L^p} &\lesssim \|u\cdot\nabla u\|_{L^p} + \|a\,(\Delta u - \nabla\Pi)\|_{L^p} + \|(1+a)B\cdot\nabla B\|_{L^p}\\ &\lesssim \|u\|_{L^2\cap L^3}\|\nabla u\|_{L^2} + \|a\|_{L^2\cap L^3}\|\Delta u - \nabla\Pi\|_{L^2}\\ &\quad + (1+\|a\|_{L^\infty})\,\|B\|_{L^2\cap L^3}\|\nabla B\|_{L^2}.\end{aligned}\tag{4.14}$$

Due to (4.7), we have

$$\begin{aligned}(4.15)\qquad &\|(\partial_t u, \partial_t B)\|^2_{L^2_t(L^2)}\\ \lesssim &\|(\nabla u_0, \nabla B_0)\|^2_{L^2}\\ &+ \|(u,B)\|_{L^\infty_t(L^2)}\|(\nabla u, \nabla B)\|_{L^\infty_t(L^2)}\|(\nabla^2 u, \nabla^2 B)\|^2_{L^2_t(L^2)}\\ \leq &C.\end{aligned}$$

which together with (4.14) gives rise to

$$\begin{aligned}(4.16)\qquad \|\Delta u - \nabla\Pi\|^2_{L^2_t(L^2)} \lesssim &\|\partial_t u\|^2_{L^2_t(L^2)} + \|u\|_{L^\infty_t(L^2)}\|\nabla u\|_{L^\infty_t(L^2)}\|\Delta u\|^2_{L^2_t(L^2)}\\ &+ \|B\|_{L^\infty_t(L^2)}\|\nabla B\|_{L^\infty_t(L^2)}\|\Delta B\|^2_{L^2_t(L^2)}\\ \leq &C.\end{aligned}$$

Therefore, thanks to $H^1(\mathbb{R}^3) \hookrightarrow L^2 \cap L^3(\mathbb{R}^3)$, together with (4.14), we get

$$\|\nabla\Pi\|_{L^1_t(L^p)} \leq C(t),$$

so that we have

$$\|u\|_{L^\infty_t(L^p)} + \|B\|_{L^\infty_t(L^p)} \leq C(t)$$

which, together with (4.15) and the classical Aubin-Lions lemma implies $u \in C([0,\infty); L^p(\mathbb{R}^3))$. □

Proposition 4.3. Under the assumptions of Theorem 1.2, there holds

$$\begin{aligned}\|u(t)\|_{L^2} + \|B(t)\|_{L^2} &\leq C\,(1+t)^{-\beta(p)},\\ \|\nabla u(t)\|_{L^2} + \|\nabla B(t)\|_{L^2} &\leq C\,(1+t)^{1/2-\beta(p)}, \quad \text{for } t \geq t_0,\\ \int_{t_0}^{\infty} (1+t)^{(1+2\,\beta(p))^-}\Big(\|(\partial_t u, \partial_t B)\|^2_{L^2} + \|(\Delta u, \Delta B)\|^2_{L^2} + \|\nabla\Pi\|^2_{L^2}\Big)\,dt &\leq C\end{aligned}$$

where C depends on $m, M, \|a_0\|_{L^2}, \|a_0\|_{L^\infty}, \|u_0\|_{L^p}$ and $\|u_0\|_{H^1}$.

Proof. Step 1 : Rough decay estimate of $\|u(t)\|_{L^2}$ and $\|B(t)\|_{L^2}$. We split the phase space $\mathbb{R}^3$ into two time-dependent regions so that

$$\begin{aligned}\|\nabla u(t)\|^2_{L^2} &= \int_{S_1(t)} |\xi|^2 |\hat{u}(\xi,t)|^2\,d\xi + \int_{S_1(t)^c} |\xi|^2 |\hat{u}(\xi,t)|^2\,d\xi,\\ \|\nabla B(t)\|^2_{L^2} &= \int_{S_2(t)} |\xi|^2 |\hat{B}(\xi,t)|^2\,d\xi + \int_{S_2(t)^c} |\xi|^2 |\hat{B}(\xi,t)|^2\,d\xi,\end{aligned}$$

where $S_1(t) := \Big\{\xi : |\xi| \leq \sqrt{\bar{\rho}/2}\,g(t)\Big\}$, $S_2(t) := \Big\{\xi : |\xi| \leq \sqrt{1/2}\,g(t)\Big\}$ with $\bar{\rho} := \sup_{(t,x)\in\mathbb{R}^+\times\mathbb{R}^3} \rho(x,t) = \sup_{x\in\mathbb{R}^3} \rho_0(x)$ and $g(t)$ satisfies $g(t) \lesssim \langle t\rangle^{-1/2}$ which will be chosen later on. Here and in what follows, we shall always denote $1+t$ by $\langle t\rangle$. Then, thanks to (3.1), we obtain

$$\begin{aligned}(4.17)\qquad &\frac{d}{dt}\left(\|\sqrt{\rho}u(t)\|^2_{L^2} + \|B(t)\|^2_{L^2}\right) + g^2(t)\left(\|\sqrt{\rho}u(t)\|^2_{L^2} + \|B(t)\|^2_{L^2}\right)\\ \leq &\bar{\rho}g^2(t)\int_{S_1(t)} |\hat{u}(\xi,t)|^2\,d\xi + g^2(t)\int_{S_2(t)} |\hat{B}(\xi,t)|^2\,d\xi.\end{aligned}$$

Next, we need to deal with the low-frequency part of u and B, we rewrite the momentum equation and magnetic field equation as follows

$$\begin{aligned}
u(t) =& e^{(t-t_0)\Delta}u(t_0) + \int_{t_0}^{t} e^{(t-t')\Delta}\mathbb{P}\Big(\nabla\cdot(-u\otimes u) + a\,(\Delta u - \nabla\Pi)\\
&+ \nabla\cdot(B\otimes B) + a\,(B\cdot\nabla B)\Big)\,dt',\\
B(t) =& e^{(t-t_0)\Delta}B(t_0) + \int_{t_0}^{t} e^{(t-t')\Delta}\Big(B\cdot\nabla u(t') - u\cdot\nabla B(t')\Big)\,dt',
\end{aligned}$$

where $\mathbb{P}$ denote the Leray projection operator ant t_0 is the positive time determined by Proposition 4.1. Taking Fourier transform with respect to x-variables leads to

$$\begin{aligned}
|\hat{u}(\xi,t)| \lesssim& e^{-(t-t_0)|\xi|^2}|\hat{u}_{t_0}(\xi)| + \int_{t_0}^{t} e^{-(t-t')|\xi|^2}\Big(|\xi|\,|\widehat{u\otimes u}(\xi)| + |\xi|\,|\widehat{B\otimes B}(\xi)|\\
&+ |\mathcal{F}(a\,(\Delta u - \nabla\Pi))(\xi)| + |\mathcal{F}(a\,B\cdot\nabla B)(\xi)|\Big)\,dt',\\
|\hat{B}(\xi,t)| \lesssim& e^{-(t-t_0)|\xi|^2}|\hat{B}_{t_0}(\xi)| + \int_{t_0}^{t} e^{-(t-t')|\xi|^2}\Big(|\xi|\,|\widehat{B\otimes u}(\xi)| + |\xi|\,|\widehat{u\otimes B}(\xi)|\Big)\,dt'.
\end{aligned}$$

which implies that

(4.18)

$$\begin{aligned}
\int_{S_1(t)} |\hat{u}(\xi,t)|^2\,d\xi \lesssim& \int_{S_1(t)} e^{-2(t-t_0)|\xi|^2}|\hat{u}_{t_0}(\xi)|^2\,d\xi + g(t)^5\left(\int_{t_0}^{t}\|\widehat{u\otimes u}\|_{L^\infty_\xi}\,dt'\right)^2\\
&+ g(t)^5\left(\int_{t_0}^{t}\|\widehat{B\otimes B}\|_{L^\infty_\xi}\,dt'\right)^2 + g(t)^3\left(\int_{t_0}^{t}\|\mathcal{F}(a\,B\cdot\nabla B)\|_{L^\infty_\xi}\,dt'\right)^2\\
&+ g(t)^3\left(\int_{t_0}^{t}\|\mathcal{F}(a\,(\Delta u - \nabla\Pi))\|_{L^\infty_\xi}\,dt'\right)^2,\\
\int_{S_2(t)} |\hat{B}(\xi,t)|^2\,d\xi \lesssim& \int_{S_2(t)} e^{-2(t-t_0)|\xi|^2}|\hat{B}_{t_0}(\xi)|^2\,d\xi + g(t)^5\left(\int_{t_0}^{t}\|\widehat{B\otimes u}\|_{L^\infty_\xi}\,dt'\right)^2.
\end{aligned}$$

Thanks to Proposition 4.1 and (3.2), we have

$$\begin{aligned}
\left(\int_{t_0}^{t}\|\mathcal{F}(a\,(\Delta u - \nabla\Pi))\|_{L^\infty_\xi}\,dt'\right)^2 \le&\\
&\|a\|^2_{L^\infty_t(L^2)}\left(\int_{t_0}^{t}\|\Delta u - \nabla\Pi\|_{L^2}\,dt'\right)^2 \lesssim t - t_0,
\end{aligned}$$

and

$$\begin{aligned}
\left(\int_{t_0}^{t}\|\mathcal{F}(a\,B\cdot\nabla B)\|_{L^\infty_\xi}\right)^2 \lesssim& \left(\int_{t_0}^{t}\|a\,B\cdot\nabla B\|_{L^1}\,dt'\right)^2\\
\lesssim& \|a\|^2_{L^\infty_t(L^\infty)}\left(\int_{t_0}^{t}\|B\|_{L^2}\|\nabla B\|_{L^2}\,dt'\right)^2\\
\lesssim& \ t - t_0,
\end{aligned}$$

while it is easy to see that

$$\begin{aligned}&\left(\int_{t_0}^t \|\widehat{u\otimes u}\|_{L^\infty_\xi}+\|\widehat{B\otimes B}\|_{L^\infty_\xi}+\|\widehat{B\otimes u}\|_{L^\infty_\xi}\,dt'\right)^2\\ \lesssim&\left(\int_{t_0}^t \|u\|_{L^2}^2\,dt'\right)^2+\left(\int_{t_0}^t \|B\|_{L^2}^2\,dt'\right)^2+\left(\int_{t_0}^t \|B\|_{L^2}\|u\|_{L^2}\,dt'\right)^2\\ \lesssim&(t-t_0)^2.\end{aligned}$$

By Proposition 4.2, we know that $u(t_0), B(t_0)\in L^p(\mathbb{R}^3)$ for $1<p<\frac{6}{5}$, and for $\frac{1}{q}:=\frac{4}{3}\beta(p)=\frac{2}{p}-1$ and $\frac{1}{p}+\frac{1}{p'}=1$,

$$\begin{aligned}\int_{S_1(t)} e^{-2(t-t_0)|\xi|^2}|\hat{u}(\xi,t_0)|^2\,d\xi&\lesssim\left(\int_{S_1(t)} e^{-2q(t-t_0)|\xi|^2}\,d\xi\right)^{1/q}\|\hat{u}(\xi,t_0)\|_{L^{p'}}^2\\ &\lesssim\|u(t_0)\|_{L^p}^2\langle t\rangle^{-2\beta(p)}\end{aligned}\tag{4.19}$$

and

$$\begin{aligned}\int_{S_2(t)} e^{-2(t-t_0)|\xi|^2}|\hat{B}(\xi,t_0)|^2\,d\xi&\lesssim\left(\int_{S_2(t)} e^{-2q(t-t_0)|\xi|^2}\,d\xi\right)^{1/q}\|\hat{B}(\xi,t_0)\|_{L^{p'}}^2\\ &\lesssim\|B(t_0)\|_{L^p}^2\langle t\rangle^{-2\beta(p)}\end{aligned}\tag{4.20}$$

where we used the Hausforff-Young inequality [24]. Then since $g(t)\lesssim\langle t\rangle^{-1/2}$, we deduce from (4.18) that

$$\int_{S_1(t)}|\hat{u}(\xi,t)|^2\,d\xi\lesssim\langle t\rangle^{-2\beta(p)}+\langle t\rangle^{-\frac{1}{2}}\lesssim\langle t\rangle^{-\frac{1}{2}}\quad\text{for } t\geq t_0,\tag{4.21}$$

and

$$\int_{S_2(t)}|\hat{B}(\xi,t)|^2\,d\xi\lesssim\langle t\rangle^{-2\beta(p)}+\langle t\rangle^{-\frac{1}{2}}\lesssim\langle t\rangle^{-\frac{1}{2}}\quad\text{for } t\geq t_0,\tag{4.22}$$

where we used the fact $\frac{1}{2}<\beta(p)<\frac{3}{4}$. Substituting (4.21) and (4.22) into (4.17) results in

$$\frac{d}{dt}\|(\sqrt{\rho}u,B)\|_{L^2}^2+g^2(t)\|(\sqrt{\rho}u,B)\|_{L^2}^2\lesssim g^2(t)\langle t\rangle^{-\frac{1}{2}}\lesssim\langle t\rangle^{-\frac{3}{2}}\quad\text{for } t\geq t_0,$$

which gives

$$e^{\int_{t_0}^t g^2(t)\,dt'}\|(\sqrt{\rho}u,B)\|_{L^2}^2\lesssim\|(\sqrt{\rho}u(t_0),B(t_0))\|_{L^2}^2+\int_{t_0}^t e^{\int_{t_0}^{t'} g(\tau)^2\,d\tau}\langle t'\rangle^{-\frac{3}{2}}\,dt'$$

Taking $g^2(t):=\frac{\alpha}{1+t}$ (with $\alpha>\frac{1}{2}$) in the above inequality, we infer

$$\left(\|\sqrt{\rho}u(t)\|_{L^2}^2+\|B(t)\|_{L^2}^2\right)\langle t\rangle^\alpha\lesssim 1+\langle t\rangle^{\alpha-\frac{1}{2}},$$

which gives

$$\|u(t)\|_{L^2}+\|B(t)\|_{L^2}\lesssim\langle t\rangle^{-\frac{1}{4}}.\tag{4.23}$$

Step 2 : Rough decay estimate of $\|\nabla u(t)\|_{L^2}$ and $\|\nabla B(t)\|_{L^2}$. We split the phase space $\mathbb{R}^3$ into two time-dependent regions so that

$$\|\nabla^2 u(t)\|_{L^2}^2 = \int_{S(t)} |\xi|^4 |\hat{u}(\xi,t)|^2\,d\xi + \int_{S(t)^c} |\xi|^2 |\widehat{\nabla u}(\xi,t)|^2\,d\xi,$$
$$\|\nabla^2 B(t)\|_{L^2}^2 = \int_{S(t)} |\xi|^4 |\hat{B}(\xi,t)|^2\,d\xi + \int_{S(t)^c} |\xi|^2 |\widehat{\nabla B}(\xi,t)|^2\,d\xi,$$

where $S(t) := \Big\{\xi : |\xi| \le \sqrt{\frac{1}{e_2}}\, g(t)\Big\}$ and $g(t) \lesssim \langle t\rangle^{-\frac{1}{2}}$, which will be chosen later on. Then we deduce from Proposition 4.1 that

$$\begin{aligned}\frac{d}{dt}\|(\nabla u, \nabla B)\|_{L^2}^2 + e_1\|(\partial_t u, \partial_t B)\|_{L^2}^2 + g^2(t)\|(\nabla u, \nabla B)\|_{L^2}^2 \\ \le g^4(t)\int_{S(t)} |\hat{u}(\xi,t)|^2 + |\hat{B}(\xi,t)|^2\,d\xi \lesssim g^4(t)\langle t\rangle^{-\frac{1}{2}} \lesssim \langle t\rangle^{-\frac{5}{2}},\end{aligned} \tag{4.24}$$

which gives

$$\begin{aligned}e^{\int_{t_0}^t g(t')^2\,dt'}\|(\nabla u, \nabla B)\|_{L^2}^2 + e_1\int_{t_0}^t e^{\int_{t_0}^{t'} g(\tau)^2\,d\tau}\|(\partial_t u, \partial_t B)\|_{L^2}^2\,dt' \\ \lesssim \|(\nabla u(t_0), \nabla B(t_0))\|_{L^2}^2 + \int_{t_0}^t e^{\int_{t_0}^{t'} g(\tau)^2\,d\tau}\langle t\rangle^{-\frac{5}{2}}\,dt'\end{aligned}$$

Taking $g^2(t) := \frac{\alpha}{1+t}$ (with $\alpha > 1$) in the above inequality, we deduce that

$$\|(\nabla u, \nabla B)\|_{L^2}^2\langle t\rangle^\alpha + e_1\int_{t_0}^t \langle t'\rangle^\alpha \|(\partial_t u, \partial_t B)\|_{L^2}^2\,dt' \lesssim 1 + \int_{t_0}^t \langle t'\rangle^{\alpha-\frac{5}{2}}\,dt'. \tag{4.25}$$

In particular, for $\alpha > \frac{3}{2}$, we get

$$\|(\nabla u, \nabla B)\|_{L^2}^2\langle t\rangle^\alpha \lesssim 1 + \langle t\rangle^{\alpha-\frac{3}{2}} \quad \text{for } t \ge t_0,$$

which implies

$$\|\nabla u(t)\|_{L^2} + \|\nabla B(t)\|_{L^2} \lesssim \langle t\rangle^{-\frac{3}{4}} \quad \text{for } t \ge t_0. \tag{4.26}$$

Moreover, it follows from (4.25) that

$$\int_{t_0}^t \langle t'\rangle^{\left(\frac{3}{2}\right)^-}\|(\partial_t u, \partial_t B)\|_{L^2}^2\,dt' \lesssim 1,$$

which ensures that

$$\begin{aligned}&\left(\int_{t_0}^t \|(\partial_t u, \partial_t B)\|_{L^2}\right)^2 \\ &\lesssim \int_{t_0}^t \langle t\rangle^{\left(\frac{3}{2}\right)^-}\|(\partial_t u, \partial_t B)\|_{L^2}^2\,dt' \int_{t_0}^t \langle t\rangle^{-\left(\frac{3}{2}\right)^-}\,dt' \lesssim 1.\end{aligned} \tag{4.27}$$

Step 3 : Improved decay estimates of $\|u(t)\|_{L^2}$, $\|\nabla u(t)\|_{L^2}$ and $\|B(t)\|_{L^2}$, $\|\nabla B(t)\|_{L^2}$. We shall utilize an iteration argument. Thanks to (4.23), we obtain

$$\begin{aligned}\left(\int_{t_0}^t \|\mathcal{F}(u\otimes u)(t')\|_{L^\infty_\xi}\,dt'\right)^2 &\lesssim \left(\int_{t_0}^t \|u(t')\|_{L^2}^2\,dt'\right)^2 \\ &\lesssim \left(\int_{t_0}^t \langle t\rangle^{-\frac{1}{2}}\,dt'\right)^2 \lesssim \langle t\rangle,\end{aligned} \tag{4.28}$$

and similar we have

$$\left(\int_{t_0}^{t} \|\mathcal{F}(u\otimes B)(t')\|_{L^\infty_\xi}\,dt'\right)^2 \lesssim \langle t\rangle. \tag{4.29}$$

Same argument as in [21], we have

$$\left(\int_{t_0}^{t} \|\mathcal{F}(a\,(\Delta u-\nabla\Pi))(t')\|_{L^\infty_\xi}\,dt'\right)^2 \le C. \tag{4.30}$$

Using (4.23) and (4.26), we obtain

$$\begin{aligned}
&\left(\int_{t_0}^{t} \|\mathcal{F}(a\,B\cdot\nabla B)(t')\|_{L^\infty_\xi}\,dt'\right)^2\\
&\lesssim \left(\int_{t_0}^{t} \|a\,B\cdot\nabla B\|_{L^1}\,dt'\right)^2\\
&\lesssim \left(\int_{t_0}^{t} \|B\|_{L^2}\|\nabla B\|_{L^2}\,dt'\right)^2\\
&\lesssim \left(\int_{t_0}^{t} \langle t\rangle^{-\frac14}\langle t\rangle^{-\frac34}\right)^2 \lesssim (\ln\langle t\rangle)^2
\end{aligned} \tag{4.31}$$

Substituting (4.19), (4.20), (4.28), (4.29), (4.30) and (4.31) into (4.18) results in

$$\int_{S(t)} |\hat u(\xi,t)|^2\,d\xi \lesssim \langle t\rangle^{-2\beta(p)} + \langle t\rangle^{-\frac32+\epsilon} \lesssim \langle t\rangle^{-2\beta(p)}, \tag{4.32}$$

and

$$\int_{S(t)} |\hat B(\xi,t)|^2\,d\xi \lesssim \langle t\rangle^{-2\beta(p)} + \langle t\rangle^{-\frac32} \lesssim \langle t\rangle^{-2\beta(p)}, \tag{4.33}$$

where $\epsilon>0$ is a small enough constant. From this and (4.17), we have

$$\frac{d}{dt}\|(\sqrt\rho u, B)\|_{L^2}^2 + g^2(t)\|(\sqrt\rho u, B)\|_{L^2}^2 \lesssim g^2(t)\langle t\rangle^{-2\beta(p)} \lesssim \langle t\rangle^{-1-2\beta(p)},$$

which implies

$$e^{\int_{t_0}^t g(t')^2\,dt'}\|(\sqrt\rho u, B)\|_{L^2}^2 \lesssim \|(\sqrt\rho u(t_0), B(t_0))\|_{L^2}^2 + \int_{t_0}^{t} e^{\int_{t_0}^{t'} g(\tau)^2\,d\tau}\langle t\rangle^{-1-2\beta(p)}\,dt'$$

Taking $g^2(t)=\frac{\alpha}{1+t}$ (with $\alpha>2\beta(p)$) in the above inequality leads to

$$\left(\|\sqrt\rho u(t)\|_{L^2}^2 + \|B(t)\|_{L^2}^2\right)\langle t\rangle^{\alpha} \lesssim 1+\langle t\rangle^{\alpha-2\beta(p)},$$

which in particular gives

$$\|u(t)\|_{L^2} + \|B(t)\|_{L^2} \lesssim \langle t\rangle^{-\beta(p)}.$$

From (4.32), (4.33) and (4.24), we infer

$$\begin{aligned}
&\frac{d}{dt}\|(\nabla u,\nabla B)\|_{L^2}^2 + e_1\|(\partial_t u,\partial_t B)\|_{L^2}^2\\
&\qquad + g^2(t)\|(\nabla u,\nabla B)\|_{L^2}^2 \lesssim g^4(t)\langle t\rangle^{-2\beta(p)} \lesssim \langle t\rangle^{-2-2\beta(p)},
\end{aligned}$$

which implies

$$e^{\int_{t_0}^t g(t')^2\,dt'}\|(\nabla u,\nabla B)\|_{L^2}^2+e_1\int_{t_0}^t e^{\int_{t_0}^{t'} g(\tau)^2\,d\tau}\|(\partial_t u,\partial_t B)\|_{L^2}^2\,dt'$$
$$\lesssim \|(\nabla u(t_0),\nabla B(t_0))\|_{L^2}^2+\int_{t_0}^t e^{\int_{t_0}^{t'} g(\tau)\,d\tau}\langle t\rangle^{-2-2\beta(p)}\,dt'$$

Taking $g^2(t)=\frac{\alpha}{1+t}$ (with $\alpha>1$) in the above inequality, we obtain

$$\|(\nabla u,\nabla B)\|_{L^2}^2\langle t\rangle^{\alpha}+e_1\int_{t_0}^t\langle t'\rangle^{\alpha}\|(\partial_t u,\partial_t B)\|_{L^2}^2\,dt'$$
$$\lesssim 1+\int_{t_0}^t\langle t'\rangle^{\alpha-2-2\beta(p)}\,dt'. \tag{4.34}$$

In particular, taking $\alpha>1+2\beta(p)$ in (4.34), we get

$$\|\nabla u(t)\|_{L^2}+\|\nabla B(t)\|_{L^2}\lesssim\langle t\rangle^{-\frac{1}{2}-\beta(p)}. \tag{4.35}$$

Taking $\alpha\in(\frac{3}{2},1+2\beta(p))$ in (4.34) results in

$$\int_{t_0}^t\langle t'\rangle^{(1+2\beta(p))^-}\left(\|\partial_t u(t')\|_{L^2}^2+\|\partial_t B(t')\|_{L^2}^2\right)dt'\le C.$$

Combining the following fact

$$\|u\cdot\nabla u\|_{L^2}^2\le C\,\|u\|_{L^\infty}^2\|\nabla u\|_{L^2}^2\le C\,\|\Delta u\|_{L^2}\|\nabla u\|_{L^2}^3,$$
$$\|B\cdot\nabla B\|_{L^2}^2\le C\,\|B\|_{L^\infty}^2\|\nabla B\|_{L^2}^2\le C\,\|\Delta B\|_{L^2}\|\nabla B\|_{L^2}^3,$$

with (4.5), we have

$$\int_{t_0}^\infty\langle t'\rangle^{(1+2\beta(p))^-}\left(\|\Delta u(t')\|_{L^2}^2+\|\Delta B(t')\|_{L^2}^2+\|\nabla\Pi(t')\|_{L^2}^2\right)dt'$$
$$\lesssim\int_{t_0}^\infty\langle t\rangle^{(1+2\beta(p))^-}\|(\partial_t u,\partial_t B)\|_{L^2}^2\,dt'+\int_{t_0}^\infty\langle t\rangle^{-\frac{1}{2}(1+2\beta(p))^-}\|(\Delta u,\Delta B)\|_{L^2}\,dt'\le C, \tag{4.36}$$

where we used (4.35) and (4.2). □

Remark 4.4. Thanks to (4.36), it is easy to observe that

$$\int_{t_0}^\infty\langle t'\rangle^{\alpha}\left(\|\Delta u(t')\|_{L^2}^2+\|\Delta B(t')\|_{L^2}^2+\|\nabla\Pi(t')\|_{L^2}^2\right)dt'\lesssim\langle t\rangle^{\alpha-(1+2\beta(p))}, \tag{4.37}$$

for $\alpha\in(1+2\beta(p),2+6\beta(p))$. Moreover, without loss of generality, we may assume that

$$\|\Delta u(t_0)\|_{L^2}+\|\Delta B(t_0)\|_{L^2}\le 1. \tag{4.38}$$

Proposition 4.5. Under the assumptions of Theorem 1.2, there holds

$$\int_{t_0}^\infty\left(\|u(t)\|_{L^\infty}+\|B(t)\|_{L^\infty}+\|\nabla u(t)\|_{L^\infty}+\|\nabla B(t)\|_{L^\infty}\right)dt\le C \tag{4.39}$$

and

$$\int_{t_0}^\infty\left(\|\Delta u(t)\|_{L^2}^\theta+\|\Delta B(t)\|_{L^2}^\theta+\|\nabla\Pi(t)\|_{L^2}^\theta\right)dt\le C, \tag{4.40}$$

for $\frac{2}{3}\le\theta\le 2$.

Proof. Step 1 : Estimate of $\|u_t(t)\|_{L^2}$, $\|u(t)\|_{\dot{H}^2}$ and $\|B_t(t)\|_{L^2}$, $\|B(t)\|_{\dot{H}^2}$. We get by first applying ∂_t to the momentum equation of (1.3) and then taking L^2 inner product of the resulting equation with $\partial_t u$ that

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\|\sqrt{\rho}u_t(t)\|_{L^2}^2+\|\nabla u_t(t)\|_{L^2}^2=&-\int_{\mathbb{R}^3}\rho_t u_t\cdot(u\cdot\nabla u)\\&-\int_{\mathbb{R}^3}\rho u_t\cdot(u_t\cdot\nabla u)+\rho_t|u_t|^2-B_t\cdot\nabla B\cdot u_t-B\cdot\nabla B_t\cdot u_t\,dx.\end{aligned}\tag{4.41}$$

Applying ∂_t to the magnetic field equation of (1.3) and then taking L^2 inner product of the resulting equation with $\partial_t B$ that

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\|B_t(t)\|_{L^2}^2+\|\nabla B_t(t)\|_{L^2}^2=&\\-\int_{\mathbb{R}^3}u_t\cdot\nabla B\cdot B_t-B_t\cdot\nabla u\cdot B_t-B\cdot\nabla u_t\cdot B_t\,dx.&\end{aligned}\tag{4.42}$$

Using similar methods in [21], we obtain

$$\left|\int_{\mathbb{R}^3}\rho_t u_t\cdot(u\cdot\nabla u)\,dx\right|\le\frac{1}{8}\|\nabla u_t\|_{L^2}^2+C\,\|\nabla u\|_{L^2}^4\|\nabla^2 u\|_{L^2}^2,$$

$$\left|\int_{\mathbb{R}^3}\rho u_t\cdot(u_t\cdot\nabla u)\,dx\right|\le\frac{1}{8}\|\nabla u_t\|_{L^2}^2+C\,\|\sqrt{\rho}u_t\|_{L^2}^2\|\nabla u\|_{L^2}^4,$$

$$\left|\int_{\mathbb{R}^3}\rho_t|u_t|^2\,dx\right|\le\frac{1}{8}\|\nabla u_t\|_{L^2}^2+C\,\|\nabla u\|_{L^2}^4\|\sqrt{\rho}u_t\|_{L^2}^2,$$

$$\left|\int_{\mathbb{R}^2}B_t\cdot\nabla B\cdot u_t\,dx\right|\le\frac{1}{8}\|\nabla u_t\|_{L^2}^2+\frac{1}{8}\|\nabla B_t\|_{L^2}^2+C\,\|B_t\|_{L^2}^2\|\nabla B\|_{L^2}^4,$$

and

$$\left|\int_{\mathbb{R}^3}B_t\cdot(B_t\cdot\nabla u)\,dx\right|\le\frac{1}{8}\|\nabla B_t\|_{L^2}^2+C\,\|B_t\|_{L^2}^2\|\nabla u\|_{L^2}^4.$$

Substituting the above five inequalities into (4.41), (4.42), we easily obtain

$$\begin{aligned}\frac{d}{dt}\|(\sqrt{\rho}u_t,B_t)\|_{L^2}^2+\|(\nabla u_t,\nabla B_t)\|_{L^2}^2\lesssim&\|\nabla u\|_{L^2}^4\|\nabla^2 u\|_{L^2}^2+\|\sqrt{\rho}u_t\|_{L^2}^2\|\nabla u\|_{L^2}^4\\&+\|B_t\|_{L^2}^2\|\nabla B\|_{L^2}^4+\|B_t\|_{L^2}^2\|\nabla u\|_{L^2}^4.\end{aligned}$$

Integrating the above inequality from t_0 to t, we get

$$\begin{aligned}&\|\sqrt{\rho}u_t(t)\|_{L^2}^2+\|B_t(t)\|_{L^2}^2+\int_{t_0}^t\|(\nabla u_t,\nabla B_t)\|_{L^2}^2\,dt'\\\le&\|\sqrt{\rho}u_t(t_0)\|_{L^2}+\|B_t(t_0)\|_{L^2}^2+C\int_{t_0}^t\|\nabla^2 u\|_{L^2}^2\|\nabla u\|_{L^2}^4\,dt'\\&+C\int_{t_0}^t\|\sqrt{\rho}u_t\|_{L^2}^2\|\nabla u\|_{L^2}^4\,dt'+C\int_{t_0}^t\|B_t\|_{L^2}^2\|\nabla B\|_{L^2}^4\,dt'\\&+C\int_{t_0}^t\|B_t\|_{L^2}^2\|\nabla u\|_{L^2}^4\,dt'.\end{aligned}$$

Applying Gronwall's inequality gives

$$\begin{aligned}&\|\sqrt{\rho}u_t(t)\|_{L^2}^2+\|B_t(t)\|_{L^2}^2+\|(\nabla u_t,\nabla B_t)\|_{L^2([t_0,t];L^2)}^2\\&\le e^{C\int_{t_0}^t\|(\nabla u,\nabla B)\|_{L^2}^4\,dt'}\left(\|(\sqrt{\rho}u_t(t_0),B_t(t_0))\|_{L^2}^2+C\int_{t_0}^t\|\nabla^2u\|_{L^2}^2\|\nabla u\|_{L^2}^4\,dt'\right)\end{aligned}$$

It then follows from (3.2) and Proposition 4.1 that

$$\begin{aligned}\int_{t_0}^t\|\nabla u\|_{L^2}^4\,dt'&\le\sup_{t'\in[t_0,t]}\|\nabla u(t')\|_{L^2}^2\|\nabla u\|_{L^2_t(L^2)}^2\le C,\\\int_{t_0}^t\|\nabla B\|_{L^2}^4\,dt'&\le\sup_{t'\in[t_0,t]}\|\nabla B(t')\|_{L^2}^2\|\nabla B\|_{L^2_t(L^2)}^2\le C,\end{aligned}$$

and

$$\int_{t_0}^t\|\nabla^2u(t')\|_{L^2}^2\|\nabla u(t')\|_{L^2}^4\,dt'\le C\sup_{t'\in[t_0,t]}\|\nabla u(t')\|_{L^2}^4\|\Delta u\|_{L^2([t_0,t];L^2)}^2\le C.$$

Applying Remark 4.4 and the standard energy estimate to the momentum and magnetic field equation of (1.3) at $t=t_0$ yields

$$\begin{aligned}\|\sqrt{\rho}u_t(t_0)\|_{L^2}^2&\lesssim\|(u\cdot\nabla u)(t_0)\|_{L^2}^2+\|(B\cdot\nabla B)(t_0)\|_{L^2}^2+\|\Delta u(t_0)\|_{L^2}^2\\&\lesssim\|u(t_0)\|_{L^4}^2\|\nabla u(t_0)\|_{L^4}^2+\|B(t_0)\|_{L^4}^2\|\nabla B(t_0)\|_{L^4}^2+\|\Delta u(t_0)\|_{L^2}^2\\&\lesssim\|u(t_0)\|_{L^2}^{1/2}\|\nabla u(t_0)\|_{L^2}^2\|\Delta u(t_0)\|_{L^2}^{3/2}\\&\quad+\|B(t_0)\|_{L^2}^{1/2}\|\nabla B(t_0)\|_{L^2}^2\|\Delta B(t_0)\|_{L^2}^{3/2}+\|\Delta u(t_0)\|_{L^2}^2\\&\le C,\end{aligned}$$

and

$$\begin{aligned}\|B_t(t_0)\|_{L^2}^2&\lesssim\|\Delta B(t_0)\|_{L^2}^2+\|(u\cdot\nabla)B(t_0)\|_{L^2}^2+\|(B\cdot\nabla)u(t_0)\|_{L^2}^2\\&\lesssim1+\|u(t_0)\|_{L^4}^2\|\nabla B(t_0)\|_{L^4}^2+\|B(t_0)\|_{L^4}^2\|\nabla u(t_0)\|_{L^4}^2\le C.\end{aligned}$$

Hence, we finally have

$$\sup_{t\ge t_0}\left(\|\sqrt{\rho}u_t(t)\|_{L^2}^2+\|B_t(t)\|_{L^2}^2\right)+\int_{t_0}^\infty\left(\|\nabla u_t\|_{L^2}^2+\|\nabla B_t\|_{L^2}^2\right)dt'\le C.\tag{4.43}$$

Notice from the momentum equation of (1.3) that

$$\begin{aligned}\|\nabla^2u(t)\|_{L^2}+\|\nabla\Pi(t)\|_{L^2}&\lesssim\|\sqrt{\rho}u_t(t)\|_{L^2}+\|(u\cdot\nabla)u(t)\|_{L^2}+\|(B\cdot\nabla)B(t)\|_{L^2}\\&\le C\Big(\|\sqrt{\rho}u_t(t)\|_{L^2}+\|u(t)\|_{L^2}\|\nabla u(t)\|_{L^2}^4\\&\qquad+\|B(t)\|_{L^2}\|\nabla B(t)\|_{L^2}^4\Big)+\frac{1}{6}\|(\nabla^2u(t),\nabla^2B(t))\|_{L^2},\end{aligned}$$

and

$$\begin{aligned}\|\nabla^2B(t)\|_{L^2}&\le C\left(\|B_t(t)\|_{L^2}+\|(u\cdot\nabla)B(t)\|_{L^2}+\|(B\cdot\nabla)u(t)\|_{L^2}\right)\\&\le C\left(\|B_t(t)\|_{L^2}+\|u(t)\|_{L^2}\|\nabla u(t)\|_{L^2}^4+\|B(t)\|_{L^2}\|\nabla B(t)\|_{L^2}^4\right)\\&\quad+\frac{1}{6}\|\nabla^2B(t)\|_{L^2}+\frac{1}{6}\|\nabla^2B(t)\|_{L^2},\end{aligned}$$

which along with Proposition 4.1 and (4.43) implies that

$$\sup_{t\geq t_0}\left(\|\nabla^2 u(t)\|_{L^2}+\|\nabla^2 B(t)\|_{L^2}+\|\nabla\Pi(t)\|_{L^2}\right)\leq C. \tag{4.44}$$

Step 2 : Estimate of $\int_{t_0}^{\infty}\|(u(t'),B(t'))\|_{L^\infty}\,dt'$, and $\int_{t_0}^{\infty}\|(\nabla u(t'),\nabla B(t'))\|_{L^\infty}dt'$. Let (v,q) solve

$$-\Delta v+\nabla q=f,\quad \mathrm{div} v=0;$$

then we have $\nabla q=-\nabla(-\Delta)^{-1}\mathrm{div} f$, which implies that for any $r\in(1,\infty)$,

$$\|\nabla q\|_{L^r}\leq C\,\|f\|_{L^r}\quad\text{and}\quad\|\Delta v\|_{L^r}\leq C\,\|f\|_{L^r}.$$

From this and the momentum equation of (1.3), we infer

$$\begin{aligned}
&\|\nabla^2 u(t)\|_{L^6}+\|\nabla\Pi(t)\|_{L^6}\\
\lesssim&\|u_t(t)\|_{L^6}+\|(u\cdot\nabla)u(t)\|_{L^6}+\|(B\cdot\nabla)B(t)\|_{L^6}\\
\lesssim&\|\nabla u_t(t)\|_{L^2}+\|\nabla u(t)\|_{L^2}^{1/2}\|\nabla^2 u(t)\|_{L^2}^{3/2}+\|\nabla B(t)\|_{L^2}^{1/2}\|\nabla^2 B(t)\|_{L^2}^{3/2}\\
&+\|u(t)\|_{L^2}^{1/4}\|\nabla u(t)\|_{L^2}^{3/4}\|\nabla^2 u(t)\|_{L^2}^{1/4}\|\nabla^2 u(t)\|_{L^6}^{3/4}\\
&+\|B(t)\|_{L^2}^{1/4}\|\nabla B(t)\|_{L^2}^{3/4}\|\nabla^2 B(t)\|_{L^2}^{1/4}\|\nabla^2 B(t)\|_{L^6}^{3/4}
\end{aligned}$$

Taking the L^2 norm for the time variables on $[t_0,t]$, we get by using (4.43) and (4.44) that

$$\begin{aligned}
&\|\nabla^2 u\|_{L^2([t_0,t];L^6)}^2+\|\nabla\Pi\|_{L^2([t_0,t];L^6)}^2\\
\leq& C\Big\{\|\nabla u_t\|_{L^2([t_0,t];L^2)}^2+\sup_{t'\in[t_0,t]}\|\nabla u(t')\|_{L^2}\sup_{t'\in[t_0,t]}\|\nabla^2 u(t')\|_{L^2}\|\nabla^2 u\|_{L^2([t_0,t];L^2)}^2\\
&+\sup_{t'\in[t_0,t]}\|u(t')\|_{L^2}^2\sup_{t'\in[t_0,t]}\|\nabla u(t')\|_{L^2}^6\|\nabla^2 u\|_{L^2([t_0,t];L^2)}^2\\
&+\sup_{t'\in[t_0,t]}\|\nabla B(t')\|_{L^2}\sup_{t'\in[t_0,t]}\|\nabla^2 B(t')\|_{L^2}\|\nabla^2 B\|_{L^2([t_0,t];L^2)}^2\\
&+\sup_{t'\in[t_0,t]}\|B(t')\|_{L^2}^2\sup_{t'\in[t_0,t]}\|\nabla B(t')\|_{L^2}^6\|\nabla^2 B\|_{L^2([t_0,t];L^2)}^2\Big\}+\frac{1}{4}\|\nabla^2 B\|_{L^2([t_0,t];L^6)}^2.
\end{aligned}\tag{4.45}$$

Using similar methods, we can easily obtain

$$\begin{aligned}
&\|\nabla^2 B\|_{L^2([t_0,t];L^6)}^2\\
\leq& C\Big\{\|\nabla B_t\|_{L^2([t_0,t];L^2)}^2+\sup_{t'\in[t_0,t]}\|\nabla u(t')\|_{L^2}\sup_{t'\in[t_0,t]}\|\nabla^2 u(t')\|_{L^2}\|\nabla^2 B\|_{L^2([t_0,t];L^2)}^2\\
&+\sup_{t'\in[t_0,t]}\|u(t')\|_{L^2}^2\sup_{t'\in[t_0,t]}\|\nabla u(t')\|_{L^2}^6\|\nabla^2 B\|_{L^2([t_0,t];L^2)}^2\\
&+\sup_{t'\in[t_0,t]}\|\nabla B(t')\|_{L^2}\sup_{t'\in[t_0,t]}\|\nabla^2 B(t')\|_{L^2}\|\nabla^2 u\|_{L^2([t_0,t];L^2)}^2\\
&+\sup_{t'\in[t_0,t]}\|B(t')\|_{L^2}^2\sup_{t'\in[t_0,t]}\|\nabla B(t')\|_{L^2}^6\|\nabla^2 u\|_{L^2([t_0,t];L^2)}^2\Big\}\\
&+\frac{1}{4}\|\nabla^2 B\|_{L^2([t_0,t];L^6)}^2+\frac{1}{4}\|\nabla^2 u\|_{L^2([t_0,t];L^6)}^2.
\end{aligned}\tag{4.46}$$

Summing up (4.45) and (4.46), we get

$$
\begin{aligned}
&\|\nabla^2 u\|^2_{L^2([t_0,t];L^6)} + \|\nabla^2 B\|^2_{L^2([t_0,t];L^6)} + \|\nabla\Pi\|^2_{L^2([t_0,t];L^6)}\\
&\lesssim \|\nabla u_t\|^2_{L^2([t_0,t];L^2)} + \|\nabla B_t\|^2_{L^2([t_0,t];L^2)}\\
&\quad + \sup_{t'\in[t_0,t]}\|\nabla u(t')\|_{L^2}\sup_{t'\in[t_0,t]}\|\nabla^2 u(t')\|_{L^2}\|(\nabla^2 u,\nabla^2 B)\|^2_{L^2([t_0,t];L^2)}\\
&\quad + \sup_{t'\in[t_0,t]}\|u(t')\|^2_{L^2}\sup_{t'\in[t_0,t]}\|\nabla u(t')\|^6_{L^2}\|(\nabla^2 u,\nabla^2 B)\|^2_{L^2([t_0,t];L^2)}\\
&\quad + \sup_{t'\in[t_0,t]}\|\nabla B(t')\|_{L^2}\sup_{t'\in[t_0,t]}\|\nabla^2 B(t')\|_{L^2}\|(\nabla^2 u,\nabla^2 B)\|^2_{L^2([t_0,t];L^2)}\\
&\quad + \sup_{t'\in[t_0,t]}\|B(t')\|^2_{L^2}\sup_{t'\in[t_0,t]}\|\nabla B(t')\|^6_{L^2}\|(\nabla^2 u,\nabla^2 B)\|^2_{L^2([t_0,t];L^2)}\\
&\leq C.
\end{aligned}
\tag{4.47}
$$

Now, let us turn to the estimate of $\int_{t_0}^t \|\nabla u(t')\|_{L^\infty}\,dt'$ and $\int_{t_0}^t \|\nabla B(t')\|_{L^\infty}\,dt'$. Indeed, simple interpolation in three dimensions gives

$$
\begin{aligned}
\int_{t_0}^t \|\nabla u(t')\|_{L^\infty}\,dt' &\lesssim \int_{t_0}^t \|\nabla u(t')\|^{1/2}_{L^6}\|\nabla^2 u(t')\|^{1/2}_{L^6}\,dt'\\
&\lesssim \int_{t_0}^t \|\Delta u(t')\|^{1/2}_{L^2}\|\nabla^2 u(t')\|^{1/2}_{L^6}\,dt'\\
&\lesssim \int_{t_0}^t \|\nabla^2 u(t')\|^2_{L^6}\,dt' + \int_{t_0}^t \|\Delta u(t')\|^{2/3}_{L^2}\,dt',
\end{aligned}
$$

and

$$
\begin{aligned}
\int_{t_0}^t \|\nabla B(t')\|_{L^\infty}\,dt' &\lesssim \int_{t_0}^t \|\nabla B(t')\|^{1/2}_{L^6}\|\nabla^2 B(t')\|^{1/2}_{L^6}\,dt'\\
&\lesssim \int_{t_0}^t \|\Delta B(t')\|^{1/2}_{L^2}\|\nabla^2 B(t')\|^{1/2}_{L^6}\,dt'\\
&\lesssim \int_{t_0}^t \|\nabla^2 B(t')\|^2_{L^6}\,dt' + \int_{t_0}^t \|\Delta B(t')\|^{2/3}_{L^2}\,dt'.
\end{aligned}
$$

Thanks to (4.36), we get

$$
\begin{aligned}
&\int_{t_0}^t \Big(\|\Delta u(t')\|^{2/3}_{L^2} + \|\Delta B(t')\|^{2/3}_{L^2} + \|\nabla\Pi(t')\|^{2/3}_{L^2}\Big)\,dt'\\
&\lesssim \bigg(\int_{t_0}^t \langle t'\rangle^{(1+2\beta(p))^-}\left(\|\Delta u(t')\|^2_{L^2} + \|\Delta B(t')\|^2_{L^2} + \|\nabla\Pi(t')\|^2_{L^2}\right)\,dt'\bigg)^{\frac{1}{3}}\\
&\leq C,
\end{aligned}
\tag{4.48}
$$

which along with (4.47) ensures that

$$
\int_{t_0}^t \|\nabla u(t)\|_{L^\infty} + \|\nabla B(t)\|_{L^\infty}\,dt' \leq C. \tag{4.49}
$$

Due to (4.5) and magnetic field equation, we have

$$\begin{aligned}
&\int_{t_0}^{t} \big(\|\Delta u(t')\|_{L^2}^2 + \|\Delta B(t')\|_{L^2}^2 + \|\nabla \Pi(t')\|_{L^2}^2\big)\, dt' \\
\leq & C\int_{t_0}^{t} \big(\|(\partial_t u(t'), \partial_t B(t'))\|_{L^2}^2 + \|(u\cdot\nabla)u(t')\|_{L^2}^2 + \|(B\cdot\nabla)B(t')\|_{L^2}^2 \\
&+ \|(u\cdot\nabla)B(t')\|_{L^2}^2 + \|(B\cdot\nabla)u(t')\|_{L^2}^2\big)\, dt' \\
\leq & C\int_{t_0}^{t} \|(\partial_t u(t'), \partial_t B(t'))\|_{L^2}^2 + \|u(t')\|_{L^6}^2(\|\nabla u(t')\|_{L^3}^2 + \|\nabla B(t')\|_{L^3}^2) \\
&+ \|B(t')\|_{L^6}^2(\|\nabla u(t')\|_{L^3}^2 + \|\nabla B(t')\|_{L^3}^2)\, dt' \\
\leq & C.
\end{aligned}$$

This together with (4.48) proves (4.40).

At last, using (4.3), we can get that

$$\begin{aligned}
\|u(t)\|_{L^\infty} \lesssim \|u(t)\|_{L^6}^{1/2}\|\nabla u(t)\|_{L^6}^{1/2} &\lesssim \|\nabla u(t)\|_{L^2} + \|\Delta u(t)\|_{L^2} \\
&\lesssim \langle t\rangle^{-\frac{1}{2}-\beta(p)} + \|\Delta u(t)\|_{L^2}, \\
\|B(t)\|_{L^\infty} \lesssim \|B(t)\|_{L^6}^{1/2}\|\nabla B(t)\|_{L^6}^{1/2} &\lesssim \|\nabla B(t)\|_{L^2} + \|\Delta B(t)\|_{L^2} \\
&\lesssim \langle t\rangle^{-\frac{1}{2}-\beta(p)} + \|\Delta B(t)\|_{L^2},
\end{aligned}$$

which along with (4.40) and $p \in (1, \frac{6}{5})$ yields

$$\int_{t_0}^{\infty} \|u(t)\|_{L^\infty} + \|B(t)\|_{L^\infty}\, dt \leq C.$$

Hence, the proof is completed. □

Proposition 4.6. Under the assumptions of Theorem 1.2, there holds

$$\|\nabla a(t)\|_{L^q} \leq C\,\|\nabla a_0\|_{L^q}, \tag{4.50}$$

and

$$\|a(t)\|_{\dot{H}^2} \leq C\,\|a_0\|_{\dot{H}^2}. \tag{4.51}$$

Proof. According to the transport equation in (1.3) and div$u = 0$, we know

$$\|\nabla a(t)\|_{L^q} \leq \|\nabla a_0\|_{L^q} + C\int_0^t \|\nabla u\|_{L^\infty}\|\nabla a\|_{L^q}\, d\tau,$$

for $q \in [1, \infty]$. Then Gronwall's inequality and Proposition 4.5 implies

$$\|\nabla a(t)\|_{L^q} \leq \|\nabla a_0\|_{L^q} e^{C\int_0^t \|\nabla u(t')\|_{L^\infty}\, dt'} \leq C\,\|\nabla a_0\|_{L^q}.$$

On the other hand, differentiating $\partial_t a + u\cdot\nabla a = 0$ twice with respect to the spatial variables, we get by a standard enery estimate that

$$\begin{aligned}
\|a(t)\|_{\dot{H}^2} &\leq \|a_0\|_{\dot{H}^2} + C\int_0^t \|\nabla u\|_{L^\infty}\|a\|_{\dot{H}^2} + \|\nabla a\|_{L^6}\|\nabla^2 u\|_{L^3}\, dt' \\
&\leq \|a_0\|_{\dot{H}^2} + C\int_0^t \|\nabla u\|_{L^\infty}\|a\|_{\dot{H}^2} \\
&\qquad + \|\Delta a\|_{L^2}(\|\Delta u\|_{L^2}^{2/3} + \|\Delta u\|_{L^6}^2)\, dt'.
\end{aligned}$$

Applying the Gronwall's inequality, Proposition 4.3, Proposition 4.5 and (4.47), we obtain

$$\begin{aligned}\|a(t)\|_{\dot{H}^2} &\le \|a_0\|_{\dot{H}^2} e^{C\int_0^t \|\nabla u\|_{L^\infty} + \|\Delta u\|_{L^2}^{2/3} + \|\Delta u\|_{L^6}^2\, dt'} \\ &\le C\|a_0\|_{\dot{H}^2}.\end{aligned}$$

Hence, the proof is completed. □

Proposition 4.7. Under the assumptions of Theorem 1.2, there holds

$$\begin{aligned}\sup_{t\ge t_0}\left(\|\nabla^2 u(t)\|_{L^2}^2 + \|\nabla^2 B(t)\|_{L^2}^2\right) + c_1\int_{t_0}^{\infty} \|\partial_t\nabla u(t)\|_{L^2}^2 + \|\partial_t \nabla B(t)\|_{L^2}^2\, dt \\ + c_2\int_{t_0}^{\infty} \|\nabla^3 u(t)\|_{L^2}^2 + \|\nabla^3 B(t)\|_{L^2}^2\, dt \le C.\end{aligned}$$

where C depend on m, M, $\|a_0\|_{L^2}$, $\|a_0\|_{L^\infty}$, $\|(u_0, B_0)\|_{L^p}$, $\|(u_0, B_0)\|_{H^2}$.

Proof. Taking spatial derivative to the momentum equation in (1.3), we have

$$\begin{aligned}\rho\partial_t\partial_j u^i + \partial_j\rho\cdot\partial_t u^i + \partial_j(\rho u)\cdot\nabla u^i \\ + \rho u\cdot\nabla\partial_j u^i - \Delta\partial_j u^i + \partial_j\partial_i\Pi = \partial_j(B\cdot\nabla B^i),\end{aligned}$$

with $i = 1, 2, 3$. Standard energy estimates yields

(4.52)
$$\begin{aligned}\|\sqrt{\rho}\partial_t\nabla u(t)\|_{L^2}^2 + \frac{1}{2}\frac{d}{dt}\|\nabla^2 u(t)\|_{L^2}^2 = -\int_{\mathbb{R}^3} \nabla\rho\,\partial_t u\,\partial_t\nabla u + \nabla(\rho u)\nabla u\,\partial_t\nabla u\, dx \\ - \int_{\mathbb{R}^3} (\rho u\cdot\nabla)\nabla u\,\partial_t\nabla u - \nabla(B\cdot\nabla B)\,\partial_t\nabla u\, dx.\end{aligned}$$

Similar argument gives

(4.53)
$$\|\partial_t\nabla B(t)\|_{L^2}^2 + \frac{1}{2}\frac{d}{dt}\|\nabla^2 B(t)\|_{L^2}^2 = -\int_{\mathbb{R}^3} \nabla(u\cdot\nabla B)\,\partial_t\nabla B - \nabla(B\cdot\nabla u)\,\partial_t\nabla B\, dx.$$

Summing (4.52), (4.53), (3.3) and Proposition 4.6, we obtain

$$\begin{aligned}&\frac{1}{2}\frac{d}{dt}\left(\|\nabla^2 u(t)\|_{L^2}^2 + \|\nabla^2 B(t)\|_{L^2}^2\right) + \|\sqrt{\rho}\partial_t\nabla u(t)\|_{L^2}^2 + \|\partial_t\nabla B(t)\|_{L^2}^2 \\ &\lesssim \|\partial_t u\|_{L^2}^2 + \|\nabla u\|_{L^2}^2\|\nabla^2 u\|_{L^2}\|\nabla^3 u\|_{L^2} + \|\nabla u\|_{L^2}^3\|\nabla^2 u\|_{L^2} \\ &\quad + \|u\|_{L^2}\|\nabla u\|_{L^2}\|\nabla^3 u\|_{L^2}^2 + \|\nabla B\|_{L^2}^2\|\nabla^2 B\|_{L^2}\|\nabla^3 B\|_{L^2} \\ &\quad + \|B\|_{L^2}\|\nabla B\|_{L^2}\|\nabla^3 B\|_{L^2}^2 + \|\nabla B\|_{L^2}^2\|\nabla^2 u\|_{L^2}\|\nabla^3 u\|_{L^2} \\ &\quad + \|u\|_{L^2}\|\nabla u\|_{L^2}\|\nabla^3 B\|_{L^2}^2 + \|B\|_{L^2}\|\nabla B\|_{L^2}\|\nabla^3 u\|_{L^2}^2.\end{aligned} \tag{4.54}$$

On the other hand, taking partial derivative to the momentum equation in (1.3) and multiplying $\frac{1}{\rho}\Delta\nabla u$ on both side, we can get

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\|\nabla^2 u(t)\|_{L^2}^2 + \|\frac{1}{\sqrt{\rho}}\Delta\nabla u\|_{L^2}^2 = \int_{\mathbb{R}^3} \nabla\rho\,\partial_t u\,\frac{1}{\rho}\,\Delta\nabla u\, dx \\ + \int_{\mathbb{R}^3} \nabla(\rho u)\,\nabla u\,\frac{1}{\rho}\,\Delta\nabla u\, dx + \int_{\mathbb{R}^3} \nabla(\nabla\Pi)\,\frac{1}{\rho}\,\Delta\nabla u\, dx \\ - \int_{\mathbb{R}^3} \nabla(B\cdot\nabla B)\,\frac{1}{\rho}\,\Delta\nabla u\, dx + \int_{\mathbb{R}^3} u\cdot\nabla(\nabla u)\,\Delta\nabla u\, dx\end{aligned} \tag{4.55}$$

by integrate over $\mathbb{R}^3$. Through similar estimate, we obtain

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\|\nabla^2 B\|_{L^2}^2 + \|\nabla^3 B\|_{L^2}^2 \\ = \int_{\mathbb{R}^3} \nabla(u\cdot\nabla B)\,\Delta\nabla B\,dx - \int_{\mathbb{R}^3}\nabla(B\cdot\nabla u)\,\Delta\nabla B\,dx.\end{aligned} \tag{4.56}$$

Combining (4.55) and (4.56), we have

$$\begin{aligned}&\frac{1}{2}\frac{d}{dt}\left(\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 B\|_{L^2}^2\right) + \|\frac{1}{\sqrt{\rho}}\nabla^3 u\|_{L^2}^2 + \|\nabla^3 B\|_{L^2}^2\\ &= \int_{\mathbb{R}^3}\nabla\rho\,\partial_t u\,\frac{1}{\rho}\,\Delta\nabla u\,dx + \int_{\mathbb{R}^3}\nabla(\rho u)\,\Delta u\,\frac{1}{\rho}\,\Delta\nabla u\,dx\\ &\quad + \int_{\mathbb{R}^3} u\cdot\nabla(\nabla u)\,\Delta\nabla u\,dx + \int_{\mathbb{R}^3}(1+a)\nabla(\nabla\Pi)\,\Delta\nabla u\,dx\\ &\quad - \int_{\mathbb{R}^3}\nabla(B\cdot\nabla B)\,\frac{1}{\rho}\,\Delta\nabla u\,dx + \int_{\mathbb{R}^3}\nabla(u\cdot\nabla B)\,\Delta\nabla B\,dx\\ &\quad - \int_{\mathbb{R}^3}\nabla(B\cdot\nabla u)\,\Delta\nabla B\,dx.\end{aligned} \tag{4.57}$$

Next, we give the estimation of each term on the right hand side of (4.57),

$$\int_{\mathbb{R}^3}\nabla\rho\,\partial_t u\,\frac{1}{\rho}\,\Delta\nabla u\,dx \lesssim \|1+a\|_{L^\infty}\|\nabla a\|_{L^\infty}\|\partial_t u\|_{L^2}\|\nabla^3 u\|_{L^2},$$

$$\begin{aligned}\int_{\mathbb{R}^3}\nabla(\rho u)\,\nabla u\,\frac{1}{\rho}\,\Delta\nabla u\,dx \lesssim &\|1+a\|_{L^\infty}\|\nabla a\|_{L^\infty}\|u\|_{L^2}^{1/2}\|\nabla u\|_{L^2}^{1/2}\|\nabla^2 u\|_{L^2}\|\nabla^3 u\|_{L^2}\\ &+\left(\|\nabla^2 u\|_{L^2}+\|\nabla^3 u\|_{L^2}\right)\|\nabla u\|_{L^2}\|\nabla^3 u\|_{L^2},\end{aligned}$$

$$\int_{\mathbb{R}^3}(u\cdot\nabla)\nabla u\,\Delta\nabla u\,dx \lesssim \|u\|_{L^2}^{1/2}\|\nabla u\|_{L^2}^{1/2}\|\nabla^3 u\|_{L^2}^2,$$

$$\begin{aligned}\int_{\mathbb{R}^3}\nabla(B\cdot\nabla B)\,\frac{1}{\rho}\,\Delta\nabla u\,dx \lesssim &\|1+a\|_{L^\infty}\|B\|_{L^2}^{1/2}\|\nabla B\|_{L^2}^{1/2}\|\nabla^3 B\|_{L^2}\|\nabla^3 u\|_{L^2}\\ &+\|1+a\|_{L^\infty}\left(\|\nabla^2 B\|_{L^2}+\|\nabla^3 B\|_{L^2}\right)\|\nabla B\|_{L^2}\|\nabla^3 u\|_{L^2},\end{aligned}$$

$$\begin{aligned}\int_{\mathbb{R}^3}\nabla(u\cdot\nabla B)\,\Delta\nabla B\,dx \lesssim &\|u\|_{L^2}^{1/2}\|\nabla u\|_{L^2}^{1/2}\|\nabla^3 B\|_{L^2}^2\\ &+\left(\|\nabla^2 u\|_{L^2}+\|\nabla^3 u\|_{L^2}\right)\|\nabla B\|_{L^2}\|\nabla^3 B\|_{L^2},\end{aligned}$$

$$\int_{\mathbb{R}^3}(1+a)\,\nabla(\nabla\Pi)\,\Delta\nabla u\,dx \lesssim \|1+a\|_{L^\infty}\|\nabla^2\Pi\|_{L^2}\|\nabla^3 u\|_{L^2}.$$

Moreover, using $\mathrm{div} u = 0$, we can get

$$
\begin{aligned}
&\|\Delta\nabla u(t)\|_{L^2} + \|\nabla(\nabla\Pi(t))\|_{L^2}\\
\le& \sqrt{2}\|\Delta\nabla u - \nabla(\nabla\Pi)\|_{L^2}\\
\le& \sqrt{2}\|\rho\partial_t\nabla u + \nabla\rho\,\partial_t u + \nabla(\rho u)\,\nabla u + \rho u\cdot\nabla(\nabla u) - \nabla(B\cdot\nabla B)\|_{L^2}\\
\le& \|\sqrt{\rho}\partial_t\nabla u\|_{L^2} + \|\partial_t u\|_{L^2} + \|u\|_{L^2}^{1/2}\|\nabla u\|_{L^2}^{1/2}\|\Delta u\|_{L^2} + \|B\|_{L^2}^{1/2}\|\nabla B\|_{L^2}^{1/2}\|\nabla^3 B\|_{L^2}\\
&+ \|u\|_{L^2}^{1/2}\|\nabla u\|_{L^2}^{1/2}\|\nabla^3 u\|_{L^2} + \|\nabla u\|_{L^2}\|\nabla^2 u\|_{L^2}^{\frac{1}{2}}\|\nabla^3 u\|_{L^2}^{\frac{1}{2}}\\
&+ \|\nabla B\|_{L^2}\|\nabla^2 B\|_{L^2}^{\frac{1}{2}}\|\nabla^3 B\|_{L^2}^{\frac{1}{2}}.
\end{aligned}
$$

Summing up all the above estimations in (4.57), then after a long and tedious calculations, we have

(4.58)
$$
\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\left(\|\nabla^2 u(t)\|_{L^2}^2 + \|\nabla^2 B(t)\|_{L^2}^2\right) + \|\frac{1}{\sqrt{\rho}}\nabla^3 u(t)\|_{L^2}^2 + \|\nabla^3 B(t)\|_{L^2}^2\\
\lesssim& \|\partial_t u\|_{L^2}^2 + \|u\|_{L^2}\|\nabla u\|_{L^2}\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 u\|_{L^2}^2\|\nabla u\|_{L^2}^2 + \|\nabla^2 B\|_{L^2}^2\|\nabla B\|_{L^2}^2\\
&+ \|\nabla B\|_{L^2}^2\|\nabla^2 u\|_{L^2}^2 + \|\nabla u\|_{L^2}^4\|\nabla^2 u\|_{L^2}^2 + \|\nabla B\|_{L^2}^4\|\nabla^2 B\|_{L^2}^2 + \|\sqrt{\rho}\partial_t\nabla u\|_{L^2}^2\\
&+ \|\nabla u\|_{L^2}^2\|\nabla^3 u\|_{L^2}^2 + \|u\|_{L^2}^{1/2}\|\nabla u\|_{L^2}^{1/2}\|\nabla^3 u\|_{L^2}^2 + \|B\|_{L^2}\|\nabla B\|_{L^2}\|\nabla^3 B\|_{L^2}^2\\
&+ \|\nabla B\|_{L^2}^2\|\nabla^3 B\|_{L^2}^2 + \|u\|_{L^2}^{1/2}\|\nabla u\|_{L^2}^{1/2}\|\nabla^3 B\|_{L^2}^2 + \|u\|_{L^2}\|\nabla u\|_{L^2}\|\nabla^3 u\|_{L^2}^2.
\end{aligned}
$$

We perform (4.54) + $\frac{1}{2}$(4.58) so that

(4.59)
$$
\begin{aligned}
&\frac{d}{dt}\left(\|\nabla^2\|_{L^2}^2 + \|\nabla^2 B\|_{L^2}^2\right) + \|\sqrt{\rho}\partial_t\nabla u\|_{L^2}^2 + \|\partial_t\nabla B\|_{L^2}^2 + \|\frac{1}{\sqrt{\rho}}\nabla^3 u\|_{L^2}^2 + \|\nabla^3 B\|_{L^2}^2\\
\lesssim& \|\partial_t u\|_{L^2}^2 + \left(\|\nabla u\|_{L^2}^4 + \|\nabla B\|_{L^2}^4\right)\left(\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 B\|_{L^2}^2\right) + \|\nabla u\|_{L^2}^3\|\nabla^2 u\|_{L^2}\\
&+ \|u\|_{L^2}\|\nabla u\|_{L^2}\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 u\|_{L^2}^2\|\nabla u\|_{L^2}^2 + \|\nabla^2 B\|_{L^2}^2\|\nabla B\|_{L^2}^2\\
&+ \|\nabla^2 u\|_{L^2}^2\|\nabla B\|_{L^2}^2 + \big(\|u\|_{L^2}\|\nabla u\|_{L^2} + \|\nabla u\|_{L^2} + \|B\|_{L^2}\|\nabla B\|_{L^2} + \|\nabla B\|_{L^2}^2\\
&+ \|u\|_{L^2}^{1/2}\|\nabla u\|_{L^2}^{1/2}\big)\left(\|\nabla^3 u\|_{L^2}^2 + \|\nabla^3 B\|_{L^2}^2\right).
\end{aligned}
$$

Note that we can take $\eta > 0$ in the proof of Proposition 4.1 smaller so that

$$
\|u_0\|_{L^2}\eta + \|B_0\|_{L^2}\eta + \|u_0\|_{L^2}^{1/2}\eta^{1/2} + \eta^2 \le \frac{c}{16C^2}
$$

where C is the constant on the right hand side of (4.59), c satisfies

$$
c\left(\|\nabla^3 u\|_{L^2}^2 + \|\nabla^3 B\|_{L^2}^2\right) \le \|\frac{1}{\sqrt{\rho}}\nabla^3 u\|_{L^2}^2 + \|\nabla^3 B\|_{L^2}^2.
$$

From the proof of Proposition 4.1, we know that $\tau^* = \infty$ as in Position 4.1. Hence, there exists two constant c_1 and c_2 such that

$$\begin{aligned}
&\frac{d}{dt}\left(\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 B\|_{L^2}^2\right) + c_1\left(\|\partial_t \nabla u\|_{L^2}^2 + \|\partial_t \nabla B\|_{L^2}^2\right) \\
&\qquad\qquad + c_2\left(\|\nabla^3 u\|_{L^2}^2 + \|\nabla^3 B\|_{L^2}^2\right) \\
\lesssim\ & \|\partial_t u\|_{L^2}^2 + \left(\|\nabla u\|_{L^2}^4 + \|\nabla B\|_{L^2}^4\right)\left(\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 B\|_{L^2}^2\right) \\
&+ \|\nabla u\|_{L^2}^3 \|\nabla^2 u\|_{L^2} + \|u\|_{L^2}\|\nabla u\|_{L^2}\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 u\|_{L^2}^2\|\nabla u\|_{L^2}^2 \\
&+ \|\nabla^2 B\|_{L^2}^2\|\nabla B\|_{L^2}^2 + \|\nabla^2 u\|_{L^2}^2\|\nabla B\|_{L^2}^2,
\end{aligned}$$

for $t \geq t_0$. Integrating the above inequality form t_0 to ∞, we then obtain

(4.60)

$$\begin{aligned}
&\sup_{t\geq t_0}\left(\|\nabla^2 u(t)\|_{L^2}^2 + \|\nabla^3 B(t)\|_{L^2}^2\right) + c_1\int_{t_0}^{\infty} \|\partial_t \nabla u\|_{L^2}^2 + \|\partial_t \nabla B(t)\|_{L^2}^2\, dt \\
&\qquad\qquad + c_2\int_{t_0}^{\infty} \|\nabla^3 u\|_{L^2}^2 + \|\nabla^3 B\|_{L^2}^2\, dt \\
\lesssim\ & \|\nabla^2 u(t_0)\|_{L^2}^2 + \|\nabla^2 B(t_0)\|_{L^2}^2 + \int_{t_0}^{\infty} \|\partial_t u\|_{L^2}^2\, dt \\
&+ \sup_{t\in[t_0,\infty]}\left(\|\nabla u\|_{L^2}^4 + \|\nabla B\|_{L^2}^4\right)\int_{t_0}^{\infty} \|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 B\|_{L^2}^2\, dt \\
&+ \sup_{t\in[t_0,\infty]} \|\nabla u(t)\|_{L^2}^3 \int_{t_0}^{\infty} \|\nabla^2 u\|_{L^2}\, dt + \sup_{t\in[t_0,\infty]} \|u(t)\|_{L^2}\|\nabla u(t)\|_{L^2}\int_{t_0}^{\infty} \|\nabla^2 u\|_{L^2}^2\, dt \\
&+ \sup_{t\in[t_0,\infty]}\left(\|\nabla u\|_{L^2}^2 + \|\nabla B(t)\|_{L^2}^2\right)\int_{t_0}^{\infty} \|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 B\|_{L^2}^2\, dt
\end{aligned}$$

At last by Proposition 4.3, the proof is completed. □

5. Global in Time Estimates for Reference Solutions

In this section, we prove the global in time estimates for the reference solution of (1.3). The proof will be based mainly on Theorem 1.2.

Proposition 5.1. Under the assumptions of Theorem 1.3, there holds

$$\|(\bar{u}, \bar{B})\|_{\tilde{L}^\infty(\mathbb{R}^+;\dot{B}^{1/2}_{2,1})} + \|(\bar{u}, \bar{B})\|_{L^1(\mathbb{R}^+;\dot{B}^{5/2}_{2,1})} \leq C, \tag{5.1}$$

for some C depending on the initial data.

Proof. According to imbedding theorems of Besov space and Theorem 1.2, we have

$$\begin{aligned}
\|\bar{u}\|_{\tilde{L}^\infty(\mathbb{R}^+;\dot{B}^{1/2}_{2,1})} &\lesssim \|\bar{u}\|_{L^\infty(\mathbb{R}^+;H^1)} \\
&\lesssim \|\bar{u}\|_{L^\infty(\mathbb{R}^+;L^2)} + \|\nabla\bar{u}\|_{L^\infty(\mathbb{R}^+;L^2)} \leq C.
\end{aligned}$$

Same considerations yields

$$\|\bar{B}\|_{\tilde{L}^\infty(\mathbb{R}^+;\dot{B}^{1/2}_{2,1})} \leq C.$$

By imbedding theorem of Besov space, we know that

$$\begin{aligned}
\|\bar{u}\|_{L^1(\mathbb{R}^+;\dot{B}^{5/2}_{2,1})} &\lesssim \|\bar{u}\|_{L^1(\mathbb{R}^+;\dot{H}^2)} + \|\bar{u}\|_{L^1(\mathbb{R}^+;\dot{H}^3)}, \\
\|\bar{B}\|_{L^1(\mathbb{R}^+;\dot{B}^{5/2}_{2,1})} &\lesssim \|\bar{B}\|_{L^1(\mathbb{R}^+;\dot{H}^2)} + \|\bar{B}\|_{L^1(\mathbb{R}^+;\dot{H}^3)}.
\end{aligned}$$

Rewrite the momentum equation in (1.3) as follows

$$-\Delta\bar{u}+\nabla\bar{\Pi}=-\frac{1}{1+\bar{a}}\,\partial_t\bar{u}-\frac{1}{1+\bar{a}}\,\bar{u}\cdot\nabla\bar{u}+\bar{B}\cdot\nabla\bar{B}.$$

From the elliptic estimate, we have

$$\begin{aligned}\|\bar{u}\|_{L^1(\mathbb{R}^+;\dot{H}^2)}\lesssim&\|\frac{1}{1+\bar{a}}\,\partial_t\bar{u}\|_{L^1(\mathbb{R}^+;L^2)}+\|\frac{1}{1+\bar{a}}\,\bar{u}\cdot\nabla\bar{u}\|_{L^1(\mathbb{R}^+;L^2)}\\&+\|\bar{B}\cdot\nabla\bar{B}\|_{L^1(\mathbb{R}^+;L^2)}\\\lesssim&\|\partial_t\bar{u}\|_{L^1(\mathbb{R}^+;L^2)}+\|\bar{u}\cdot\nabla\bar{u}\|_{L^1(\mathbb{R}^+;L^2)}+\|\bar{B}\cdot\nabla\bar{B}\|_{L^1(\mathbb{R}^+;L^2)}.\end{aligned}$$

Next, we estimate the three terms on the right hand side of the above inequality. Simple calculations yields

$$\begin{aligned}\int_{t_0}^{\infty}\|\partial_t\bar{u}\|_{L^2}\,dt\lesssim&\left(\int_{t_0}^{\infty}\langle t\rangle^{(1+2\beta(p))^-}\|\partial_t\bar{u}\|_{L^2}^2\,dt\right)^{1/2}\left(\int_{t_0}^{\infty}\langle t\rangle^{-(1+2\beta(p))^-}\,dt\right)^{1/2}\\\le&C,\end{aligned}$$

$$\begin{aligned}\int_{t_0}^{\infty}\|\bar{u}\cdot\nabla\bar{u}\|_{L^2}\,dt\lesssim&\int_{t_0}^{\infty}\|\bar{u}\|_{L^4}\|\nabla\bar{u}\|_{L^4}\,dt\\\lesssim&\int_{t_0}^{\infty}\|\bar{u}\|_{L^2}^{1/4}\|\nabla\bar{u}\|_{L^2}\|\nabla^2\bar{u}\|_{L^2}^{3/4}\,dt\\\lesssim&\|\bar{u}\|_{L^2}^{1/4}\left(\int_{t_0}^{\infty}\|\nabla\bar{u}\|_{L^2}^4\,dt\right)^{1/4}\left(\int_{t_0}^{\infty}\|\nabla^2\bar{u}\|_{L^2}\,dt\right)^{3/4}\\\le&C.\end{aligned}$$

$$\begin{aligned}\int_{t_0}^{\infty}\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}\,dt\lesssim&\|\bar{B}\|_{L^2}^{1/4}\left(\int_{t_0}^{\infty}\|\nabla\bar{B}\|_{L^2}^4\,dt\right)^{1/4}\left(\int_{t_0}^{\infty}\|\nabla^2\bar{B}\|_{L^2}\,dt\right)^{3/4}\\\le&C.\end{aligned}$$

Now we give the estimates about $\|\bar{u}\|_{L^1(\mathbb{R}^+;\dot{H}^3)}$,

$$\begin{aligned}\|\bar{u}\|_{L^1(\mathbb{R}^+;\dot{H}^3)}\lesssim&\left\|\nabla\left(\frac{1}{1+\bar{a}}\,\partial_t\bar{u}\right)\right\|_{L^1(\mathbb{R}^+;L^2)}+\left\|\nabla\left(\frac{1}{1+\bar{a}}\,\bar{u}\cdot\nabla\bar{u}\right)\right\|_{L^1(\mathbb{R}^+;L^2)}\\&+\left\|\nabla\left(\bar{B}\cdot\nabla\bar{B}\right)\right\|_{L^1(\mathbb{R}^+;L^2)}\\\lesssim&\|\nabla\bar{a}\cdot\partial_t\bar{u}\|_{L^1(\mathbb{R}^+;L^2)}+\|\partial_t\nabla\bar{u}\|_{L^1(\mathbb{R}^+;L^2)}+\|\nabla\bar{a}\cdot\bar{u}\cdot\nabla\bar{u}\|_{L^1(\mathbb{R}^+;L^2)}\\&+\|\nabla\bar{u}\cdot\nabla\bar{u}\|_{L^1(\mathbb{R}^+;L^2)}+\|\bar{u}\cdot\nabla(\nabla\bar{u})\|_{L^1(\mathbb{R}^+;L^2)}\\&+\|\nabla\bar{B}\cdot\nabla\bar{B}\|_{L^1(\mathbb{R}^+;L^2)}+\|\bar{B}\cdot\nabla(\nabla\bar{B})\|_{L^1(\mathbb{R}^+;L^2)}.\end{aligned}$$

Now, we give the estimates of the right hand side. According to Proposition 4.3 to Proposition 4.7, we have

$$\int_{t_0}^{\infty}\|\nabla\bar{a}\cdot\partial_t\bar{u}\|_{L^2}\,dt\lesssim\int_{t_0}^{\infty}\|\partial_t\bar{u}\|_{L^2}\,dt\le\,C,$$

$$\int_{t_0}^{\infty}\|\partial_t\nabla\bar{u}\|_{L^2}\,dt\le\,C,$$

$$
\begin{aligned}
&\int_{t_0}^{\infty} \|\nabla \bar{a}\cdot \bar{u}\cdot \nabla \bar{u}\|_{L^2}\,dt\\
\lesssim& \|\nabla \bar{a}\|_{L^\infty(\mathbb{R}^+;L^2)}\int_{t_0}^{\infty} \|\bar{u}\cdot \nabla \bar{u}\|_{L^2}\,dt\\
\lesssim& \|\nabla \bar{a}\|_{L^\infty(\mathbb{R}^+;L^2)}\int_{t_0}^{\infty} \|\bar{u}\|_{L^2}^{\frac{1}{2}}\|\nabla \bar{u}\|_{L^2}^{\frac{1}{2}}\|\nabla^2 \bar{u}\|_{L^2}\,dt\\
\lesssim& \|\nabla \bar{a}\|_{L^\infty(\mathbb{R}^+;L^2)}\|\bar{u}\|_{L^\infty(\mathbb{R}^+;L^2)}^{\frac{1}{2}}\|\nabla \bar{u}\|_{L^\infty(\mathbb{R}^+;L^2)}^{\frac{1}{2}}\int_{t_0}^{\infty} \|\nabla^2 \bar{u}\|_{L^2}\,dt\\
\leq& C,
\end{aligned}
$$

$$
\begin{aligned}
\int_{t_0}^{\infty} \|\nabla \bar{u}\cdot \nabla \bar{u}\|_{L^2}\,dt &\lesssim \int_{t_0}^{\infty} \|\nabla \bar{u}\|_{L^4}\|\nabla \bar{u}\|_{L^4}\,dt\\
&\lesssim \int_{t_0}^{\infty} \|\nabla \bar{u}\|_{L^2}^{1/4}\|\nabla^2 \bar{u}\|_{L^2}^{3/4}\|\nabla \bar{u}\|_{L^2}^{1/4}\|\nabla^2 \bar{u}\|_{L^2}^{3/4}\,dt\\
&\lesssim \int_{t_0}^{\infty} \|\nabla \bar{u}\|_{L^2}^{1/2}\|\nabla^2 \bar{u}\|_{L^2}^{3/2}\,dt \leq C,
\end{aligned}
$$

$$
\begin{aligned}
&\int_{t_0}^{\infty} \|\bar{u}\cdot \nabla(\nabla \bar{u})\|_{L^2}\,dt\\
\lesssim& \int_{t_0}^{\infty} \|\bar{u}\|_{L^\infty}\|\nabla^2 \bar{u}\|_{L^2}\,dt\\
\lesssim& \int_{t_0}^{\infty} \|\nabla \bar{u}\|_{L^2}\|\nabla^2 \bar{u}\|_{L^2}\,dt + \int_{t_0}^{\infty} \|\nabla^2 \bar{u}\|_{L^2}^2\,dt \leq C.
\end{aligned}
$$

Similar to the above estimates of $\bar{u}$, we can obtain

$$
\int_{t_0}^{\infty} \|\nabla \bar{B}\cdot \nabla \bar{B}\|_{L^2}\,dt \leq C \quad \text{and} \quad \int_{t_0}^{\infty} \|\bar{B}\cdot \nabla(\nabla \bar{B})\|_{L^2}\,dt \leq C.
$$

Summing up all the above estimates, we finally obtain

$$
\|\bar{u}\|_{L^1(\mathbb{R}^+;\dot{B}^{5/2}_{2,1})} \leq C.
$$

Using similar ideas as for $\bar{u}$, we have

$$
\begin{aligned}
\|\bar{B}\|_{L^1(\mathbb{R}^+;\dot{H}^2)} &\leq \|\partial_t \bar{B}\|_{L^1(\mathbb{R}^+;L^2)} + \|\bar{u}\cdot \nabla \bar{B}\|_{L^1(\mathbb{R}^+;L^2)} + \|\bar{B}\cdot \nabla \bar{u}\|_{L^1(\mathbb{R}^+;L^2)}\\
&\leq C,
\end{aligned}
$$

and

$$
\begin{aligned}
&\|\bar{B}\|_{L^1(\mathbb{R}^+;\dot{H}^3)}\\
\leq& \|\partial_t \nabla \bar{B}\|_{L^1(\mathbb{R}^+;L^2)} + \|\nabla\left(\bar{B}\cdot \nabla \bar{u}\right)\|_{L^1(\mathbb{R}^+;L^2)} + \|\nabla\left(\bar{u}\cdot \nabla \bar{B}\right)\|_{L^1(\mathbb{R}^+;L^2)}\\
\leq& C.
\end{aligned}
$$

So we also obtain that

$$
\|\bar{B}\|_{L^1(\mathbb{R}^+;\dot{B}^{5/2}_{2,1})} \leq C.
$$

Hence, finally the proof of Proposition 5.1 is completed. □

Proposition 5.2. Under the assumptions of Theorem 1.3, there hold

$$\|\bar{a}\|_{\tilde{L}^\infty(\mathbb{R}^+;B^{5/2}_{2,1})} \leq C \tag{5.2}$$

and

$$\|\bar{u}\|_{L^\infty(\mathbb{R}^+;L^p)} + \|\bar{u}\|_{\tilde{L}^\infty(\mathbb{R}^+;B^2_{2,1})} + \|\bar{u}\|_{L^1(\mathbb{R}^+;\dot{B}^4_{2,1})} + \|\nabla\bar{\Pi}\|_{L^1(\mathbb{R}^+;B^2_{2,1})} \leq C, \tag{5.3}$$

$$\|\bar{B}\|_{L^\infty(\mathbb{R}^+;L^p)} + \|\bar{B}\|_{\tilde{L}^\infty(\mathbb{R}^+;B^2_{2,1})} + \|\bar{B}\|_{L^1(\mathbb{R}^+;\dot{B}^4_{2,1})} \leq C. \tag{5.4}$$

Proof. Thanks to Proposition 5.1, we get by applying (2.3) to the transport equation in (1.2) that

$$\|\bar{a}\|_{\tilde{L}^\infty_t(B^{5/2}_{2,1})} \leq \|\bar{a}_0\|_{B^{5/2}_{2,1}} \exp\left\{C\int_0^t \|\bar{u}(\tau)\|_{\dot{B}^{5/2}_{2,1}}\,d\tau\right\} \leq C. \tag{5.5}$$

Next, let us turn to the estimates of $\bar{u}$ and $\bar{B}$. Indeed, from (4.13), we know that

$$\begin{aligned}\|\bar{u}\|_{L^\infty_t(L^p)} + \|\bar{B}\|_{L^\infty_t(L^p)} \lesssim{}& \|(u_0,B_0)\|_{L^p} + \int_0^t \|(u,B)\|_{L^2\cap L^3}\|(\nabla u,\nabla B)\|_{L^2}\,dt'\\ &+ \|\nabla\Pi\|_{L^1_t(L^p)}.\end{aligned}$$

Then by (4.14), Proposition 4.6, Theorem 1.2 and the above inequality, we deduce that

$$\|\bar{u}\|_{L^\infty(\mathbb{R}^+;L^p)} + \|\bar{B}\|_{L^\infty(\mathbb{R}^+;L^p)} \leq C. \tag{5.6}$$

On the other hand, applying Proposition 2.6 to the momentum and magnetic field equation of (1.2) ensures that

$$\begin{aligned}&\|\bar{u}\|_{\tilde{L}^\infty_T(\dot{B}^0_{2,1})} + \|\bar{u}\|_{\tilde{L}^1_T(\dot{B}^2_{2,1})} + \|\bar{B}\|_{\tilde{L}^\infty_T(\dot{B}^0_{2,1})} + \|\bar{B}\|_{\tilde{L}^1_T(\dot{B}^2_{2,1})}\\ &\lesssim \exp\left(C\int_0^T \|\bar{u}\|_{\dot{B}^{5/2}_{2,1}} + \|\bar{B}\|_{\dot{B}^{5/2}_{2,1}}\,dt\right)\Big(\|\bar{u}_0\|_{\dot{B}^0_{2,1}} + \|\bar{B}_0\|_{\dot{B}^0_{2,1}} + \|\bar{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^1_T(\dot{B}^0_{2,1})}\\ &\quad + \|\bar{a}\|_{\tilde{L}^\infty_T(\dot{H}^{3/2})}\|\nabla\bar{\Pi}\|_{L^1_T(L^2)} + \|\bar{a}\|_{L^\infty_T(\dot{H}^2)}\|\bar{u}\|_{L^1_T(\dot{H}^{3/2})}\Big).\end{aligned}$$

We know that

$$\begin{aligned}\|\bar{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^1_T(\dot{B}^0_{2,1})} &\lesssim \int_0^T \|\bar{a}\|_{\dot{B}^1_{2,1}}\|\bar{B}\cdot\nabla\bar{B}\|_{\dot{B}^{1/2}_{2,1}}\,dt\\ &\lesssim \int_0^T \|\bar{B}\|_{\dot{B}^{1/2}_{2,1}}\|\nabla\bar{B}\|_{\dot{B}^{3/2}_{2,1}}\,dt \leq C.\end{aligned}$$

Combining the above estimation, (5.5) and Proposition 4.5, we obtain

$$\|\bar{u}\|_{\tilde{L}^\infty_T(\dot{B}^0_{2,1})} + \|\bar{B}\|_{\tilde{L}^\infty_T(\dot{B}^0_{2,1})} \leq C. \tag{5.7}$$

Due to just a minor change of the proof in [21], we can get the following estimation. So, we postpone the proof in the appendix and give the estimation first

$$\begin{aligned}&\|\nabla\bar{u}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|\nabla\bar{B}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|\nabla\bar{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\\ &\qquad + \|\nabla\bar{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} + \|\nabla\bar{\Pi}\|_{L^1_t(\dot{B}^{3/2}_{2,1})} \leq C.\end{aligned} \tag{5.8}$$

Differentiating the momentum equation and magnetic field equation of (1.2) with respect to the spatial variables gives rise to

$$\begin{aligned}
&\partial_t\partial_i\bar{u}+\bar{u}\cdot\nabla\partial_i\bar{u}-\bar{B}\cdot\nabla\partial_i\bar{B}-(1+\bar{a})\Delta\partial_i\bar{u}+(1+\bar{a})\nabla\partial_i\Pi\\
&\qquad\qquad=-\partial_i\bar{u}\cdot\nabla\bar{u}+\partial_i\bar{a}\,\Delta\bar{u}-\partial_i\bar{a}\,\nabla\bar{\Pi}+\partial_i\bar{B}\cdot\nabla\bar{B}\\
&\qquad\qquad\quad+\partial_i\bar{a}\,\bar{B}\cdot\nabla\bar{B}+\bar{a}\,\partial_i\bar{B}\cdot\nabla\bar{B}+\bar{a}\,\bar{B}\cdot\nabla\partial_i\bar{B},\\
&\partial_t\partial_i\bar{B}-\Delta\partial_i\bar{B}+\bar{u}\cdot\nabla\partial_i\bar{B}-\bar{B}\cdot\nabla\partial_i\bar{u}=\partial_i\bar{B}\cdot\nabla\bar{u}-\partial_i\bar{u}\cdot\nabla\bar{B}.
\end{aligned}\tag{5.9}$$

Using Remark 2.9 and Gronwall's inequality, we will obtain

$$\begin{aligned}
&\|\nabla\bar{u}\|_{\tilde{L}^\infty_T(\dot{B}^1_{2,1})}+\|\nabla\bar{u}\|_{\tilde{L}^1_T(\dot{B}^3_{2,1})}+\|\nabla\bar{B}\|_{\tilde{L}^\infty_T(\dot{B}^1_{2,1})}+\|\nabla\bar{B}\|_{\tilde{L}^1_T(\dot{B}^3_{2,1})}\\
&\lesssim\exp\left\{C\int_0^T\|\bar{u}\|_{\dot{B}^{5/2}_{2,1}}+\|\bar{B}\|_{\dot{B}^{5/2}_{2,1}}\,dt\right\}\Big\{\|\nabla\bar{u}_0\|_{\dot{B}^1_{2,1}}+\|\nabla\bar{B}_0\|_{\dot{B}^1_{2,1}}\\
&\quad+\|\bar{a}\|_{\tilde{L}^\infty_T(\dot{B}^2_{2,1})}\Big(\|\bar{u}\|_{L^1_T(\dot{B}^{7/2}_{2,1})}+\|\Pi\|_{L^1_T(\dot{B}^{5/2}_{2,1})}+\|\bar{B}\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})}\|\bar{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\Big)\\
&\quad+\|\bar{a}\|_{\tilde{L}^\infty_t(\dot{B}^2_{2,1})}\|\bar{B}\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})}\|\bar{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}+\|\bar{a}\|_{\tilde{L}^\infty_T(\dot{B}^1_{2,1})}\|\bar{B}\|_{\tilde{L}^\infty_T(\dot{B}^{3/2}_{2,1})}\|\nabla\bar{B}\|_{L^1_T(\dot{B}^{5/2}_{2,1})}\Big\}.
\end{aligned}$$

Using Proposition 5.1, (5.5), (5.8) and product laws in Besov space, we get

$$\begin{aligned}
&\|\nabla\bar{u}\|_{\tilde{L}^\infty_T(\dot{B}^1_{2,1})}+\|\nabla\bar{u}\|_{\tilde{L}^1_T(\dot{B}^3_{2,1})}+\|\nabla\bar{B}\|_{\tilde{L}^\infty_T(\dot{B}^1_{2,1})}+\|\nabla\bar{B}\|_{\tilde{L}^1_T(\dot{B}^3_{2,1})}\\
&\le C+C\left(\|\bar{u}\|_{L^1_T(\dot{B}^{7/2}_{2,1})}+\|\bar{B}\|_{L^1_T(\dot{B}^{7/2}_{2,1})}+\|\bar{\Pi}\|_{L^1_T(\dot{B}^{5/2}_{2,1})}\right)\\
&\le C+C\left(\|\Delta\bar{u}\|_{L^1_T(L^2)}+\|\Delta\bar{B}\|_{L^1_T(L^2)}+\|\nabla\bar{\Pi}\|_{L^1_T(L^2)}\right)\\
&\quad+\eta\Big(\|\bar{u}\|_{L^1_T(\dot{B}^4_{2,1})}+\|\bar{B}\|_{L^1_T(\dot{B}^4_{2,1})}+\|\bar{\Pi}\|_{L^1_T(\dot{B}^3_{2,1})}\Big).
\end{aligned}\tag{5.10}$$

Notice that $\mathrm{div}\bar{u}=0$, we get by taking div to (5.9) that

$$\begin{aligned}
\mathrm{div}\left((1+\bar{a})\nabla\partial_i\bar{\Pi}\right)=&-\mathrm{div}\partial_i\left[(\bar{u}\cdot\nabla)\bar{u}\right]+\mathrm{div}\partial_i\left[\bar{a}\,\Delta\bar{u}\right]-\mathrm{div}\left[\partial_i\bar{a}\,\nabla\bar{\Pi}\right]\\
&+\mathrm{div}\partial_i\left(\bar{B}\cdot\nabla\bar{B}\right)+\mathrm{div}\left(\partial_i\bar{a}\,\bar{B}\cdot\nabla\bar{B}\right)\\
&+\mathrm{div}\left(\bar{a}\,\partial_i\bar{B}\cdot\nabla\bar{B}\right)+\mathrm{div}\left(\bar{a}\,\bar{B}\cdot\nabla\partial_i\bar{B}\right)
\end{aligned}$$

From this and (2.5), we deduce that

$$\begin{aligned}
\|\nabla^2\bar{\Pi}\|_{L^1_t(\dot{B}^1_{2,1})}\lesssim&\|\partial_i\left(\bar{u}\cdot\nabla\bar{u}\right)\|_{L^1_t(\dot{B}^1_{2,1})}+\|\partial_i\left(\bar{a}\,\Delta\bar{u}\right)\|_{L^1_t(\dot{B}^1_{2,1})}\\
&+\|\partial_i\bar{a}\,\nabla\bar{\Pi}\|_{L^1_t(\dot{B}^1_{2,1})}+\|\partial_i\left(\bar{B}\cdot\nabla\bar{B}\right)\|_{L^1_t\dot{B}^1_{2,1}}\\
&+\|\partial_i\bar{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^1_t(\dot{B}^1_{2,1})}+\|\bar{a}\,\partial_i\bar{B}\cdot\nabla\bar{B}\|_{L^1_t\dot{B}^1_{2,1}}\\
&+\|\bar{a}\,\bar{B}\cdot\nabla\partial_i\bar{B}\|_{L^1_t\dot{B}^1_{2,1}}+\|\bar{a}\|_{\tilde{L}^\infty_t(\dot{H}^2)}\|\nabla^2\bar{\Pi}\|_{\tilde{L}^1_t(\dot{H}^{1/2})}.
\end{aligned}$$

Applying the product laws in Besov space yields that for any $\epsilon > 0$

$$
\begin{aligned}
&\|\nabla^2\bar{\Pi}\|_{L^1_t(\dot{B}^1_{2,1})}\\
\lesssim&\|\nabla\bar{u}\|_{L^\infty_t(\dot{B}^1_{2,1})}\|\nabla\bar{u}\|_{L^1_t(\dot{B}^{3/2}_{2,1})}+\|\bar{u}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\nabla^2\bar{u}\|_{L^1_t(\dot{B}^1_{2,1})}\\
&+\|\nabla\bar{B}\|_{L^\infty_t(\dot{B}^1_{2,1})}\|\nabla\bar{B}\|_{L^1_t(\dot{B}^{3/2}_{2,1})}+\|\bar{B}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\nabla^2\bar{B}\|_{L^1_t(\dot{B}^1_{2,1})}\\
&+\|\nabla\bar{a}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\Delta\bar{u}\|_{L^1_t(\dot{B}^1_{2,1})}+\|\bar{a}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\nabla^3\bar{u}\|_{L^1_t(\dot{B}^1_{2,1})}\\
&+\|\nabla\bar{a}\|_{\tilde{L}^\infty_t(\dot{B}^1_{2,1})}\|\bar{B}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\nabla\bar{B}\|_{L^1_t(\dot{B}^{3/2}_{2,1})}\\
&+\|\bar{a}\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})}\|\nabla\bar{B}\|_{L^\infty_t(\dot{B}^1_{2,1})}\|\nabla\bar{B}\|_{L^1_t(\dot{B}^{3/2}_{2,1})}\\
&+\|\bar{a}\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})}\|\bar{B}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\nabla^2\bar{B}\|_{L^1_t(\dot{B}^1_{2,1})}\\
&+\|\bar{a}\|_{\tilde{L}^\infty_t(\dot{B}^2_{2,1})}\Big(\epsilon\|\nabla\bar{\Pi}\|_{L^1_t(\dot{B}^2_{2,1})}+C\,\|\nabla\bar{\Pi}\|_{L^1_t(L^2)}\Big)
\end{aligned}
\tag{5.11}
$$

Combining (5.8), Proposition 5.1 and taking $\epsilon > 0$ in (5.11) small enough, we will have

$$
\begin{aligned}
\|\nabla\bar{\Pi}\|_{L^1_t(\dot{B}^2_{2,1})}\leq& C\,\big(1+\|\nabla\bar{u}\|_{L^\infty_t(\dot{B}^1_{2,1})}+\|\nabla\bar{B}\|_{L^\infty_t(\dot{B}^1_{2,1})}\\
&+\|\bar{u}\|_{L^1_t(\dot{B}^4_{2,1})}+\|\bar{B}\|_{L^1_t(\dot{B}^4_{2,1})}\big).
\end{aligned}
\tag{5.12}
$$

Finally, taking η in (5.10) small enough and substituting (5.12) into (5.10), we get

$$
\|\nabla\bar{u}\|_{\tilde{L}^\infty_t(\dot{B}^1_{2,1})}+\|\nabla\bar{u}\|_{\tilde{L}^1_t(\dot{B}^3_{2,1})}+\|\nabla\bar{B}\|_{\tilde{L}^\infty_t(\dot{B}^1_{2,1})}+\|\nabla\bar{B}\|_{\tilde{L}^1_t(\dot{B}^3_{2,1})}\leq C.
$$

This along with (5.6) and (5.7) completes the proof of the proposition. □

6. Stability of the Global Large Solutions

In this section we will give the proof of Theorem 1.3. Denoting $\tilde{u} := u - \bar{u}$, $\tilde{B} := B - \bar{B}$ and $\tilde{a} := a - \bar{a}$, we have

$$
\begin{cases}
\partial_t\tilde{a}+(\bar{u}+\tilde{u})\nabla\tilde{a}=-\tilde{u}\cdot\nabla\bar{a},\\
\partial_t\tilde{u}+\tilde{u}\cdot\nabla\tilde{u}+\tilde{u}\cdot\nabla\bar{u}+\bar{u}\cdot\nabla\tilde{u}-(1+\bar{a}+\tilde{a})(\Delta\tilde{u}-\nabla\tilde{\Pi})\\
\qquad\qquad -(1+\bar{a}+\tilde{a})(\tilde{B}\cdot\nabla\tilde{B})-(1+\bar{a}+\tilde{a})(\tilde{B}\cdot\nabla\bar{B}+\bar{B}\cdot\nabla\tilde{B})\\
\qquad\qquad\qquad\qquad =\tilde{a}(\Delta\bar{u}-\nabla\bar{\Pi})+\tilde{a}\,(\bar{B}\cdot\nabla\bar{B}),\\
\partial_t\tilde{B}-\Delta\tilde{B}+\tilde{u}\cdot\nabla\tilde{B}-\tilde{B}\cdot\nabla\tilde{u}=\tilde{B}\cdot\nabla\bar{u}+\bar{B}\nabla\tilde{u}-\tilde{u}\nabla\bar{B}-\bar{u}\cdot\nabla\tilde{B},\\
\operatorname{div}\tilde{u}=\operatorname{div}\tilde{B}=0,\\
(\tilde{a},\tilde{u},\tilde{B})|_{t=0}=(\tilde{a}_0,\tilde{u}_0,\tilde{B}_0).
\end{cases}
\tag{6.1}
$$

Then the proof of Theorem 1.3 is equivalent to the proof of the global well posedness of (6.1) with small enough initial data $(\tilde{a}_0,\tilde{u}_0)$. Indeed, according to the coupled parabolic hyperbolic theory [22], it is standard to prove that there exists a positive time $\tilde{T}^*$ such that (1.2) with initial data $(\bar{a}_0+\tilde{a}_0,\bar{u}_0+\tilde{u}_0,\bar{B}_0+\tilde{B}_0)$ has a unique solution (a,u,B) with

$$
\begin{aligned}
&a\in C([0,\tilde{T}^*);B^{7/2}_{2,1}(\mathbb{R}^3)),\\
&u\in C([0,\tilde{T}^*);B^2_{2,1})\cap L^1_{\rm loc}((0,\tilde{T}^*);\dot{B}^4_{2,1}(\mathbb{R}^3)),\\
&B\in C([0,\tilde{T}^*);B^2_{2,1})\cap L^1_{\rm loc}((0,\tilde{T}^*);\dot{B}^4_{2,1}(\mathbb{R}^3)).
\end{aligned}
$$

Then $(\tilde{a},\tilde{u},\tilde{B})$ with

$$\begin{aligned}&\tilde{a}\in C([0,\tilde{T}^*);B^{7/2}_{2,1}(\mathbb{R}^3)),\\&\tilde{u}\in C([0,\tilde{T}^*);B^{2}_{2,1})\cap L^1_{\rm loc}((0,\tilde{T}^*);\dot{B}^{4}_{2,1}(\mathbb{R}^3)),\\&\tilde{B}\in C([0,\tilde{T}^*);B^{2}_{2,1})\cap L^1_{\rm loc}((0,\tilde{T}^*);\dot{B}^{4}_{2,1}(\mathbb{R}^3)).\end{aligned}$$

solves (6.1) on $[0,\tilde{T}^*)$. Without loss of generality, we may assume that $\tilde{T}^*$ is the maximal time of the existence to this solution. The aim of what follows is to prove that $\tilde{T}^*=\infty$ and $(\tilde{a},\tilde{u},\tilde{B})$ remains small for all $t>0$.

Lemma 6.1. *Let*

$$V(t)=2\int_0^t\left(\|\nabla\bar{u}(\tau)\|_{L^\infty}+\|\nabla\bar{B}(\tau)\|_{L^\infty}\right)\,d\tau.$$

Then under the assumption of Theorem 1.3, we have

$$\begin{aligned}&\frac{d}{dt}\Big[e^{-V(t)}\Big(\|(\sqrt{\rho}\tilde{u}(t)\|^2_{L^2}+\|\tilde{B}(t)\|^2_{L^2}\Big)\Big]\\&\qquad\qquad+c_0g^2(t)e^{-V(t)}\Big(\|\sqrt{\rho}\tilde{u}(t)\|^2_{L^2}+\|\tilde{B}(t)\|^2_{L^2}\Big)\\&\le Ce^{-V(t)}\bigg\{g^2(t)\int_{S_1(t)}e^{-2t|\xi|^2}|\hat{\tilde{u}}_0(\xi)|^2\,d\xi+g^2(t)\int_{S_2(t)}e^{-2t|\xi|^2}|\hat{\tilde{B}}_0(\xi)|^2\,d\xi\\&\quad+\|\tilde{\rho}\|^2_{L^3}\|\Delta\bar{u}-\nabla\bar{\Pi}\|^2_{L^2}+\|\tilde{\rho}\|^2_{L^3}\|\bar{B}\cdot\nabla\bar{B}\|^2_{L^2}\\&\quad+g(t)^7\Big(\|\tilde{u}\|^2_{L^2_t(L^2)}+\|\tilde{B}\|^2_{L^2_t(L^2)}\Big)\Big(\|\bar{u}\|^2_{L^2_t(L^2)}+\|\bar{B}\|^2_{L^2_t(L^2)}\Big)\\&\quad+g(t)^7\Big(\|\tilde{B}\|^2_{L^2_t(L^2)}+\|\tilde{u}\|^2_{L^2_t(L^2)}\Big)^2\\&\quad+g(t)^5\|\tilde{B}\|^2_{L^2_t(L^2)}\Big(\|\nabla\bar{B}\|^2_{L^2_t(L^2)}+\|\nabla\tilde{B}\|^2_{L^2_t(L^2)}\Big)\\&\quad+g(t)^5\Big(\|\Delta\tilde{u}\|^2_{L^1_t(L^2)}+\|\nabla\tilde{\Pi}\|^2_{L^1_t(L^2)}+\|\tilde{\rho}\|^2_{L^\infty_t(L^2)}\|\Delta\bar{u}-\nabla\bar{\Pi}\|^2_{L^1_t(L^2)}\Big)\\&\quad+g(t)^5\|\tilde{\rho}\|^2_{L^\infty_t(L^2)}\|\bar{B}\cdot\nabla\bar{B}\|^2_{L^1_t(L^2)}\bigg\},\end{aligned}\tag{6.2}$$

for $t<\tilde{T}^*$, where the time dependent phase space region $S_1(t)$, $S_2(t)$ is given as in the proof of Proposition 4.3. Here and in what follows, we shall always denote

$$\rho:=\frac{1}{1+\bar{a}+\tilde{a}},\quad \bar{\rho}:=\frac{1}{1+\bar{a}},\quad \tilde{\rho}:=\rho-\bar{\rho}.$$

Proof. Thanks to (6.1), $(\rho,\tilde{u},\tilde{B})$ solves

$$\begin{cases}\partial_t\rho+\operatorname{div}(\rho u)=0,\\ \rho\partial_t\tilde{u}+\rho u\cdot\nabla\tilde{u}-B\cdot\nabla\tilde{B}+\rho\tilde{u}\cdot\nabla\bar{u}-\tilde{B}\cdot\nabla\bar{B}-\Delta\tilde{u}+\nabla\tilde{\Pi}\\ \qquad\qquad=-\frac{\tilde{\rho}}{\rho}(\Delta\bar{u}-\nabla\bar{\Pi})-\frac{\tilde{\rho}}{\rho}\bar{B}\cdot\nabla\bar{B},\\ \partial_t\tilde{B}-\Delta\tilde{B}+u\cdot\nabla\tilde{B}-B\cdot\nabla\tilde{u}=\tilde{B}\cdot\nabla\bar{u}-\tilde{u}\cdot\nabla\bar{B},\\ \operatorname{div}u=\operatorname{div}B=0,\end{cases}\tag{6.3}$$

from which we get by a standard energy estimate that

$$\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\left(\|\sqrt{\rho}\tilde{u}(t)\|_{L^2}^2+\|\tilde{B}(t)\|_{L^2}^2\right)+\left(\|\nabla\tilde{u}(t)\|_{L^2}^2+\|\nabla\tilde{B}(t)\|_{L^2}^2\right)\\
=&-\int_{\mathbb{R}^3}\rho\,\tilde{u}\cdot\nabla\bar{u}\,\tilde{u}\,dx+\int_{\mathbb{R}^3}\tilde{B}\cdot\nabla\bar{u}\,\tilde{B}\,dx-\int_{\mathbb{R}^3}\frac{\tilde{\rho}}{\bar{\rho}}(\Delta\bar{u}-\nabla\bar{\Pi})\,\tilde{u}\,dx\\
&-\int_{\mathbb{R}^3}\frac{\tilde{\rho}}{\bar{\rho}}\,\bar{B}\cdot\nabla\bar{B}\,\tilde{u}\,dx\\
\le&\|\nabla\bar{u}\|_{L^\infty}\int_{\mathbb{R}^3}\rho|\tilde{u}|^2+|\tilde{B}|^2\,dx+C\|\tilde{\rho}\|_{L^3}\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\\
&+C\|\tilde{\rho}\|_{L^3}\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}.
\end{aligned}$$

This gives

(6.4)
$$\begin{aligned}
&\frac{d}{dt}\left(e^{-V(t)}\left(\|\sqrt{\rho}\tilde{u}(t)\|_{L^2}^2+\|\tilde{B}(t)\|_{L^2}^2\right)\right)+e^{-V(t)}\left(\|\nabla\tilde{u}(t)\|_{L^2}^2+\|\nabla\tilde{B}(t)\|_{L^2}^2\right)\\
\le&Ce^{-V(t)}\|\tilde{\rho}\|_{L^3}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+Ce^{-V(t)}\|\tilde{\rho}\|_{L^3}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2,
\end{aligned}$$

which along with a similar derivation of (4.17) ensures

$$\begin{aligned}
&\frac{d}{dt}\left(e^{-V(t)}\left(\|\sqrt{\rho}\tilde{u}(t)\|_{L^2}^2+\|\tilde{B}(t)\|_{L^2}^2\right)\right)\\
&\qquad\qquad+c_0g^2(t)e^{-V(t)}\left(\|\sqrt{\rho}\tilde{u}(t)\|_{L^2}^2+\|\tilde{B}(t)\|_{L^2}^2\right)\\
\le&Ce^{-V(t)}\|\tilde{\rho}\|_{L^3}^2\left(\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\right)\\
&\qquad+Ce^{-V(t)}g^2(t)\left(\int_{S_1(t)}|\hat{\tilde{u}}(\xi)|^2\,d\xi+\int_{S_2(t)}|\hat{\tilde{B}}(\xi)|^2\,d\xi\right),
\end{aligned}\tag{6.5}$$

with $c_0\le\min(\frac{1}{\rho},1)$ and the time dependent phase space region $S_1(t)$, $S_2(t)$ being the same as the one in (4.17).

Now, we need to give the estimate of $\int_{S_1(t)}|\hat{\tilde{u}}(\xi)|^2\,d\xi$ and $\int_{S_2(t)}|\hat{\tilde{B}}(\xi)|^2\,d\xi$. Rewrite the second equation in (6.3) as

$$\begin{aligned}
\tilde{u}(t)=&e^{t\Delta}\tilde{u}_0+\int_0^t e^{(t-\tau)\Delta}\mathbb{P}\Big(-\nabla\cdot(\tilde{u}\otimes\tilde{u}+\bar{u}\otimes\tilde{u}+\tilde{u}\otimes\bar{u})\\
&+\left(\frac{1}{\rho}-1\right)\Delta\tilde{u}-\left(\frac{1}{\rho}-1\right)\nabla\tilde{\Pi}-\frac{\tilde{\rho}}{\rho\bar{\rho}}(\Delta\bar{u}-\nabla\bar{\Pi})\\
&+\frac{1}{\rho}B\cdot\nabla\tilde{B}+\frac{1}{\rho}\tilde{B}\cdot\nabla\bar{B}-\frac{\tilde{\rho}}{\rho\bar{\rho}}\,\bar{B}\cdot\nabla\bar{B}\Big)\,d\tau.
\end{aligned}$$

Taking the Fourier transformation with respect to the x variables and integrating the resulting equation over $S_1(t)$, we obtain

$$
\begin{aligned}
&\int_{S_1(t)} |\hat{\tilde{u}}(\xi)|^2\,d\xi\\
&\lesssim \int_{S_1(t)} e^{-2t|\xi|^2}|\hat{\tilde{u}}_0(\xi)|^2\,d\xi + g(t)^5\bigg(\int_0^t \|\mathcal{F}(\tilde{u}\otimes\tilde{u})\|_{L^\infty_\xi}\\
&\quad + \|\mathcal{F}(\bar{u}\otimes\tilde{u})\|_{L^\infty_\xi}\,d\tau\bigg)^2 + g(t)^3\bigg(\int_0^t \left\|\mathcal{F}\left(\left(\frac{1}{\rho}-1\right)\Delta\tilde{u}\right)\right\|_{L^\infty_\xi}\\
&\quad + \left\|\mathcal{F}\left(\left(\frac{1}{\rho}-1\right)\Delta\tilde{\Pi}\right)\right\|_{L^\infty_\xi} + \left\|\mathcal{F}\left(\frac{\tilde{\rho}}{\rho\bar{\rho}}(\Delta\bar{u}-\nabla\bar{\Pi})\right)\right\|_{L^\infty_\xi}\,d\tau\bigg)^2\\
&\quad + g(t)^3\bigg(\int_0^t \left\|\mathcal{F}\left(\frac{1}{\rho}\nabla\cdot\left(\tilde{B}\otimes\tilde{B}\right)\right)\right\|_{L^\infty_\xi} + \left\|\mathcal{F}\left(\frac{1}{\rho}\nabla\cdot\left(\bar{B}\otimes\tilde{B}\right)\right)\right\|_{L^\infty_\xi}\\
&\quad + \left\|\mathcal{F}\left(\frac{\tilde{\rho}}{\rho\bar{\rho}}\,\bar{B}\cdot\nabla\bar{B}\right)\right\|_{L^\infty_\xi}\,d\tau\bigg)^2.
\end{aligned}
\tag{6.6}
$$

Notice that

$$\int_0^t \|\mathcal{F}(\tilde{u}\otimes\tilde{u})(\tau)\|_{L^\infty_\xi}\,d\tau \le C\int_0^t \|\tilde{u}(\tau)\|_{L^2}^2\,d\tau,$$

$$\int_0^t \|\mathcal{F}(\tilde{u}\otimes\bar{u})(\tau)\|_{L^\infty_\xi}\,d\tau \le C\left(\int_0^t \|\tilde{u}(\tau)\|_{L^2}^2\right)^{\frac{1}{2}}\left(\int_0^t \|\bar{u}(\tau)\|_{L^2}^2\right)^{\frac{1}{2}},$$

$$\int_0^t \left\|\mathcal{F}\left(\left(\frac{1}{\rho}-1\right)\Delta\tilde{u}\right)(\tau)\right\|_{L^\infty_\xi}\,d\tau \le C\int_0^t \|\Delta\tilde{u}\|_{L^2}\,d\tau,$$

$$\int_0^t \left\|\mathcal{F}\left(\left(\frac{1}{\rho}-1\right)\nabla\tilde{\Pi}\right)\right\|_{L^\infty_\xi} \le C\int_0^t \|\nabla\tilde{\Pi}\|_{L^2}\,d\tau,$$

$$\int_0^t \left\|\mathcal{F}\left(\frac{\tilde{\rho}}{\bar{\rho}\rho}\left(\Delta\bar{u}-\nabla\bar{\Pi}\right)\right)\right\|_{L^\infty_\xi}\,d\tau \le C\|\tilde{\rho}\|_{L^\infty_t(L^2)}\int_0^t \|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}\,d\tau,$$

$$
\begin{aligned}
\int_0^t \left\|\mathcal{F}\left(\frac{1}{\rho}\nabla\left(\tilde{B}\otimes\tilde{B}\right)\right)\right\|_{L^\infty_\xi}\,d\tau &\le C\int_0^t \|\frac{1}{\rho}\nabla\cdot\left(\tilde{B}\otimes\tilde{B}\right)\|_{L^1}\,d\tau\\
&\le C\left(\int_0^t \|\tilde{B}(\tau)\|_{L^2}^2\,d\tau\right)^{\frac{1}{2}}\left(\int_0^t \|\nabla\tilde{B}(\tau)\|_{L^2}^2\,d\tau\right)^{\frac{1}{2}},
\end{aligned}
$$

$$\int_0^t \left\|\mathcal{F}\left(\frac{1}{\rho}\nabla\left(\tilde{B}\otimes\bar{B}\right)\right)\right\|_{L^\infty_\xi}\,d\tau \le C\left(\int_0^t \|\tilde{B}(\tau)\|_{L^2}^2\,d\tau\right)^{\frac{1}{2}}\left(\int_0^t \|\nabla\bar{B}(\tau)\|_{L^2}^2\,d\tau\right)^{\frac{1}{2}},$$

$$\int_0^t \left\|\frac{\tilde{\rho}}{\rho\bar{\rho}}\,\bar{B}\,\nabla\bar{B}\right\|_{L^1}\,d\tau \le C\,\|\tilde{\rho}\|_{L^\infty_t(L^2)}\int_0^t \|\bar{B}\cdot\nabla\bar{B}\|_{L^2}\,d\tau.$$

Then, we rewrite the third equation in (6.3) as

$$\begin{aligned}\tilde{B}(t) =& e^{t\Delta}\tilde{B}_0 + \int_0^t e^{(t-\tau)\Delta}\Big[\nabla\cdot\Big(\tilde{B}\otimes\bar{u} + \bar{B}\otimes\tilde{u} - \tilde{u}\otimes\bar{B}\\ &- \bar{u}\otimes\tilde{B} + \tilde{B}\otimes\tilde{u} - \tilde{u}\otimes\tilde{B}\Big)\Big]\,d\tau.\end{aligned}$$

Taking the Fourier transformation with respect to the x variables and integrating the resulting equation over $S(t)$, we obtain

$$\begin{aligned}\int_{S_2(t)} |\hat{\tilde{B}}(\xi)|^2\,d\xi \lesssim& \int_{S_2} e^{-2t|\xi|^2}|\hat{\tilde{B}}_0|^2\,d\xi\\ &+ g(t)^5\bigg(\int_0^t \|\mathcal{F}(\tilde{B}\otimes\bar{u})\|_{L^\infty_\xi} + \|\mathcal{F}(\bar{B}\otimes\tilde{u})\|_{L^\infty_\xi}\\ &+ \|\mathcal{F}(\tilde{u}\otimes\bar{B})\|_{L^\infty_\xi} + \|\mathcal{F}(\bar{u}\otimes\tilde{B})\|_{L^\infty_\xi}\\ &+ \|\mathcal{F}(\tilde{B}\otimes\tilde{u})\|_{L^\infty_\xi} + \|\mathcal{F}(\tilde{u}\otimes\tilde{B})\|_{L^\infty_\xi}\bigg)^2.\end{aligned}\tag{6.7}$$

Similar to the estimates used above, we have

$$\begin{aligned}\int_{S_2(t)} |\hat{\tilde{B}}(\xi)|^2\,d\xi \lesssim& \int_{S_2(t)} e^{-2t|\xi|^2}|\hat{\tilde{B}}_0(\xi)|^2\,d\xi\\ &+ g(t)^5\left(\|\tilde{u}\|^2_{L^2_t(L^2)} + \|\tilde{B}\|^2_{L^2_t(L^2)}\right)\left(\|\bar{u}\|^2_{L^2_t(L^2)} + \|\bar{B}\|^2_{L^2_t(L^2)}\right)\\ &+ g(t)^5\left(\|\tilde{B}\|^2_{L^2_t(L^2)} + \|\tilde{u}\|^2_{L^2_t(L^2)}\right)^2\end{aligned}\tag{6.8}$$

Combining all the above estimates, we complete the proof. □

Lemma 6.2. *Let*

$$\begin{aligned}U(t) :=& C\int_0^t \Big(\|\Delta\bar{u}(\tau)\|_{L^2}\|\nabla\bar{u}(\tau)\|_{L^2} + \|\bar{u}(\tau)\|^2_{L^\infty}\\ &+ \|\Delta\bar{B}(\tau)\|_{L^2}\|\nabla\bar{B}(\tau)\|_{L^2} + \|\bar{B}(\tau)\|^2_{L^\infty}\Big)\,d\tau.\end{aligned}$$

If

$$\sup_{t\in[0,\bar{T})}\Big[\|\tilde{u}(t)\|_{L^2}\|\nabla\tilde{u}(t)\|_{L^2} + \|\tilde{B}(t)\|_{L^2}\|\nabla\tilde{B}(t)\|_{L^2}\Big] \le \nu\tag{6.9}$$

for some $\bar{T}\le\tilde{T}^*$ and some sufficiently small positive constant ν, then under the assumptions of Theorem 1.3, we have

$$\begin{aligned}&\frac{d}{dt}\Big(e^{-U(t)}\|(\nabla\tilde{u},\nabla\tilde{B})\|^2_{L^2}\Big) + e^{-U(t)}\|(\sqrt{\rho}\partial_t\tilde{u},\partial_t\tilde{B})\|^2_{L^2}\\ &\qquad + c_0e^{-U(t)}\|(\Delta\tilde{u},\Delta\tilde{B})\|^2_{L^2}\\ &\le C\,e^{-U(t)}\left(\|\tilde{\rho}\|^2_{L^\infty}\|\Delta\bar{u}-\nabla\bar{\Pi}\|^2_{L^2} + \|\tilde{\rho}\|^2_{L^\infty}\|\bar{B}\cdot\nabla\bar{B}\|^2_{L^2}\right)\qquad \text{for } t<\bar{T}.\end{aligned}\tag{6.10}$$

Proof. Taking the L^2 inner product between the second equation of (6.3) and $\partial_t\tilde{u}$, we obtain

$$\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\|\nabla\tilde{u}\|_{L^2}^2+\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}^2\\
\lesssim&\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}\|\sqrt{\rho}\|_{L^\infty}\left(\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^2}+\|\tilde{u}\cdot\nabla\bar{u}\|_{L^2}+\|\bar{u}\cdot\nabla\tilde{u}\|_{L^2}\right)\\
&+\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}\left\|\frac{\tilde{\rho}}{\bar{\rho}\sqrt{\rho}}\right\|_{L^\infty}\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}+\left\|\frac{\tilde{\rho}}{\bar{\rho}}\right\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\\
&+\|\tilde{B}\cdot\nabla\tilde{B}\|_{L^2}^2+\|\tilde{B}\cdot\nabla\bar{B}\|_{L^2}^2+\|\bar{B}\cdot\nabla\tilde{B}\|_{L^2}^2.
\end{aligned}$$

Notice that

$$\begin{aligned}
&\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^2}+\|\tilde{u}\cdot\nabla\bar{u}\|_{L^2}+\|\bar{u}\cdot\nabla\tilde{u}\|_{L^2}\\
\lesssim&\|\tilde{u}\|_{L^2}^{\frac{1}{2}}\|\nabla\tilde{u}\|_{L^2}^{\frac{1}{2}}\|\Delta\tilde{u}\|_{L^2}+\|\nabla\tilde{u}\|_{L^2}\left(\|\nabla\bar{u}\|_{L^2}^{\frac{1}{2}}\|\Delta\bar{u}\|_{L^2}^{\frac{1}{2}}+\|\bar{u}\|_{L^\infty}\right),\\
&\|\tilde{B}\cdot\nabla\tilde{B}\|_{L^2}+\|\tilde{B}\cdot\nabla\bar{B}\|_{L^2}+\|\bar{B}\cdot\nabla\tilde{B}\|_{L^2}\\
\lesssim&\|\tilde{B}\|_{L^2}^{\frac{1}{2}}\|\nabla\tilde{B}\|_{L^2}^{\frac{1}{2}}\|\Delta\tilde{B}\|_{L^2}+\|\nabla\tilde{B}\|_{L^2}\left(\|\nabla\bar{B}\|_{L^2}^{\frac{1}{2}}\|\Delta\bar{B}\|_{L^2}^{\frac{1}{2}}+\|\bar{B}\|_{L^\infty}\right),
\end{aligned}\tag{6.11}$$

which ensures

$$\begin{aligned}
&\frac{d}{dt}\|\nabla\tilde{u}\|_{L^2}^2+\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}^2\\
\lesssim&\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\Delta\tilde{u}\|_{L^2}^2+\|\nabla\tilde{u}\|_{L^2}^2\Big(\|\Delta\bar{u}\|_{L^2}\|\nabla\bar{u}\|_{L^2}+\|\bar{u}\|_{L^\infty}^2\Big)\\
&\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\Delta\tilde{B}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big(\|\Delta\bar{B}\|_{L^2}\|\nabla\bar{B}\|_{L^2}+\|\bar{B}\|_{L^\infty}^2\Big)\\
&+\|\tilde{\rho}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2.
\end{aligned}\tag{6.12}$$

By taking the L^2 inner product between (6.3) and $\Delta\tilde{u}$, we get

$$\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\|\nabla\tilde{u}\|_{L^2}^2+\|\rho\|_{L^\infty}^{-1}\|\Delta\tilde{u}\|_{L^2}^2\\
\le&\|\Delta\tilde{u}\|_{L^2}\Big(\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^2}+\|\tilde{u}\cdot\nabla\bar{u}\|_{L^2}+\|\bar{u}\cdot\nabla\tilde{u}\|_{L^2}\\
&+\|\rho^{-1}\|_{L^\infty}(\|\tilde{B}\cdot\nabla\tilde{B}\|_{L^2}+\|\bar{B}\cdot\nabla\tilde{B}\|_{L^2}+\|\tilde{B}\cdot\nabla\bar{B}\|_{L^2})\\
&\left\|\frac{\tilde{\rho}}{\rho\bar{\rho}}\right\|_{L^\infty}\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}+\|\rho^{-1}\|_{L^\infty}\|\nabla\bar{\Pi}\|_{L^2}+\left\|\frac{\tilde{\rho}}{\rho\bar{\rho}}\right\|_{L^\infty}\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}\Big).
\end{aligned}$$

Thanks to the momentum equation of (6.3) and $\operatorname{div}\tilde{u}=0$, we have

$$\begin{aligned}
\|\nabla\tilde{\Pi}\|_{L^2}+\|\Delta\tilde{u}\|_{L^2}\lesssim&\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^2}+\|\tilde{u}\cdot\nabla\bar{u}\|_{L^2}+\|\bar{u}\cdot\nabla\tilde{u}\|_{L^2}\\
&+\|\tilde{B}\cdot\nabla\tilde{B}\|_{L^2}+\|\tilde{B}\cdot\nabla\bar{B}\|_{L^2}+\|\bar{B}\cdot\nabla\tilde{B}\|_{L^2}\\
&+\|\tilde{\rho}\|_{L^\infty}\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}+\|\tilde{\rho}\|_{L^\infty}\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}\\
&+\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}.
\end{aligned}\tag{6.13}$$

This gives

$$
\begin{aligned}
&\frac{d}{dt}\|\nabla\tilde{u}\|_{L^2}^2+c_0\|\Delta\tilde{u}\|_{L^2}^2\\
\lesssim&\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\Delta\tilde{u}\|_{L^2}^2+\|\nabla\tilde{u}\|_{L^2}^2\Big(\|\nabla\bar{u}\|_{L^2}\|\Delta\bar{u}\|_{L^2}+\|\bar{u}\|_{L^\infty}^2\Big)\\
&\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\Delta\tilde{B}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big(\|\nabla\bar{B}\|_{L^2}\|\Delta\bar{B}\|_{L^2}+\|\bar{B}\|_{L^\infty}^2\Big)\\
&\|\tilde{\rho}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}^2+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2.
\end{aligned}\tag{6.14}
$$

Combining (6.12) and (6.14), we arrive at

$$
\begin{aligned}
&\frac{d}{dt}\|\nabla\tilde{u}\|_{L^2}^2+\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}^2+\left(c_0-C\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\right)\|\Delta\tilde{u}\|_{L^2}^2\\
\lesssim&\|\nabla\tilde{u}\|_{L^2}^2(\|\Delta\bar{u}\|_{L^2}\|\nabla\bar{u}\|_{L^2}+\|\bar{u}\|_{L^\infty}^2)+\|\tilde{\rho}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\\
&+\|\nabla\tilde{B}\|_{L^2}^2(\|\Delta\bar{B}\|_{L^2}\|\nabla\bar{B}\|_{L^2}+\|\bar{B}\|_{L^\infty}^2)+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\\
&+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\Delta\tilde{B}\|_{L^2}^2
\end{aligned}\tag{6.15}
$$

for $t<\tilde{T}^*$.

Taking the L^2 inner product between the third equation of (6.3) and $\partial_t\tilde{B}$, we have

$$
\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\|\nabla\tilde{B}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\\
\lesssim&\|\tilde{u}\cdot\nabla\tilde{B}\|_{L^2}^2+\|\bar{u}\cdot\nabla\tilde{B}\|_{L^2}^2+\|\bar{B}\cdot\nabla\tilde{u}\|_{L^2}^2+\|\tilde{B}\cdot\nabla\tilde{u}\|_{L^2}^2\\
&+\|\tilde{B}\cdot\nabla\bar{u}\|_{L^2}^2+\|\tilde{u}\cdot\nabla\bar{B}\|_{L^2}^2.
\end{aligned}
$$

Noticing that

$$
\begin{aligned}
&\|\tilde{u}\cdot\nabla\tilde{B}\|_{L^2}+\|\bar{u}\cdot\nabla\tilde{B}\|_{L^2}+\|\tilde{u}\cdot\nabla\bar{B}\|_{L^2}\\
&\qquad\qquad+\|\bar{B}\cdot\nabla\tilde{u}\|_{L^2}+\|\tilde{B}\cdot\nabla\tilde{u}\|_{L^2}+\|\tilde{B}\cdot\nabla\bar{u}\|_{L^2}\\
\lesssim&\|\tilde{u}\|_{L^2}^{\frac{1}{2}}\|\nabla\tilde{u}\|_{L^2}^{\frac{1}{2}}\|\Delta\tilde{B}\|_{L^2}+\|\nabla\tilde{u}\|_{L^2}\|\nabla\bar{B}\|_{L^2}^{\frac{1}{2}}\|\Delta\bar{B}\|_{L^2}^{\frac{1}{2}}\\
&+\|\tilde{B}\|_{L^2}^{\frac{1}{2}}\|\nabla\tilde{B}\|_{L^2}^{\frac{1}{2}}\|\Delta\tilde{u}\|_{L^2}+\|\nabla\tilde{B}\|_{L^2}\|\nabla\bar{u}\|_{L^2}^{\frac{1}{2}}\|\Delta\bar{u}\|_{L^2}^{\frac{1}{2}}\\
&+\|\nabla\tilde{B}\|_{L^2}\|\bar{u}\|_{L^\infty}+\|\nabla\tilde{u}\|_{L^2}\|\bar{B}\|_{L^\infty}
\end{aligned}
$$

which ensures

$$
\begin{aligned}
&\frac{d}{dt}\|\nabla\tilde{B}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\\
\lesssim&\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\Delta\tilde{B}\|_{L^2}^2+\|\nabla\tilde{u}\|_{L^2}^2\|\nabla\bar{B}\|_{L^2}\|\Delta\bar{B}\|_{L^2}\\
&\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\Delta\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\|\nabla\bar{u}\|_{L^2}\|\Delta\bar{u}\|_{L^2}\\
&+\|\nabla\tilde{B}\|_{L^2}^2\|\bar{u}\|_{L^\infty}^2+\|\nabla\tilde{u}\|_{L^2}^2\|\bar{B}\|_{L^\infty}^2.
\end{aligned}\tag{6.16}
$$

By taking the L^2 inner product between (6.3) and $\Delta\tilde{B}$, we get

$$
\begin{aligned}
\frac{1}{2}\frac{d}{dt}\|\nabla\tilde{B}\|_{L^2}^2+\|\Delta\tilde{B}\|_{L^2}^2\le&\|\Delta\tilde{B}\|_{L^2}\Big(\|\tilde{u}\cdot\nabla\tilde{B}\|_{L^2}+\|\bar{u}\cdot\nabla\tilde{B}\|_{L^2}\\
&+\|\tilde{B}\cdot\nabla\tilde{u}\|_{L^2}+\|\bar{B}\cdot\nabla\tilde{u}\|_{L^2}\\
&+\|\tilde{B}\cdot\nabla\bar{u}\|_{L^2}+\|\tilde{u}\cdot\nabla\bar{B}\|_{L^2}\Big).
\end{aligned}
$$

Combining this and (6.16), we arrive at

$$\begin{aligned}
&\frac{d}{dt}\|\nabla\tilde{B}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2+(1-C\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2})\|\Delta\tilde{B}\|_{L^2}^2\\
\lesssim&\|\nabla\tilde{u}\|_{L^2}^2\|\nabla\bar{B}\|_{L^2}\|\Delta\bar{B}\|_{L^2}+\|\nabla\tilde{B}\|_{L^2}^2\|\bar{u}\|_{L^\infty}^2+\|\nabla\tilde{u}\|_{L^2}^2\|\bar{B}\|_{L^\infty}^2\\
&+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\Delta\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\|\nabla\bar{u}\|_{L^2}\|\Delta\bar{u}\|_{L^2}.
\end{aligned}\tag{6.17}$$

Hence, by (6.15) and (6.17), we obtain that

$$\begin{aligned}
&\frac{d}{dt}\Big(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big)+\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\\
&\quad+(c_0-C\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}-C\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2})\|\Delta\tilde{u}\|_{L^2}^2\\
&\quad+(1-C\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}-C\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2})\|\Delta\tilde{B}\|_{L^2}^2\\
\lesssim&\Big(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big)\Big(\|\Delta\bar{u}\|_{L^2}\|\nabla\bar{u}\|_{L^2}+\|\Delta\bar{B}\|_{L^2}\|\nabla\bar{B}\|_{L^2}\\
&+\|\bar{u}\|_{L^\infty}^2+\|\bar{B}\|_{L^\infty}^2\Big)+\|\tilde{\rho}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2
\end{aligned}$$

for $t<\tilde{T}^*$. For small enough ν, we obtain

$$\begin{aligned}
&\frac{d}{dt}\Big(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big)+\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\\
&\qquad\qquad+c_0\|\Delta\tilde{u}\|_{L^2}^2+\|\Delta\tilde{B}\|_{L^2}^2\\
\lesssim&\Big(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big)\Big(\|\Delta\bar{u}\|_{L^2}\|\nabla\bar{u}\|_{L^2}+\|\Delta\bar{B}\|_{L^2}\|\nabla\bar{B}\|_{L^2}\\
&+\|\bar{u}\|_{L^\infty}^2+\|\bar{B}\|_{L^\infty}^2\Big)+\|\tilde{\rho}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2.
\end{aligned}$$

Then simple calculations leads to

$$\begin{aligned}
&\frac{d}{dt}\left(e^{-U(t)}\left(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\right)\right)+e^{-U(t)}\left(\|\sqrt{\rho}\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\right)\\
&\qquad\qquad+e^{-U(t)}\left(c_0\|\Delta\tilde{u}\|_{L^2}^2+\|\Delta\tilde{B}\|_{L^2}^2\right)\\
\le&C\,e^{-U(t)}\|\tilde{\rho}\|_{L^\infty}^2\Big(\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\Big)
\end{aligned}$$

for $t<\bar{T}$. □

Proposition 6.3. Under the assumptions in Theorem 1.3, there exist constants C and c so that if

$$\delta_0:=\|\tilde{u}_0\|_{H^1}+\|\tilde{u}_0\|_{L^p}+\|\tilde{B}_0\|_{H^1}+\|\tilde{B}_0\|_{L^p}+\|\tilde{\rho}_0\|_{L^2}+\|\tilde{\rho}_0\|_{L^\infty}<c,$$

then there hold

$$\begin{aligned}
&\|\tilde{u}(t)\|_{L^2}+\|\tilde{B}(t)\|_{L^2}\le C\delta_0\langle t\rangle^{-\beta(p)},\\
&\|\nabla\tilde{u}\|_{L^2}+\|\nabla\tilde{B}\|_{L^2}\le C\delta_0\langle t\rangle^{-\frac{1}{2}-\beta(p)}\quad\text{for } t<\tilde{T}^*,\\
&\int_0^{\tilde{T}^*}\|\Delta\tilde{u}\|_{L^2}+\|\Delta\tilde{B}\|_{L^2}+\|\nabla\tilde{\Pi}\|_{L^2}\,dt\le C\delta_0,\\
&\int_0^{\tilde{T}^*}\|\tilde{u}\|_{L^\infty}+\|\tilde{B}\|_{L^\infty}\,dt\le C\delta_0,
\end{aligned}\tag{6.18}$$

and

$$\int_0^{\tilde{T}^*} \left(\|(\partial_t \tilde{u}, \partial_t \tilde{B})\|_{L^2}^2 + \|(\Delta \tilde{u}, \Delta \tilde{B})\|_{L^2}^2 + \|\nabla \tilde{\Pi}\|_{L^2}^2 \right) \langle t \rangle^{(1+2\beta(p))^-} \, d\tau \le C\delta_0 \tag{6.19}$$

for $\beta(p) = \frac{3}{4}\left(\frac{2}{p} - 1\right)$.

Proof. Let $\xi(t) := \sup_{t' \in [0,t]} \left(\|\tilde{\rho}(t)\|_{L^2} + \|\tilde{\rho}(t)\|_{L^\infty} \right)$. Integrating (6.4) and (6.10) over $[0,t]$ for $t \le \bar{T}$ that

$$\begin{aligned} e^{-V(t)} \left(\|\tilde{u}\|_{L^2}^2 + \|\tilde{B}\|_{L^2}^2 \right) \lesssim{}& \|\tilde{u}_0\|_{L^2}^2 + \|\tilde{B}_0\|_{L^2}^2 \\ &+ \int_0^t e^{-V(t')} \|\tilde{\rho}\|_{L^3}^2 \|\Delta \bar{u} - \nabla \bar{\Pi}\|_{L^2}^2 \, dt' \\ &+ \int_0^t e^{-V(t')} \|\tilde{\rho}\|_{L^3}^2 \|\bar{B} \cdot \nabla \bar{B}\|_{L^2}^2 \, dt'. \end{aligned}$$

and

$$\begin{aligned} & e^{-U(t)} \|(\nabla \tilde{u}, \nabla \tilde{B})\|_{L^2}^2 + \int_0^t e^{-U(t')} \|(\sqrt{\rho}\partial_t \tilde{u}, \partial_t \tilde{B})\|_{L^2}^2 + c_0 e^{-U(t')} \|(\Delta \tilde{u}, \Delta \tilde{B})\|_{L^2}^2 \, dt' \\ \lesssim{}& \|(\nabla \tilde{u}_0, \nabla \tilde{B}_0)\|_{L^2}^2 + \int_0^t e^{-V(t')} \|\Delta \bar{u} - \nabla \bar{\Pi}\|_{L^2}^2 \, dt' + \int_0^t e^{-V(t')} \|\bar{B} \cdot \nabla \bar{B}\|_{L^2}^2 \, dt', \end{aligned}$$

where $V(t)$ and $U(t)$ defined as in Lemma 6.1 and 6.2 respectively. From the decay properties of the reference solution, we have

$$\begin{aligned} & \|\tilde{u}(t)\|_{L^2} + \|\tilde{B}(t)\|_{L^2} + \|\nabla \tilde{u}\|_{L_t^2(L^2)} + \|\nabla \tilde{B}\|_{L_t^2(L^2)} \\ \lesssim{}& \|(\tilde{u}_0, \tilde{B}_0)\|_{L^2} + \sup_{t' \in [0,t]} \|\rho(t')\|_{L^3} \le C \left(\delta_0 + \xi(t) \right), \\ & \|\nabla \tilde{u}(t)\|_{L^2} + \|\nabla \tilde{B}(t)\|_{L^2} + \|\sqrt{\rho}\partial_t \tilde{u}\|_{L_t^2(L^2)} + \|\partial_t \tilde{B}\|_{L_t^2(L^2)} \\ & \quad + \|\Delta \tilde{u}\|_{L_t^2(L^2)} + \|\Delta \tilde{B}\|_{L_t^2(L^2)} \le C \left(\delta_0 + \xi(t) \right). \end{aligned} \tag{6.20}$$

Now, we use Schonbek's strategy in [23] to prove (6.18) and (6.19).

Step 1: Rough decay estimates of $\|\tilde{u}(t)\|_{L^2}$, $\|\nabla \tilde{u}(t)\|_{L^2}$ and $\|\tilde{B}(t)\|_{L^2}$, $\|\nabla \tilde{B}(t)\|_{L^2}$. Let $S_1(t) := \left\{ \xi : |\xi| \le \sqrt{\frac{2}{c_0}} g(t) \right\}$, $S_2(t) := \left\{ \xi : |\xi| \le \sqrt{2} g(t) \right\}$ with $g(t)$ satisfying $g(t) \lesssim \langle t \rangle^{-\frac{1}{2}}$, which will be chosen later on. Then thanks to Lemma 6.1, (6.20), (4.19) and (4.20), we obtain for $t \le \bar{T}$

$$\begin{aligned} & \frac{d}{dt} \left[e^{-V(t)} \left(\|\sqrt{\rho}\tilde{u}\|_{L^2}^2 + \|\tilde{B}\|_{L^2}^2 \right) \right] + c_0 g^2(t) e^{-V(t)} \left(\|\sqrt{\rho}\tilde{u}\|_{L^2}^2 + \|\tilde{B}\|_{L^2}^2 \right) \\ \le{}& C \Big[g^2(t) \delta_0^2 \langle t \rangle^{-2\beta(p)} + \xi(t)^2 \|\Delta \bar{u} - \nabla \bar{\Pi}\|_{L^2}^2 + \xi(t)^2 \|\bar{B} \cdot \nabla \bar{B}\|_{L^2}^2 \\ & + g(t)^5 \left(\delta_0 + \xi(t) \right)^2 \langle t \rangle \Big] \end{aligned}$$

provided that $\xi(t)\le 1$ for $t\le \bar{T}$, which will be justified later on. In what follows, we shall always assume that $t\le \bar{T}$. Then, we have

$$\begin{aligned}
&e^{c_0\int_0^t g(t')^2\,dt'}e^{-V(t)}\Big(\|\sqrt{\rho}\tilde{u}\|_{L^2}^2+\|\tilde{B}\|_{L^2}^2\Big)\\
\lesssim&\|(\sqrt{\rho_0}\tilde{u}_0,\tilde{B}_0)\|_{L^2}^2+\delta_0^2\int_0^t e^{c_0\int_0^{t'} g(\tau)^2\,d\tau}\langle t\rangle^{-1-2\beta(p)}\,dt'\\
&+\xi(t)^2\int_0^t e^{c_0\int_0^{t'} g(\tau)^2\,d\tau}\left(\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\right)\,dt'\\
&+(\delta_0+\xi(t))^2\int_0^t e^{c_0\int_0^{t'} g(\tau)^2\,d\tau}\langle t\rangle^{-\frac{3}{2}}\,dt'.
\end{aligned}$$

Taking $g^2(t)=\frac{\alpha}{c_0(1+t)}$ with $\frac{1}{2}<\alpha<1+2\beta(p)$ in the above inequality gives

$$\|\tilde{u}(t)\|_{L^2}+\|\tilde{B}(t)\|_{L^2}\le C(\delta_0+\xi(t))\langle t\rangle^{-\frac{1}{4}}, \tag{6.21}$$

Applying a similar procedure to (6.10) ensures that

$$\begin{aligned}
&\frac{d}{dt}\Big[e^{-U(t)}\Big(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big)\Big]+2g^2(t)e^{-U(t)}\Big(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big)\\
&\qquad+e^{-U(t)}\Big(\|\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\Big)\\
\lesssim&g^4(t)e^{-U(t)}\int_{S_1(t)}|\hat{\tilde{u}}(\xi)|^2\,d\xi+g^4(t)e^{-U(t)}\int_{S_2(t)}|\hat{\tilde{B}}(\xi)|^2\,d\xi\\
&+e^{-U(t)}\|\tilde{\rho}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+e^{-U(t)}\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2
\end{aligned}$$

which together with (6.6), (6.7) implies that

$$\begin{aligned}
&\frac{d}{dt}\Big[e^{-U(t)}\Big(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big)\Big]+2g^2(t)e^{-U(t)}\Big(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\Big)\\
&\qquad+e^{-U(t)}\Big(\|\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\Big)\\
\lesssim&e^{-U(t)}\bigg\{g^4(t)\int_{S_1(t)}e^{-2t|\xi|^2}|\hat{\tilde{u}}_0(\xi)|^2\,d\xi+g^4(t)\int_{S_2(t)}e^{-2t|\xi|^2}|\hat{\tilde{B}}_0|^2\,d\xi\\
&+g(t)^9\Big(\|\tilde{u}\|_{L^2_t(L^2)}^2+\|\tilde{B}\|_{L^2_t(L^2)}^2\Big)\Big(\|\bar{u}\|_{L^2_t(L^2)}^2+\|\bar{B}\|_{L^2_t(L^2)}^2\Big)\\
&+g(t)^9\Big(\|\tilde{B}\|_{L^2_t(L^2)}^2+\|\tilde{u}\|_{L^2_t(L^2)}^2\Big)^2\\
&+g(t)^7\|\tilde{B}\|_{L^2_t(L^2)}^2\Big(\|\nabla\bar{B}\|_{L^2_t(L^2)}^2+\|\nabla\tilde{B}\|_{L^2_t(L^2)}^2\Big)\\
&+g(t)^7\Big(\|\Delta\tilde{u}\|_{L^1_t(L^2)}^2+\|\nabla\tilde{\Pi}\|_{L^1_t(L^2)}^2+\|\tilde{\rho}\|_{L^\infty_t(L^2)}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^1_t(L^2)}^2\Big)\\
&+g(t)^7\|\tilde{\rho}\|_{L^\infty_t(L^2)}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^1_t(L^2)}^2+\|\tilde{\rho}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\\
&+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\bigg\}
\end{aligned}\tag{6.22}$$

Substituting (6.21) into (7.17), we arrive at

$$\begin{aligned}
&\frac{d}{dt}\left(e^{-U(t)}\left(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\right)\right)+2e^{-U(t)}g^2(t)\left(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\right)\\
&\qquad\qquad+e^{-U(t)}\left(\|\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\right)\\
&\lesssim(\delta_0+\xi(t))^2\langle t\rangle^{-\frac{5}{2}}+\xi(t)^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\xi(t)^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2,
\end{aligned}$$

from which we deduce that

$$\begin{aligned}
&e^{2\int_0^t g(\tau)^2\,d\tau}e^{-U(t)}\left(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\right)\\
&\qquad\qquad+\int_0^t e^{2\int_0^{t'} g(\tau)^2\,d\tau}e^{-U(t')}\left(\|\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\right)dt'\\
\lesssim&\|\nabla\tilde{u}_0\|_{L^2}^2+\|\nabla\tilde{B}_0\|_{L^2}^2+\xi(t)^2\int_0^t e^{2\int_0^{t'} g(\tau)^2\,d\tau}\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\,dt'\\
&+\xi(t)^2\int_0^t e^{2\int_0^{t'} g(\tau)^2\,d\tau}\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\,dt'\\
&+(\delta_0+\xi(t))^2\int_0^t e^{2\int_0^{t'} g(\tau)^2\,d\tau}\langle t\rangle^{-\frac{5}{2}}\,dt'.
\end{aligned}$$

Taking $g^2(t)=\frac{\alpha}{2(1+t)}$ with $\alpha>1$ in the above inequality yields

$$\begin{aligned}
&\langle t\rangle^{\alpha}\left(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\right)+\int_0^t\langle t\rangle^{\alpha}\left(\|\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\right)dt'\\
\lesssim&\delta_0^2+\xi(t)^2\int_0^t\langle t'\rangle^{\alpha}\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\,dt'+\xi(t)^2\int_0^t\langle t'\rangle^{\alpha}\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\,dt'\\
&+\Big(\delta_0+\xi(t)\Big)^2\int_0^t\langle t'\rangle^{\alpha-\frac{5}{2}}\,dt'.
\end{aligned}\tag{6.23}$$

Taking $\alpha\in(\frac{3}{2},1+2\beta(p))$ in (6.23) yields

$$\|\nabla\tilde{u}(t)\|_{L^2}+\|\nabla\tilde{B}(t)\|_{L^2}\lesssim(\delta_0+\xi(t))\langle t\rangle^{-\frac{3}{4}}.\tag{6.24}$$

Step 2 : Improved decay estimates of $\|\tilde{u}(t)\|_{L^2}$, $\|\nabla\tilde{u}(t)\|_{L^2}$ and $\|\tilde{B}(t)\|_{L^2}$, $\|\nabla\tilde{B}(t)\|_{L^2}$. On the other hand, taking $\alpha\in[1,\frac{3}{2})$ in (6.23), we obtain

$$\int_0^t\langle t'\rangle^{\left(\frac{3}{2}\right)^-}\left(\|\partial_t\tilde{u}(t')\|_{L^2}^2+\|\partial_t\tilde{B}(t')\|_{L^2}^2\right)dt'\lesssim(\delta_0+\xi(t))^2\tag{6.25}$$

which implies

$$\int_0^t\|\partial_t\tilde{u}(t')\|_{L^2}+\|\partial_t\tilde{B}(t')\|_{L^2}\,dt'\lesssim\delta_0+\xi(t).\tag{6.26}$$

Next, we give the estimates about $\|\Delta\tilde{u}\|^2_{L^1_t(L^2)}+\|\nabla\tilde{\Pi}\|^2_{L^1_t(L^2)}$. Notice that

$$\begin{aligned}
&\int_0^t \|\tilde{u}\cdot\nabla\tilde{u}\|_{L^2}+\|\tilde{u}\cdot\nabla\bar{u}\|_{L^2}+\|\bar{u}\cdot\nabla\tilde{u}\|_{L^2}\,dt'\\
&\lesssim \int_0^t \|\tilde{u}\|_{L^2}^{\frac12}\|\nabla\tilde{u}\|_{L^2}^{\frac12}\|\Delta\tilde{u}\|_{L^2}+\|\tilde{u}\|_{L^2}^{\frac12}\|\nabla\tilde{u}\|_{L^2}^{\frac12}\|\Delta\bar{u}\|_{L^2}\\
&\qquad\qquad +\|\bar{u}\|_{L^2}^{\frac12}\|\nabla\bar{u}\|_{L^2}^{\frac12}\|\Delta\tilde{u}\|_{L^2}\,dt'\\
&\lesssim \Big(\delta_0+\xi(t)\Big)\ln\langle t\rangle,
\end{aligned}$$

and

$$\begin{aligned}
&\int_0^t \|\tilde{B}\cdot\nabla\tilde{B}\|_{L^2}+\|\tilde{B}\cdot\nabla\bar{B}\|_{L^2}+\|\bar{B}\cdot\nabla\tilde{B}\|_{L^2}\,dt'\\
&\lesssim \int_0^t \|\tilde{B}\|_{L^2}^{\frac12}\|\nabla\tilde{B}\|_{L^2}^{\frac12}\|\Delta\tilde{B}\|_{L^2}+\|\tilde{B}\|_{L^2}^{\frac12}\|\nabla\tilde{B}\|_{L^2}^{\frac12}\|\Delta\bar{u}\|_{L^2}\\
&\qquad\qquad +\|\bar{B}\|_{L^2}^{\frac12}\|\nabla\bar{B}\|_{L^2}^{\frac12}\|\Delta\tilde{B}\|_{L^2}\,dt'\\
&\lesssim \Big(\delta_0+\xi(t)\Big)\ln\langle t\rangle.
\end{aligned}$$

Substituting the above inequality and (6.26) into (6.13) results in

$$\left(\int_0^t \|\Delta\tilde{u}\|_{L^2}+\|\nabla\tilde{\Pi}\|_{L^2}\,dt'\right)^2 \le C\left(\delta_0+\xi(t)\right)^2\ln^2\langle t\rangle. \tag{6.27}$$

Applying (6.21) gives

$$\int_0^t \|\tilde{u}\|_{L^2}^2+\|\tilde{B}\|_{L^2}^2\,dt' \lesssim \Big(\delta_0+\xi(t)\Big)^2\langle t\rangle^{\frac12}. \tag{6.28}$$

Plugging (6.27) and (6.28) into (6.22), we obtain

$$\begin{aligned}
&\frac{d}{dt}\left[e^{-U(t)}\left(\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\right)\right]+2g^2(t)e^{-U(t)}\left[\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla\tilde{B}\|_{L^2}^2\right]\\
&\qquad\qquad +e^{-U(t)}\left[\|\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\right]\\
&\lesssim g^4(t)\delta_0^2\langle t\rangle^{-2\beta(p)}+\Big(\delta_0+\xi(t)\Big)^2\langle t\rangle^{-\frac72}\ln^2\langle t\rangle+\xi^2(t)\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\\
&\quad+\xi^2(t)\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\\
&\lesssim \Big(\delta_0+\xi(t)\Big)^2\langle t\rangle^{-2-2\beta(p)}+\xi^2(t)\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\xi^2(t)\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2.
\end{aligned}$$

Taking $g^2(t)=\frac{\alpha}{c_0(1+t)}$ with $1+2\beta(p)<\alpha<2+6\beta(p)$ in the above inequality and integrating the resulting inequality over $[0,t]$, we arrive at

$$\|\nabla\tilde{u}(t)\|_{L^2}+\|\nabla\tilde{B}(t)\|_{L^2}\lesssim\Big(\delta_0+\xi(t)\Big)\langle t\rangle^{-\frac12-\beta(p)} \tag{6.29}$$

for $t<\bar{T}$. Now, taking $g^2(t)=\frac{\alpha}{c_0(1+t)}$ with $\alpha<1+2\beta(p)$, we deduce from Theorem 1.2 that

$$\int_0^t\langle t'\rangle^{(1+2\beta(p))^-}\left(\|\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\right)dt'\lesssim(\delta_0+\xi(t))^2\quad\text{for } t\le\bar{T}. \tag{6.30}$$

Plugging (6.27) and (6.28) into (6.2), we arrive at

$$\begin{aligned}&\frac{d}{dt}\left[e^{-V(t)}\left(\|\sqrt{\rho}\tilde{u}\|_{L^2}^2+\|\tilde{B}\|_{L^2}^2\right)\right]+c_0g^2(t)e^{-V(t)}\left(\|\sqrt{\rho}\tilde{u}\|_{L^2}^2+\|\tilde{B}\|_{L^2}^2\right)\\&\lesssim\Big(\delta_0^2+\xi(t)\Big)^2\langle t\rangle^{-1-2\beta(p)}+\xi^2(t)\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2+\xi^2(t)\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\end{aligned}$$

Then, taking $g^2(t)=\frac{\alpha}{c_0(1+t)}$ with $1+2\beta(p)<\alpha<2+6\beta(p)$ in the above inequality and integrating the resulting inequality over $[0,t]$, we reach

$$\|\tilde{u}(t)\|_{L^2}+\|\tilde{B}(t)\|_{L^2}\lesssim\Big(\delta_0+\xi(t)\Big)\langle t\rangle^{-\beta(p)}. \tag{6.31}$$

Step 3 : Time integral estimates of $\|\nabla\bar{\Pi}\|_{L^2}$, $\|\Delta\tilde{u}(t)\|_{L^2}$ and $\|\Delta\tilde{B}(t)\|_{L^2}$. It follows from (6.11) and (6.13) that

$$\begin{aligned}\|\Delta\tilde{\Pi}\|_{L^2}+\|\Delta\tilde{u}\|_{L^2}\lesssim&\|\partial_t\tilde{u}\|_{L^2}+\|\tilde{u}\|_{L^2}^{\frac12}\|\nabla\tilde{u}\|_{L^2}^{\frac12}\|\Delta\tilde{u}\|_{L^2}+\|\tilde{B}\|_{L^2}^{\frac12}\|\nabla\tilde{B}\|_{L^2}^{\frac12}\|\Delta\tilde{B}\|_{L^2}\\&+\|\tilde{\rho}\|_{L^\infty}\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}+\|\tilde{\rho}\|_{L^\infty}\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}\\&+\|\nabla\tilde{u}\|_{L^2}\left(\|\nabla\bar{u}\|_{L^2}^{\frac12}\|\Delta\bar{u}\|_{L^2}^{\frac12}+\|\bar{u}\|_{L^\infty}\right)\\&+\|\nabla\tilde{B}\|_{L^2}\left(\|\nabla\bar{B}\|_{L^2}^{\frac12}\|\Delta\bar{B}\|_{L^2}^{\frac12}+\|\bar{B}\|_{L^\infty}\right),\end{aligned}$$

and from the equation of $\tilde{B}$, we have

$$\begin{aligned}\|\Delta\tilde{B}\|_{L^2}\lesssim&\|\partial_t\tilde{B}\|_{L^2}+\|\nabla\tilde{B}\|_{L^2}\|\bar{u}\|_{L^\infty}+\|\nabla\tilde{u}\|_{L^2}\|\bar{B}\|_{L^\infty}\\&+\|\tilde{u}\|_{L^2}^{\frac12}\|\nabla\tilde{u}\|_{L^2}^{\frac12}\|\Delta\tilde{B}\|_{L^2}+\|\tilde{B}\|_{L^2}^{\frac12}\|\nabla\tilde{B}\|_{L^2}^{\frac12}\|\Delta\tilde{u}\|_{L^2}\\&+\|\nabla\tilde{u}\|_{L^2}\|\nabla\bar{B}\|_{L^2}^{\frac12}\|\Delta\bar{B}\|_{L^2}^{\frac12}+\|\nabla\tilde{B}\|_{L^2}\|\nabla\bar{u}\|_{L^2}^{\frac12}\|\Delta\bar{u}\|_{L^2}^{\frac12}.\end{aligned}$$

Inequality (6.9) along with

$$\begin{aligned}\|\bar{u}\|_{L^\infty}&\lesssim\|\bar{u}\|_{L^6}^{\frac12}\|\nabla\bar{u}\|_{L^6}^{\frac12}\lesssim\|\nabla\bar{u}\|_{L^2}^{\frac12}\|\Delta\bar{u}\|_{L^2}^{\frac12},\\\|\bar{B}\|_{L^\infty}&\lesssim\|\bar{B}\|_{L^6}^{\frac12}\|\nabla\bar{B}\|_{L^6}^{\frac12}\lesssim\|\nabla\bar{B}\|_{L^2}^{\frac12}\|\Delta\bar{B}\|_{L^2}^{\frac12},\end{aligned}$$

implies for $t\leq\bar{T}$

$$\begin{aligned}&\|\nabla\tilde{\Pi}\|_{L^2}+\|\Delta\tilde{u}\|_{L^2}+\|\Delta\tilde{B}\|_{L^2}\\\lesssim&\|\partial_t\tilde{u}\|_{L^2}+\|\partial_t\tilde{B}\|_{L^2}+\|\tilde{\rho}\|_{L^\infty}\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}+\|\tilde{\rho}\|_{L^\infty}\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}\\&+\left(\|\nabla\tilde{u}\|_{L^2}+\|\nabla\tilde{B}\|_{L^2}\right)\left(\|\nabla\bar{u}\|_{L^2}^{\frac12}\|\Delta\bar{u}\|_{L^2}^{\frac12}+\|\nabla\bar{B}\|_{L^2}^{\frac12}\|\Delta\bar{B}\|_{L^2}^{\frac12}\right)\end{aligned}$$

Therefore, thanks to Theorem (1.2) and (6.30), we obtain

$$\begin{aligned}
&\int_0^t \langle t'\rangle^{(1+2\beta(p))^-}\left(\|\Delta\tilde{u}\|_{L^2}^2+\|\Delta\tilde{B}\|_{L^2}^2+\|\nabla\tilde{\Pi}\|_{L^2}^2\right)dt'\\
\lesssim&\int_0^t \langle t'\rangle^{(1+2\beta(p))^-}\left(\|\partial_t\tilde{u}\|_{L^2}^2+\|\partial_t\tilde{B}\|_{L^2}^2\right)dt'\\
&+(\delta_0+\xi(t))^2\int_0^t\|\nabla\bar{u}\|_{L^2}\|\Delta\bar{u}\|_{L^2}+\|\nabla\bar{B}\|_{L^2}\|\Delta\bar{B}\|_{L^2}\,dt'\\
&+(\delta_0+\xi(t))^2\int_0^t \langle t'\rangle^{(1+2\beta(p))^-}\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\,dt'\\
&+(\delta_0+\xi(t))^2\int_0^t \langle t'\rangle^{(1+2\beta(p))^-}\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2\,dt'\\
\le&C\left(\delta_0+\xi(t)\right)^2
\end{aligned}$$

for $t\le\bar{T}$. So finally, we obtain that

$$\int_0^t \langle t'\rangle^{(1+2\beta(p))^-}\left(\|\Delta\tilde{u}\|_{L^2}^2+\|\Delta\tilde{B}\|_{L^2}^2+\|\nabla\tilde{\Pi}\|_{L^2}^2\right)dt'\le C\left(\delta_0+\xi(t)\right)^2 \tag{6.32}$$

which leads to

$$\int_0^t \|\Delta\tilde{u}\|_{L^2}+\|\Delta\tilde{B}\|_{L^2}+\|\nabla\tilde{\Pi}\|_{L^2}\,dt'\le C\left(\delta_0+\xi(t)\right) \tag{6.33}$$

for $t\le\bar{T}$.

Step 4 : Estimate of $\int_0^t\|\tilde{u}(t')\|\,dt'$ and $\int_0^t\|\tilde{B}(t')\|_{L^\infty}\,dt'$. Thanks to (6.29), we have

$$\begin{aligned}
\|\tilde{u}(t)\|_{L^\infty}&\lesssim\|\tilde{u}(t)\|_{L^6}^{\frac12}\|\nabla\tilde{u}(t)\|_{L^6}^{\frac12}\\
&\lesssim\|\nabla\tilde{u}(t)\|_{L^2}^{\frac12}\|\Delta\tilde{u}(t)\|_{L^2}^{\frac12}\\
&\le C_\eta\left(\delta_0+\xi(t)\right)\langle t\rangle^{-\frac12-\beta(p)}+\eta\|\Delta\tilde{u}(t)\|_{L^2},
\end{aligned} \tag{6.34}$$

and

$$\|\tilde{B}(t)\|_{L^\infty}\le C_\eta\left(\delta_0+\xi(t)\right)\langle t\rangle^{-\frac12-\beta(p)}+\eta\|\Delta\tilde{B}(t)\|_{L^2}, \tag{6.35}$$

for any $\eta>0$. It is easy to observe from the transport equation that

$$\partial_t\tilde{\rho}+u\cdot\nabla\tilde{\rho}=-\tilde{u}\cdot\nabla\bar{\rho}.$$

From this and $\operatorname{div}u=0$, we deduce that

$$\|\tilde{\rho}(t)\|_{L_t^\infty(L^q)}\le\|\tilde{\rho}_0\|_{L^q}+\|\nabla\bar{\rho}\|_{L_t^\infty(L^q)}\|\tilde{u}\|_{L_t^1(L^\infty)}$$

for any $q\in[1,\infty]$. From the above and (5.2), we get

$$\xi(t)\le\delta_0+C\|\tilde{u}\|_{L_t^1(L^\infty)}. \tag{6.36}$$

From (6.32), we know that

$$\begin{aligned}
&\int_0^t\|\Delta\tilde{u}\|_{L^2}+\|\Delta\tilde{B}\|_{L^2}\,dt'\\
\lesssim&\left(\int_0^t\left(\|\Delta\tilde{u}\|_{L^2}^2+\|\Delta\tilde{B}\|_{L^2}^2\right)\langle t'\rangle^{(1+2\beta(p))^-}\,dt'\right)^{\frac12}\\
\le&C\left(\delta_0+\xi(t)\right).
\end{aligned}$$

Plugging (6.36) into (6.34) and integrating the resulting equation over $[0,t]$, we obtain

$$\begin{aligned}
&\|\tilde{u}\|_{L^1_t(L^\infty)} + \|\tilde{B}\|_{L^1_t(L^\infty)} \\
&\leq C_\eta \delta_0 + C_\eta \int_0^t \langle t' \rangle^{-\frac{1}{2}-\beta(p)} \int_0^{t'} \|\tilde{u}\|_{L^\infty} + \|\tilde{B}\|_{L^\infty} \, d\tau \, dt' \\
&\quad + \eta \left(\|\Delta \tilde{u}\|_{L^1_t(L^2)} + \|\Delta \tilde{B}\|_{L^1_t(L^2)} \right) \\
&\leq C_\eta \delta_0 + C_\eta \int_0^t \langle t' \rangle^{-\frac{1}{2}-\beta(p)} \int_0^{t'} \|\tilde{u}\|_{L^\infty} + \|\tilde{B}\|_{L^\infty} \, d\tau \, dt' \\
&\quad + \eta \left(\delta_0 + \|\tilde{u}\|_{L^1_t(L^\infty)} + \|\tilde{B}\|_{L^1_t(L^\infty)} \right).
\end{aligned} \tag{6.37}$$

Taking $\eta = \frac{1}{2C}$. Let $m(t) = \|\tilde{u}\|_{L^1_t(L^\infty)} + \|\tilde{B}\|_{L^1_t(L^\infty)}$, we obtain

$$m(t) \leq C \left(\delta_0 + \int_0^t \langle t' \rangle^{-\frac{1}{2}-\beta(p)} m(t') \, dt' \right)$$

for $t \leq \bar{T}$. Applying the Gronwall's inequality gives

$$m(t) = \|\tilde{u}\|_{L^1_t(L^\infty)} + \|\tilde{B}\|_{L^1_t(L^\infty)} \leq C\delta_0$$

for $t \leq \bar{T}$. From this and (6.36), we deduce that

$$\xi(t) \leq C\delta_0 \quad \text{for } t \leq \bar{T}. \tag{6.38}$$

Thanks to (6.29), (6.30) and (6.32), we arrive at (6.18) and (6.19) for $t \leq \bar{T}$. Moreover, (6.20) and (6.38) ensures that

$$\begin{aligned}
&\|\tilde{u}(t)\|_{L^2} + \|\tilde{B}(t)\|_{L^2} + \|\nabla \tilde{u}(t)\|_{L^2} + \|\nabla \tilde{B}(t)\|_{L^2} \\
&\qquad + \|\sqrt{\rho}\partial_t \tilde{u}\|_{L^2_t(L^2)} + \|\partial_t \tilde{B}\|_{L^2_t(L^2)} + \|\Delta \tilde{u}\|_{L^2_t(L^2)} + \|\Delta \tilde{B}\|_{L^2_t(L^2)} \leq C\delta_0
\end{aligned}$$

for $t \leq \bar{T}$. Now, let ν be the small constant determined in (6.9). We take δ_0 sufficiently small such that

$$\|\tilde{u}(t)\|_{L^2} \|\nabla \tilde{u}(t)\|_{L^2} + \|\tilde{B}(t)\|_{L^2} \|\nabla \tilde{B}(t)\|_{L^2} \leq C\delta_0^2 \leq \frac{\nu}{2}$$

for $t \leq \bar{T}$. This shows that $\bar{T}$ can be any time smaller than $\tilde{T}^*$. This completes the proof of the proposition. □

Proposition 6.4. Under the assumptions of Theorem 1.3, there exist constants C and c so that if

$$A_0 := \|\tilde{u}_0\|_{H^1} + \|\tilde{u}_0\|_{L^p} + \|\tilde{B}_0\|_{H^1} + \|\tilde{B}_0\|_{L^p} + \|\tilde{a}_0\|_{B^{3/2}_{2,1}} \leq c,$$

then we have

$$\begin{aligned}
&\|\tilde{a}\|_{\tilde{L}^\infty_t(B^{3/2}_{2,1})} + \|\tilde{u}\|_{\tilde{L}^\infty_t(L^p)} + \|\tilde{u}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} \\
&\qquad + \|\tilde{B}\|_{\tilde{L}^\infty_t(L^p)} + \|\tilde{B}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} \leq CA_0
\end{aligned} \tag{6.39}$$

for all $t < \tilde{T}^*$.

Proof. Note that $\delta_0 \lesssim A_0$ and $\tilde{a} = a - \bar{a}$. Thanks to Proposition 5.1, Proposition 5.2 and Proposition 6.3, we get by applying (2.7) to the transport equations in (1.2) and using Gronwall's inequality that

$$\|a\|_{\tilde{L}^\infty_t(\dot{H}^2)} \leq C \exp\Big\{C\|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\Big\}\|a_0\|_{\dot{H}^2}. \tag{6.40}$$

Similarly applying (2.7) to the transport equations in (6.1) and using Gronwall's inequality that

$$\|\tilde{a}\|_{\tilde{L}^\infty_t(B^{3/2}_{2,1})} \leq CA_0 \exp\Big\{C\|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\Big\} \tag{6.41}$$

for any $t < \tilde{T}^*$. Applying Lemma 2.6 to the second equation in (6.1) yields

$$\begin{aligned}
&\|\tilde{u}\|_{\tilde{L}^\infty_t(\dot{B}^{\frac{1}{2}}_{2,1})} + \|\tilde{u}\|_{\tilde{L}^1_t(\dot{B}^{5/2}_{2,1})} + \|\tilde{B}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{B}\|_{\tilde{L}^1_t(\dot{B}^{5/2}_{2,1})} \\
&\leq C \exp\Big(C\|\bar{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} + C\|\bar{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\Big)\bigg\{\|\tilde{u}_0\|_{\dot{B}^{\frac{1}{2}}_{2,1}} + \|\tilde{B}_0\|_{\dot{B}^{1/2}_{2,1}} \\
&\quad + \|\tilde{u}\cdot\nabla\bar{u}\|_{\tilde{L}^1_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{u}\cdot\nabla\tilde{u}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} + \|a\,\bar{B}\cdot\nabla\tilde{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \\
&\quad + \|(1+a)(\tilde{B}\cdot\nabla\bar{B} + \tilde{B}\cdot\nabla\tilde{B})\|_{L^1_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{a}(\Delta\bar{u} - \nabla\bar{\Pi})\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \\
&\quad + \|\tilde{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{B}\cdot\nabla\bar{u}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{B}\cdot\nabla\tilde{u}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \\
&\quad + \|\tilde{u}\cdot\nabla\bar{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{u}\cdot\nabla\tilde{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} + \|a\|_{L^\infty_t(\dot{H}^2)}\|\tilde{u}\|_{L^1_t(\dot{B}^2_{2,1})} \\
&\quad + \|a\|_{L^\infty_t(\dot{H}^2)}\|\nabla\tilde{\Pi}\|_{L^1_t(L^2)}\bigg\}.
\end{aligned} \tag{6.42}$$

Next, we list some useful estimates

$$\begin{aligned}
&\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim \|\tilde{u}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})}\|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}, \\
&\|(1+a)\tilde{B}\cdot\nabla\tilde{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim (1+\|a\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})})\|\tilde{B}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})}\|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}, \\
&\|\tilde{B}\cdot\nabla\tilde{u}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim \|\tilde{B}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})}\|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}, \\
&\|\tilde{u}\cdot\nabla\tilde{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim \|\tilde{u}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})}\|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}, \\
&\|\tilde{u}\cdot\nabla\bar{u}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim \|\tilde{u}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})}\|\nabla\bar{u}\|_{L^1_t(\dot{B}^{3/2}_{2,1})}, \\
&\|a\,\bar{B}\cdot\nabla\tilde{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim \|a\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\bar{B}\|_{L^\infty_t(\dot{B}^1_{2,1})}\|\tilde{B}\|_{L^1_t(\dot{B}^2_{2,1})}, \\
&\|(1+a)\tilde{B}\cdot\nabla\bar{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim (1+\|a\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})})\|\tilde{B}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})}\|\bar{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}, \\
&\|\tilde{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim \|\tilde{a}\|_{L^1_t(\dot{B}^{3/2}_{2,1})}\|\bar{B}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})}\|\bar{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}, \\
&\|\tilde{u}\cdot\nabla\bar{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim \|\tilde{u}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})}\|\bar{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}, \\
&\|\tilde{a}\,(\Delta\bar{u} - \nabla\bar{\Pi})\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \lesssim \|\tilde{a}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\Delta\bar{u} - \nabla\bar{\Pi}\|_{L^1_t(\dot{B}^{1/2}_{2,1})} \\
&\qquad\qquad \leq CA_0 \exp\Big\{C\|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\Big\},
\end{aligned}$$

which along with (6.18), (6.40), (6.42) and the following inequalities

$$\begin{aligned}
&\|\tilde{u}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})} \lesssim \|\tilde{u}\|^{\frac{1}{2}}_{L^\infty_t(L^2)}\|\nabla\tilde{u}\|^{\frac{1}{2}}_{L^\infty_t(L^2)},\\
&\|\tilde{u}\|_{L^1_t(\dot{B}^{2}_{2,1})} \lesssim \|\tilde{u}\|^{\frac{2}{3}}_{L^1_t(\dot{B}^{5/2}_{2,1})}\|\tilde{u}\|^{\frac{1}{3}}_{L^1_t(\dot{H}^1)},\\
&\|\tilde{B}\|_{L^\infty_t(\dot{B}^{1/2}_{2,1})} \lesssim \|\tilde{B}\|^{\frac{1}{2}}_{L^\infty_t(L^2)}\|\nabla\tilde{B}\|^{\frac{1}{2}}_{L^\infty_t(L^2)},\\
&\|\tilde{B}\|_{L^1_t(\dot{B}^{2}_{2,1})} \lesssim \|\tilde{B}\|^{\frac{2}{3}}_{L^1_t(\dot{B}^{5/2}_{2,1})}\|\tilde{B}\|^{\frac{1}{3}}_{L^1_t(\dot{H}^1)},
\end{aligned}$$

imply that

$$\begin{aligned}
&\|\tilde{u}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} + \|\tilde{B}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\\
&\le C\bigg[\|\tilde{u}_0\|_{\dot{B}^{1/2}_{2,1}} + \|\tilde{B}_0\|_{\dot{B}^{1/2}_{2,1}} + \delta_0\|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} + \delta_0\|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} + A_0\\
&\quad + A_0\exp\Big(C\|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\Big)\bigg] + \frac{1}{2}\|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} + \frac{1}{2}\|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}.
\end{aligned}\tag{6.43}$$

This gives

$$\begin{aligned}
&\|\tilde{u}\|_{\tilde{L}^\infty_t(\dot{B}^{\frac{1}{2}}_{2,1})} + \|\tilde{B}\|_{\tilde{L}^\infty_t(\dot{B}^{\frac{1}{2}}_{2,1})} + \left(\frac{1}{2} - C\delta_0\right)\Big(\|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} + \|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\Big)\\
&\lesssim \|\tilde{u}_0\|_{\dot{B}^{1/2}_{2,1}} + \|\tilde{B}_0\|_{\dot{B}^{1/2}_{2,1}} + A_0 + A_0\exp\Big(C\|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\Big).
\end{aligned}$$

Taking $A_0 < c$ sufficiently small, we deduce

$$\begin{aligned}
&\|\tilde{a}\|_{\tilde{L}^\infty_t(B^{3/2}_{2,1})} + \|\tilde{u}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\\
&\qquad + \|\tilde{B}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})} + \|\tilde{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})} \le CA_0
\end{aligned}\tag{6.44}$$

for any $t < \tilde{T}^*$.

On the other hand, by multiplying by $|\tilde{u}^i|^{p-1}\mathrm{sgn}(\tilde{u}^i)$ the $\tilde{u}^i$ equation in (6.1) for $i = 1, 2, 3$ and integrating the resulting equation over $\mathbb{R}^3$, we obtain

$$\begin{aligned}
\frac{d}{dt}\|\tilde{u}\|^p_{L^p} \lesssim \|\tilde{u}\|^{p-1}_{L^p}\Big\{&\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^p} + \|\tilde{u}\cdot\nabla\bar{u}\|_{L^p} + \|\bar{u}\cdot\nabla\tilde{u}\|_{L^p}\\
&+ \|(\bar{a}+\tilde{a})(\Delta\tilde{u} - \nabla\tilde{\Pi})\|_{L^p} + \|\nabla\tilde{\Pi}\|_{L^p}\\
&+ \|\tilde{a}\,(\Delta\bar{u} - \nabla\bar{\Pi})\|_{L^p} + \|\tilde{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^p}\\
&+ \|(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\tilde{B}\|_{L^p} + \|(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\bar{B}\|_{L^p}\\
&+ \|(1+\bar{a}+\tilde{a})\bar{B}\cdot\nabla\tilde{B}\|_{L^p}\Big\}.
\end{aligned}\tag{6.45}$$

Similarly, for $\tilde{B}$, we have

$$\begin{aligned}
\frac{d}{dt}\|\tilde{B}\|^p_{L^p} \lesssim \|\tilde{B}\|^{p-1}_{L^p}\Big\{&\|\tilde{u}\cdot\nabla\tilde{B}\|_{L^p} + \|\tilde{B}\cdot\nabla\tilde{u}\|_{L^p} + \|\tilde{B}\cdot\nabla\bar{u}\|_{L^p}\\
&+ \|\bar{B}\cdot\nabla\tilde{u}\|_{L^p} + \|\tilde{u}\cdot\nabla\bar{B}\|_{L^p} + \|\bar{u}\cdot\nabla\tilde{B}\|_{L^p}\Big\}
\end{aligned}\tag{6.46}$$

Notice that applying the operator div to the second equation in (6.1) results in

$$\begin{aligned}\Delta\tilde{\Pi} =&\mathrm{div}\Big\{-\tilde{u}\cdot\nabla\tilde{u}-\tilde{u}\cdot\nabla\bar{u}-\bar{u}\cdot\nabla\tilde{u}+(\bar{a}+\tilde{a})(\Delta\tilde{u}-\nabla\tilde{\Pi})\\&+(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\tilde{B}+(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\bar{B}\\&+(1+\bar{a}+\tilde{a})\bar{B}\cdot\nabla\tilde{B}+\tilde{a}\,(\Delta\bar{u}-\nabla\bar{\Pi})+\tilde{a}\,\bar{B}\cdot\nabla\bar{B}\Big\}\end{aligned}$$

which together with the elliptic estimates implies that

$$\begin{aligned}\|\nabla\tilde{\Pi}\|_{L^p}\lesssim&\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^p}+\|\tilde{u}\dot{\nabla}\bar{u}\|_{L^p}+\|\bar{u}\cdot\nabla\tilde{u}\|_{L^p}\\&+\|(\bar{a}+\tilde{a})(\Delta\tilde{u}-\nabla\tilde{\Pi})\|_{L^p}+\|(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\tilde{B}\|_{L^p}\\&+\|(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\bar{B}\|_{L^p}+\|(1+\bar{a}+\tilde{a})\bar{B}\cdot\nabla\tilde{B}\|_{L^p}\\&+\|\tilde{a}\,(\Delta\bar{u}-\nabla\bar{\Pi})\|_{L^p}+\|\tilde{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^p}.\end{aligned}$$

As a consequence, we deduce from (6.45) that

$$\begin{aligned}\|\tilde{u}\|_{L^\infty_t(L^p)}\lesssim&\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^1_t(L^p)}+\|\tilde{u}\cdot\nabla\bar{u}\|_{L^1_t(L^p)}+\|\bar{u}\cdot\nabla\tilde{u}\|_{L^1_t(L^p)}\\&+\|(\bar{a}+\tilde{a})(\Delta\tilde{u}-\nabla\tilde{\Pi})\|_{L^1_t(L^p)}+\|\tilde{a}\,(\Delta\bar{u}-\nabla\bar{\Pi})\|_{L^1_t(L^p)}\\&+\|\tilde{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^1_t(L^p)}+\|(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\tilde{B}\|_{L^1_t(L^p)}\\&+\|(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\bar{B}\|_{L^1_t(L^p)}+\|(1+\bar{a}+\tilde{a})\bar{B}\cdot\nabla\tilde{B}\|_{L^1_t(L^p)}\\&+\|\tilde{u}_0\|_{L^p}.\end{aligned}\tag{6.47}$$

However, since $p\in(1,\frac{6}{5})$, it follows from Proposition 6.3 and Theorem 1.2 that

$$\begin{aligned}&\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^1_t(L^p)}+\|\tilde{u}\cdot\nabla\bar{u}\|_{L^1_t(L^p)}\\\lesssim&\|\tilde{u}\|_{L^\infty_t(L^{\frac{2p}{2-p}})}\Big(\|\nabla\tilde{u}\|_{L^1_t(L^2)}+\|\nabla\bar{u}\|_{L^1_t(L^2)}\Big)\\\lesssim&\|\tilde{u}\|_{L^\infty_t(H^{1/2})}\lesssim\delta_0\lesssim A_0,\end{aligned}$$

where we used the Sobolev embedding inequality $\|u\|_{L^{\frac{2p}{2-p}}}\lesssim\|u\|_{H^{1/2}}$. Similarly, we have

$$\begin{aligned}&\|\tilde{B}\cdot\nabla\tilde{B}\|_{L^1_t(L^p)}+\|\tilde{B}\cdot\nabla\bar{B}\|_{L^1_t(L^p)}\lesssim A_0,\\&\|\tilde{u}\cdot\nabla\tilde{B}\|_{L^1_t(L^p)}+\|\tilde{u}\cdot\nabla\bar{B}\|_{L^1_t(L^p)}\lesssim A_0,\\&\|\tilde{B}\cdot\nabla\tilde{u}\|_{L^1_t(L^p)}+\|\tilde{B}\cdot\nabla\bar{u}\|_{L^1_t(L^p)}\lesssim A_0,\end{aligned}$$

and

$$\begin{aligned}&\|(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\tilde{B}\|_{L^1_t(L^p)}+\|(1+\bar{a}+\tilde{a})\tilde{B}\cdot\nabla\bar{B}\|_{L^1_t(L^p)}\\\lesssim&\big(1+\|\bar{a}\|_{L^\infty_t(L^\infty)}+\|\tilde{a}\|_{L^\infty_t(L^\infty)}\big)\Big(\|\tilde{B}\cdot\nabla\tilde{B}\|_{L^1_t(L^p)}+\|\tilde{B}\cdot\nabla\bar{B}\|_{L^1_t(L^p)}\Big)\lesssim A_0.\end{aligned}$$

Also, we can obtain

$$\begin{aligned}\|\bar{u}\cdot\nabla\tilde{u}\|_{L^1_t(L^p)}\lesssim&\|\bar{u}\|_{L^\infty_t(L^{\frac{2p}{2-p}})}\|\nabla\tilde{u}\|_{L^1_t(L^2)}\\\lesssim&\|\bar{u}\|_{L^\infty_t(H^{1/2})}\|\nabla\tilde{u}\|_{L^1_t(L^2)}\lesssim A_0.\end{aligned}$$

Similarly, we have

$$\begin{aligned}&\|\bar{B}\cdot\nabla\tilde{B}\|_{L^1_t(L^p)}+\|\bar{B}\cdot\nabla\tilde{u}\|_{L^1_t(L^p)}+\|\bar{u}\cdot\nabla\tilde{B}\|_{L^1_t(L^p)}\lesssim A_0,\\&\|(1+\bar{a}+\tilde{a})\bar{B}\cdot\nabla\tilde{B}\|_{L^1_t(L^p)}\lesssim(1+\|\bar{a}\|_{L^\infty_t(L^\infty)}\\&\qquad\qquad+\|\tilde{a}\|_{L^\infty_t(L^\infty)})\|\bar{B}\|_{L^\infty_t(H^{1/2})}\|\nabla\tilde{B}\|_{L^1_t(L^2)}\lesssim A_0,\end{aligned}$$

and

$$\|(\bar{a}+\tilde{a})(\Delta\tilde{u}-\nabla\tilde{\Pi})\|_{L^1_t(L^p)}+\|\tilde{a}\,(\Delta\bar{u}-\nabla\bar{\Pi})\|_{L^1_t(L^p)}+\|\tilde{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^1_t(L^p)}\lesssim A_0.$$

Plugging the above estimates into (6.46) and (6.47), we arrive at

$$\|\tilde{u}\|_{L^\infty_t(L^p)}+\|\tilde{B}\|_{L^\infty_t(L^p)}\lesssim A_0.$$

This along with (6.44) complete the proof. □

Proposition 6.5. Under the assumption of Theorem 1.3, there holds

$$\|\tilde{a}(t)\|_{L^\infty}\le CA_0 \tag{6.48}$$

and

$$\|\nabla\tilde{a}(t)\|_{L^\infty}\le C\left(\|\nabla\tilde{a}_0\|_{L^\infty}+\|\bar{a}_0\|_{B^{7/2}_{2,1}}\right), \tag{6.49}$$

where A_0 defined as before.

Proof. Firstly, let us give the equation of $\tilde{a}$

$$\partial_t\tilde{a}+u\cdot\nabla\tilde{a}=-\tilde{u}\cdot\nabla\bar{a}.$$

Using basic estimates about transport equations, we have

$$\|\tilde{a}(t)\|_{L^\infty}\le Ce^{C\int_0^t\|\nabla u\|_{L^\infty}dt'}\left(\|\tilde{a}_0\|_{L^\infty}+\int_0^t\|\tilde{u}\cdot\nabla\bar{a}\|_{L^\infty}\,dt'\right)$$

Moreover, we obtain

$$\|\tilde{a}(t)\|_{L^\infty}\lesssim\|\tilde{a}_0\|_{L^\infty}+\|\nabla\bar{a}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}A_0.$$

By the definition of A_0 and Proposition 5.2, we get

$$\|\tilde{a}(t)\|_{L^\infty}\le CA_0.$$

Similarly, we give the equation of $\nabla\tilde{a}$

$$\partial_t\nabla\tilde{a}+u\cdot\nabla(\nabla\tilde{a})=-\tilde{u}\cdot\nabla(\nabla\bar{a}).$$

Then classical estimates about transport equations, Proposition 4.5 and Proposition 6.3 gives

$$\|\nabla\tilde{a}(t)\|_{L^\infty}\lesssim\|\nabla\tilde{a}_0\|_{L^\infty}+\|\bar{a}\|_{L^\infty_t(\dot{B}^{7/2}_{2,1})}.$$

Due to Lemma 2.3, we obtain

$$\|\bar{a}(t)\|_{B^{7/2}_{2,1}}\lesssim\|\bar{a}_0\|_{B^{7/2}_{2,1}},$$

where we used (5.8). From this, we easily obtain

$$\|\nabla\tilde{a}(t)\|_{L^\infty}\le C\left(\|\nabla\tilde{a}_0\|_{L^\infty}+\|\bar{a}_0\|_{B^{7/2}_{2,1}}\right).$$

□

Proposition 6.6. Under the assumptions of Theorem 1.3, there holds

$$\sup_{t\geq t_0}\left(\|\nabla^2\tilde{u}(t)\|_{L^2}^2+\|\nabla^2\tilde{B}(t)\|_{L^2}^2\right)+\int_{t_0}^{\infty}\|\partial_t\nabla\tilde{u}(t)\|_{L^2}^2+\|\partial_t\nabla\tilde{B}(t)\|_{L^2}^2\,dt \\ +\int_{t_0}^{\infty}\|\nabla^3\tilde{u}(t)\|_{L^2}^2+\|\nabla^3\tilde{B}(t)\|_{L^2}^2\,dt\leq C,$$

where C depends on the initial data.

Proof. Taking derivative to the second and third equation of (6.3), we obtain

$$\begin{aligned}\rho\partial_t\partial_j\tilde{u}^i-\Delta\partial_j\tilde{u}^i+\partial_j\partial_i\tilde{\Pi}&=-\partial_j\rho\,\partial_t\tilde{u}^i-\partial_j\rho\,u\cdot\nabla\tilde{u}^i-\rho\,\partial_j u\cdot\nabla\tilde{u}^i\\&\quad-\rho\,u\cdot\nabla\partial_j\tilde{u}^i+\partial_j B\cdot\nabla\tilde{B}^i+B\cdot\nabla\partial_j\tilde{B}^i-\partial_j\rho\,\tilde{u}\cdot\nabla\bar{u}^i-\rho\,\partial_j\tilde{u}\cdot\nabla\bar{u}^i\\&\quad-\rho\,\tilde{u}\cdot\nabla\partial_j\bar{u}^i+\partial_j\tilde{B}\cdot\nabla\bar{B}^i+\tilde{B}\cdot\nabla\partial_j\bar{B}^i-\frac{\tilde{\rho}}{\bar{\rho}}(\Delta\partial_j\bar{u}^i-\partial_j\partial_i\tilde{\Pi})\\&\quad-\partial_j\left(\frac{\tilde{\rho}}{\bar{\rho}}\right)(\Delta\bar{u}^i-\partial_i\tilde{\Pi})-\frac{\tilde{\rho}}{\bar{\rho}}\,\partial_j\bar{B}\cdot\nabla\bar{B}^i-\frac{\tilde{\rho}}{\bar{\rho}}\,\bar{B}\cdot\nabla\partial_j\bar{B}^i\\&\quad-\partial_j\left(\frac{\tilde{\rho}}{\bar{\rho}}\right)\bar{B}\cdot\nabla\bar{B}^i\end{aligned}\tag{6.50}$$

and

$$\begin{aligned}\partial_t\partial_j\tilde{B}^i-\Delta\partial_i\tilde{B}^i&=-\partial_j u\cdot\nabla\tilde{B}^i-u\cdot\nabla\partial_j\tilde{B}^i+\partial_j B\cdot\nabla\tilde{u}^i+B\cdot\nabla\partial_j\tilde{u}^i\\&\quad-\partial_j\tilde{B}\cdot\nabla\bar{u}^i-\tilde{B}\cdot\nabla\partial_j\bar{u}^i+\partial_j\tilde{u}\cdot\nabla\bar{B}^i+\tilde{u}\cdot\nabla\partial_j\bar{B}^i.\end{aligned}\tag{6.51}$$

Multiplying (6.50) with $\partial_t\partial_j\tilde{u}^i$ and integrating over $\mathbb{R}^3$, we obtain

$$\begin{aligned}\|\sqrt{\rho}\partial_t\nabla\tilde{u}(t)\|_{L^2}^2+\frac{1}{2}\frac{d}{dt}\|\nabla^2\tilde{u}(t)\|_{L^2}^2&=-\int_{\mathbb{R}^3}\nabla\rho\,\partial_t\tilde{u}\,\partial_t\nabla\tilde{u}\,dx\\&\quad-\int_{\mathbb{R}^3}\nabla\rho\,\tilde{u}\cdot\nabla\tilde{u}\,\partial_t\nabla\tilde{u}\,dx-\int_{\mathbb{R}^3}\nabla\rho\,\bar{u}\cdot\nabla\tilde{u}\,\partial_t\nabla\tilde{u}\,dx\\&\quad-\int_{\mathbb{R}^3}\rho\,\nabla\tilde{u}\cdot\nabla\tilde{u}\,\partial_t\nabla\tilde{u}\,dx-\int_{\mathbb{R}^3}\rho\,\nabla\bar{u}\cdot\nabla\tilde{u}\,\partial_t\nabla\tilde{u}\,dx\\&\quad-\int\mathbb{R}^3\rho\,u\cdot\nabla\nabla\tilde{u}\,\partial_t\nabla\tilde{u}\,dx+\int_{\mathbb{R}^3}\nabla\tilde{B}\cdot\nabla\tilde{B}\,\partial_t\nabla\tilde{u}\,dx\\&\quad+\int_{\mathbb{R}^3}\nabla\bar{B}\cdot\nabla\tilde{B}\,\partial_t\nabla\tilde{u}\,dx+\int_{\mathbb{R}^3}\tilde{B}\cdot\nabla\nabla\tilde{B}\,\partial_t\nabla\tilde{u}\,dx\\&\quad+\int_{\mathbb{R}^3}\bar{B}\cdot\nabla\nabla\tilde{B}\,\partial_t\nabla\tilde{u}\,dx-\int_{\mathbb{R}^3}\nabla\rho\,\tilde{u}\cdot\nabla\bar{u}\,\partial_t\nabla\tilde{u}\,dx\\&\quad-\int_{\mathbb{R}^3}\rho\,\nabla\tilde{u}\cdot\nabla\bar{u}\,\partial_t\nabla\tilde{u}\,dx-\int_{\mathbb{R}^3}\rho\,\tilde{u}\cdot\nabla\nabla\bar{u}\,dx\\&\quad+\int_{\mathbb{R}^3}\nabla\tilde{B}\cdot\nabla\bar{B}\,\partial_t\nabla\tilde{u}\,dx+\int_{\mathbb{R}^3}\tilde{B}\cdot\nabla\nabla\bar{B}\,\partial_t\nabla\tilde{u}\,dx\\&\quad-\int_{\mathbb{R}^3}\frac{\tilde{\rho}}{\bar{\rho}}(\Delta\partial_j\bar{u}^i-\partial_j\partial_i\tilde{\Pi})\,\partial_t\nabla\tilde{u}\,dx-\int_{\mathbb{R}^3}\partial_j\left(\frac{\tilde{\rho}}{\bar{\rho}}\right)(\Delta\bar{u}^i-\partial_i\tilde{\Pi})\,\partial_t\nabla\tilde{u}\,dx\\&\quad-\int_{\mathbb{R}^3}\frac{\tilde{\rho}}{\bar{\rho}}\,\nabla\bar{B}\cdot\nabla\bar{B}\,dx-\int_{\mathbb{R}^3}\frac{\tilde{\rho}}{\bar{\rho}}\,\bar{B}\cdot\nabla\nabla\bar{B}\,\partial_t\nabla\tilde{u}\,dx\\&\quad-\int_{\mathbb{R}^3}\partial_j\left(\frac{\tilde{\rho}}{\bar{\rho}}\right)\bar{B}\cdot\nabla\bar{B}\,dx.\end{aligned}\tag{6.52}$$

For $\tilde{B}$, using same process, we have

$$
\begin{aligned}
\|\partial_t\nabla\tilde{B}(t)\|_{L^2}^2+\frac{1}{2}\frac{d}{dt}\|\nabla^2\tilde{B}(t)\|_{L^2}^2=&-\int_{\mathbb{R}^3}\nabla u\cdot\nabla\tilde{B}\,\partial_t\tilde{B}\,dx\\
&-\int_{\mathbb{R}^3}u\cdot\nabla\nabla\tilde{B}\,\partial_t\nabla\tilde{B}\,dx+\int_{\mathbb{R}^3}\nabla B\cdot\nabla\tilde{u}\,\partial_t\nabla\tilde{B}\,dx\\
&+\int_{\mathbb{R}^3}B\cdot\nabla\nabla\tilde{u}\,\partial_t\nabla\tilde{B}\,dx-\int_{\mathbb{R}^3}\nabla\tilde{B}\cdot\nabla\bar{u}\,\partial_t\nabla\tilde{B}\,dx\\
&-\int_{\mathbb{R}^3}\tilde{B}\cdot\nabla\nabla\bar{u}\,\partial_t\nabla\tilde{B}\,dx+\int_{\mathbb{R}^3}\nabla\tilde{u}\cdot\nabla\bar{B}\,\partial_t\nabla\tilde{B}\,dx\\
&+\int_{\mathbb{R}^3}\tilde{u}\cdot\nabla\nabla\bar{B}\,\partial_t\nabla\tilde{B}\,dx.
\end{aligned}\tag{6.53}
$$

Combining (6.52) and (6.53), we obtain

$$
\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\left(\|\nabla^2\tilde{u}(t)\|_{L^2}^2+\|\nabla^2\tilde{B}(t)\|_{L^2}^2\right)+\|\sqrt{\rho}\,\partial_t\nabla\tilde{u}\|_{L^2}^2+\|\partial_t\nabla\tilde{B}\|_{L^2}^2\\
\le&\|\partial_t\tilde{u}\|_{L^2}^2+\left(\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^2}+\|\bar{u}\cdot\nabla\tilde{u}\|_{L^2}+\|\tilde{u}\cdot\nabla\bar{u}\|_{L^2}\right)^2\\
&+\|\nabla\tilde{u}\|_{L^2}^2\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}+\|\nabla\bar{u}\|_{L^2}^2\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}\\
&+\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}^2+\|\nabla\bar{u}\|_{L^2}\|\nabla^2\bar{u}\|_{L^2}\|\nabla^2\tilde{u}\|_{L^2}^2\\
&+\|\nabla\tilde{B}\|_{L^2}^2\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}+\|\nabla\bar{B}\|_{L^2}^2\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}\\
&+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}^2+\|\nabla\bar{B}\|_{L^2}\|\nabla^2\bar{B}\|_{L^2}\|\nabla^2\tilde{B}\|_{L^2}^2\\
&+\|\nabla\bar{u}\|_{L^2}^2\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}+\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\nabla^3\bar{u}\|_{L^2}^2\\
&\|\nabla\bar{B}\|_{L^2}^2\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\nabla^3\bar{B}\|_{L^2}^2\\
&+\|\tilde{\rho}\|_{L^\infty}^2\|\nabla(\Delta\bar{u}-\nabla\bar{\Pi})\|_{L^2}^2+\|\nabla\tilde{a}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\\
&+\|\tilde{\rho}\|_{L^\infty}^2\|\nabla\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\nabla\bar{B}\|_{L^2}^2\\
&+\|\nabla\tilde{a}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2+\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}^2\\
&+\|\nabla\bar{u}\|_{L^2}^2\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}+\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}^2\\
&+\|\nabla\bar{u}\|_{L^2}\|\nabla^2\bar{u}\|_{L^2}\|\nabla^2\tilde{B}\|_{L^2}^2+\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}^2\\
&+\|\nabla\bar{B}\|_{L^2}^2\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}^2\\
&+\|\nabla\bar{B}\|_{L^2}\|\nabla^2\bar{B}\|_{L^2}\|\nabla^2\tilde{u}\|_{L^2}^2+\|\nabla\bar{u}\|_{L^2}^2\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}\\
&+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\nabla^3\bar{u}\|_{L^2}^2+\|\nabla\bar{B}\|_{L^2}^2\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}\\
&+\|\tilde{u}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\nabla^3\bar{B}\|_{L^2}^2.
\end{aligned}
$$

Multiplying $\frac{1}{\rho}\,\Delta\tilde{u}$ and $\Delta\tilde{u}$ to (6.50) and (6.51) separately and doing some basic energy estimates, we obtain

$$
\begin{aligned}
&\frac{1}{2}\frac{d}{dt}\left(\|\nabla^2\tilde{u}\|_{L^2}^2+\|\nabla^2\tilde{B}\|_{L^2}^2\right)+\|\rho\|_{L^\infty}^{-1}\|\nabla^3\tilde{u}\|_{L^2}^2+\|\nabla^3\tilde{B}\|_{L^2}^2\\
\le&\|\partial_t\nabla\tilde{u}\|_{L^2}^2+\left(\|\tilde{u}\cdot\nabla\tilde{u}\|_{L^2}+\|\bar{u}\cdot\nabla\tilde{u}\|_{L^2}+\|\tilde{u}\cdot\nabla\bar{u}\|_{L^2}\right)^2\\
&+\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}^2+\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}\|\nabla\bar{u}\|_{L^2}^2\\
&+\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}^2+\|\nabla\bar{u}\|_{L^2}\|\nabla^2\bar{u}\|_{L^2}\|\nabla^2\tilde{u}\|_{L^2}^2\\
&+\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}^2+\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}\|\nabla\bar{B}\|_{L^2}^2
\end{aligned}
$$

$$
\begin{aligned}
&+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}^2+\|\nabla\bar{B}\|_{L^2}\|\nabla^2\bar{B}\|_{L^2}\|\nabla^2\tilde{B}\|_{L^2}^2\\
&+\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^3}\|\nabla\bar{u}\|_{L^2}^2+\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\nabla^3\bar{u}\|_{L^2}^2\\
&+\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}\|\nabla\bar{B}\|_{L^2}^2+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\nabla^3\bar{B}\|_{L^2}^2\\
&+\|\tilde{\rho}\|_{L^\infty}^2\|\nabla(\Delta\bar{u}-\nabla\bar{\Pi})\|_{L^2}^2+\|\nabla\tilde{a}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\\
&+\|\tilde{\rho}\|_{L^\infty}^2\|\nabla\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\nabla\bar{B}\|_{L^2}^2\\
&+\|\nabla\tilde{a}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2+\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}^2\\
&+\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}\|\nabla\bar{u}\|_{L^2}^2+\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}^2\\
&+\|\nabla\bar{u}\|_{L^2}\|\nabla^2\bar{u}\|_{L^2}\|\nabla^2\tilde{B}\|_{L^2}^2+\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}^2\\
&+\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}\|\nabla\bar{B}\|_{L^2}^2+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}^2\\
&+\|\nabla\bar{B}\|_{L^2}\|\nabla^2\bar{B}\|_{L^2}\|\nabla^2\tilde{B}\|_{L^2}^2+\|\nabla^2\tilde{B}\|_{L^2}\|\nabla^3\tilde{B}\|_{L^2}\|\nabla\bar{u}\|_{L^2}^2\\
&+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\|\nabla^3\bar{u}\|_{L^2}^2+\|\nabla^2\tilde{u}\|_{L^2}\|\nabla^3\tilde{u}\|_{L^2}\|\nabla\bar{B}\|_{L^2}^2\\
&+\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\nabla^3\bar{B}\|_{L^2}^2.
\end{aligned}
$$

At this point, we got two completed inequalities. The second inequality times a small number then plus the first inequality yieds

$$
\begin{aligned}
&\frac{d}{dt}\left(\|\nabla^2\tilde{u}(t)\|_{L^2}^2+\|\nabla^2\tilde{B}(t)\|_{L^2}^2\right)+\|\sqrt{\rho}\partial_t\nabla\tilde{u}\|_{L^2}^2+\|\partial_t\nabla\tilde{B}\|_{L^2}^2\\
&+\left(\frac{c_0}{2}-C\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}-C\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\right)\left(\|\nabla^3\tilde{u}\|_{L^2}^2+\|\nabla^3\tilde{B}\|_{L^2}^2\right)\\
\le\ &\left(\|\nabla\bar{u}\|_{L^2}\|\nabla^2\bar{u}\|_{L^2}+\|\nabla\bar{B}\|_{L^2}\|\nabla^2\bar{B}\|_{L^2}\right)\left(\|\nabla^2\tilde{u}\|_{L^2}^2+\|\nabla^2\tilde{B}\|_{L^2}^2\right)\\
&+\left(\|\nabla\tilde{u}\|_{L^2}^4+\|\nabla\tilde{B}\|_{L^2}^4+\|\nabla\bar{u}\|_{L^2}^4+\|\nabla\bar{B}\|_{L^2}^4\right)\left(\|\nabla^2\tilde{u}\|_{L^2}^2+\|\nabla^2\tilde{B}\|_{L^2}^2\right.\\
&\left.+\|\nabla^2\bar{u}\|_{L^2}^2+\|\nabla^2\bar{B}\|_{L^2}^2\right)+\|\partial_t\tilde{u}\|_{L^2}^2+\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\nabla^2\tilde{u}\|_{L^2}^2\\
&+\|\nabla\tilde{u}\|_{L^2}^2\|\nabla\bar{u}\|_{L^2}\|\nabla^2\bar{u}\|_{L^2}+\left(\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\right)\left(\|\nabla^3\bar{u}\|_{L^2}^2\right.\\
&\left.+\|\nabla^3\bar{B}\|_{L^2}^2\right)+\|\tilde{\rho}\|_{L^\infty}^2\|\nabla(\Delta\bar{u}-\nabla\bar{\Pi})\|_{L^2}^2+\|\nabla\tilde{a}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\\
&+\|\tilde{\rho}\|_{L^\infty}^2\|\nabla\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\nabla\bar{B}\|_{L^2}^2+\|\nabla\tilde{a}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2.
\end{aligned}
$$

Taking c in Theorem 1.3 small enough, we have

$$
\begin{aligned}
&\frac{d}{dt}\left(\|\nabla^2\tilde{u}(t)\|_{L^2}^2+\|\nabla^2\tilde{B}(t)\|_{L^2}^2\right)+\|\sqrt{\rho}\partial_t\nabla\tilde{u}\|_{L^2}^2+\|\partial_t\nabla\tilde{B}\|_{L^2}^2\\
&\quad+\|\nabla^3\tilde{u}\|_{L^2}^2+\|\nabla^3\tilde{B}\|_{L^2}^2\\
\le\ &\left(\|\nabla\bar{u}\|_{L^2}\|\nabla^2\bar{u}\|_{L^2}+\|\nabla\bar{B}\|_{L^2}\|\nabla^2\bar{B}\|_{L^2}\right)\left(\|\nabla^2\tilde{u}\|_{L^2}^2+\|\nabla^2\tilde{B}\|_{L^2}^2\right)\\
&+\left(\|\nabla\tilde{u}\|_{L^2}^4+\|\nabla\tilde{B}\|_{L^2}^4+\|\nabla\bar{u}\|_{L^2}^4+\|\nabla\bar{B}\|_{L^2}^4\right)\left(\|\nabla^2\tilde{u}\|_{L^2}^2+\|\nabla^2\tilde{B}\|_{L^2}^2\right.\\
&\left.+\|\nabla^2\bar{u}\|_{L^2}^2+\|\nabla^2\bar{B}\|_{L^2}^2\right)+\|\partial_t\tilde{u}\|_{L^2}^2+\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}\|\nabla^2\tilde{u}\|_{L^2}^2\\
&+\|\nabla\tilde{u}\|_{L^2}^2\|\nabla\bar{u}\|_{L^2}\|\nabla^2\bar{u}\|_{L^2}+\left(\|\tilde{u}\|_{L^2}\|\nabla\tilde{u}\|_{L^2}+\|\tilde{B}\|_{L^2}\|\nabla\tilde{B}\|_{L^2}\right)\left(\|\nabla^3\bar{u}\|_{L^2}^2\right.\\
&\left.+\|\nabla^3\bar{B}\|_{L^2}^2\right)+\|\tilde{\rho}\|_{L^\infty}^2\|\nabla(\Delta\bar{u}-\nabla\bar{\Pi})\|_{L^2}^2+\|\nabla\tilde{a}\|_{L^\infty}^2\|\Delta\bar{u}-\nabla\bar{\Pi}\|_{L^2}^2\\
&+\|\tilde{\rho}\|_{L^\infty}^2\|\nabla\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2+\|\tilde{\rho}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\nabla\bar{B}\|_{L^2}^2+\|\nabla\tilde{a}\|_{L^\infty}^2\|\bar{B}\cdot\nabla\bar{B}\|_{L^2}^2.
\end{aligned}
$$

Integrating the above inequality, using decay estimates about reference solution and perturbed solution, we obtain

$$\sup_{t\geq t_0}\Big(\|\nabla^2\tilde{u}(t)\|_{L^2}^2+\|\nabla^2\tilde{B}(t)\|_{L^2}^2\Big)+\int_{t_0}^{\infty}\|\sqrt{\rho}\,\partial_t\nabla\tilde{u}\|_{L^2}^2+\|\partial_t\nabla\tilde{B}\|_{L^2}^2\,dt$$
$$+\int_{t_0}^{\infty}\|\nabla^3\tilde{u}\|_{L^2}^2+\|\nabla^3\tilde{B}\|_{L^2}^2\,dt\leq C.$$

Hence, the proof of Proposition 6.6 is completed. □

Now, we can complete the proof of Theorem 1.3 as following.

Proof. According to the statement at the beginning of this section, given initial data $(\bar{a}_0+\tilde{a}_0,\bar{u}_0+\tilde{u}_0,\bar{B}_0+\tilde{B}_0)$, (1.2) has a unique solution (a,u,B) on $[0,\tilde{T}^*)$ such that

$$\begin{aligned}&a\in C([0,\tilde{T}^*);B^{7/2}_{2,1}(\mathbb{R}^3)),\\&u\in C([0,\tilde{T}^*);B^{2}_{2,1})\cap L^1_{\rm loc}((0,\tilde{T}^*);\dot{B}^{4}_{2,1}(\mathbb{R}^3)),\\&B\in C([0,\tilde{T}^*);B^{2}_{2,1})\cap L^1_{\rm loc}((0,\tilde{T}^*);\dot{B}^{4}_{2,1}(\mathbb{R}^3)).\end{aligned}$$

We need only prove the maximal existence time $\tilde{T}^*=\infty$. Indeed, according to all the decay estimates for reference solution and perturbed solution, we repeat the argument used in the proof of Proposition 5.1 and Proposition 5.2 to prove that $\tilde{T}^*=\infty$. Then a standard interpolation argument gives (1.10) and (1.11). This completes the proof of Theorem 1.3. □

7. Appendix

For completeness, in this section, we give the proof of some Lemmas and Propositions. Firstly, we give the proof of Remark 2.9.

Proof. Using the same method as in the proof of Lemma 2.6, we can get the result. The only difference is that we allow the parameter $s=1$, so we only give the estimates related to $s<1$ which is required in the proof of Lemma 2.6. Notice that

$$\begin{aligned}{\rm div}[\Delta_q,a]\nabla u=&\Delta_q{\rm div}\,\mathcal{R}(a,\nabla u)+\Delta_q{\rm div}\,T_{\nabla u}a\\&-{\rm div}R(a,\Delta_q\nabla u)-{\rm div}[T_a,\Delta_q]\nabla u.\end{aligned}$$

Applying Lemma 2.1, we have

$$\begin{aligned}&\|\Delta_q{\rm div}\,\mathcal{R}(a,\nabla u)(t)\|_{L^2}\\\lesssim&\,2^{\frac{5}{2}q}\sum_{k\geq q-5}\|\Delta_k a(t)\|_{L^2}\|\tilde{\Delta}_k\nabla u(t)\|_{L^2}\\\lesssim&\,2^{\frac{5}{2}q}\sum_{k\geq q-5}2^{-2k}2^{-k-\frac{1}{2}k}2^{2k}\|\Delta_k a(t)\|_{L^2}2^{k+\frac{3}{2}k}\|\tilde{\Delta}_k u(t)\|_{L^2}\\\lesssim&\,2^{\frac{5}{2}q}\sum_{k\geq q-5}d_k(t)2^{-k(1+\frac{5}{2})}\|a(t)\|_{\dot{B}^2_{2,1}}\|u(t)\|_{\dot{B}^{5/2}_{2,1}}\\\lesssim&\,d_q(t)2^{-q}\|a(t)\|_{\dot{B}^2_{2,1}}\|u(t)\|_{\dot{B}^{5/2}_{2,1}},\end{aligned}$$

and

$$\begin{aligned}
&\|\mathrm{div}\, R(a,\Delta_q\nabla u)(t)\|_{L^2}\\
\lesssim&\sum_{|k-q|\le 1}\|\nabla\Delta_k a(t)\|_{L^2}\|\tilde{\Delta}_k\Delta_q\nabla u(t)\|_{L^\infty}\\
&\qquad\qquad+\sum_{|k-q|\le 1}\|\Delta_k a(t)\|_{L^2}\|\tilde{\Delta}_k\Delta_q\Delta u(t)\|_{L^\infty}\\
\lesssim&\sum_{|k-q|\le 1}2^k\|\Delta_k a(t)\|_{L^2}2^{q\frac{5}{2}}\|\tilde{\Delta}_k u(t)\|_{L^2}\\
\lesssim& d_q(t)2^{-q}\|a(t)\|_{\dot{B}^2_{2,1}}\|u(t)\|_{\dot{B}^{5/2}_{2,1}}.
\end{aligned}$$

Note that

$$\begin{aligned}
\|S_{k-1}\nabla u(t)\|_{L^\infty}&\lesssim\sum_{j\le k-2}\|\Delta_j\nabla u(t)\|_{L^\infty}\\
&\lesssim\sum_{j\le k-2}2^{\frac{5}{2}j}\|\Delta_j u(t)\|_{L^2}\\
&\lesssim\|u(t)\|_{\dot{B}^{5/2}_{2,1}},
\end{aligned}$$

this along with Lemma 2.1 leads to

$$\begin{aligned}
\|\Delta_q\mathrm{div}\, T_{\nabla u}a(t)\|_{L^2}&\lesssim 2^q\sum_{|k-q|\le 4}\|\Delta_k a(t)\|_{L^2}\|S_{k-1}\nabla u(t)\|_{L^\infty}\\
&\lesssim 2^q\sum_{|q-k|\le 4}d_q(t)2^{-2q}\|u(t)\|_{\dot{B}^{5/2}_{2,1}}\|a(t)\|_{\dot{B}^2_{2,1}}\\
&\lesssim d_q(t)2^{-q}\|u(t)\|_{\dot{B}^{5/2}_{2,1}}\|a(t)\|_{\dot{B}^2_{2,1}}.
\end{aligned}$$

Notice that

$$\begin{aligned}
\|\nabla S_{k-1}a(t)\|_{L^\infty}&\lesssim\sum_{j\le k-2}\|\nabla\Delta_j a(t)\|_{L^\infty}\\
&\lesssim\sum_{j\le k-2}2^{\frac{1}{2}j}2^{2j}\|\Delta_j a(t)\|_{L^2}\lesssim 2^{\frac{k}{2}}\|a(t)\|_{\dot{B}^2_{2,1}},
\end{aligned}$$

this gives

$$\begin{aligned}
\|\mathrm{div}\,[T_a,\Delta_q]\nabla u(t)\|_{L^2}&\lesssim\sum_{|q-k|\le 4}\|\nabla S_{k-1}a(t)\|_{L^\infty}\|\nabla\Delta_k u(t)\|_{L^2}\\
&\lesssim\sum_{|q-k|\le 4}d_k(t)2^{\frac{1}{2}k}2^{-\frac{3}{2}k}\|u(t)\|_{\dot{B}^{5/2}_{2,1}}\|a(t)\|_{\dot{B}^2_{2,1}}\\
&\lesssim d_q(t)2^{-q}\|u(t)\|_{\dot{B}^{5/2}_{2,1}}\|a(t)\|_{\dot{B}^2_{2,1}}.
\end{aligned}$$

Summing up all the above estimates, we arrive at

$$\|\mathrm{div}\,[\Delta_q,a]\nabla u(t)\|_{L^2}\lesssim d_q(t)2^{-q}\|a(t)\|_{\dot{B}^2_{2,1}}\|u(t)\|_{\dot{B}^{5/2}_{2,1}}.$$

Similarly, we have

$$\begin{aligned}
[\nabla a\cdot\nabla,\Delta_q]u(t)=&[T_{\nabla a},\Delta_q]\nabla u-\Delta_q\mathcal{R}(\nabla a,\nabla u)\\
&-\Delta_q(T_{\nabla u}\nabla a)+R(\nabla a,\nabla\Delta_q u).
\end{aligned}$$

Then following the same line of reasoning as above, we obtain

$$\|[\nabla a\cdot\nabla,\Delta_q]u(t)\|_{L^2}\lesssim d_q(t)2^{-q}\|a(t)\|_{\dot{B}^2_{2,1}}\|u(t)\|_{\dot{B}^{5/2}_{2,1}}.$$

Finally, we have

$$\|\nabla a(t)\Pi(t)\|_{\dot{B}^1_{2,1}}\lesssim\|\nabla a(t)\|_{\dot{B}^1_{2,1}}\|\Pi(t)\|_{\dot{B}^{3/2}_{2,1}}.$$

At this point, we list all the different estimates related to the restriction of s. □

Next, we will give the proof of (5.8).

Proof. Applying Lemma 2.6 to (5.9) together with Gronwall's inequality gives

$$\begin{aligned}&\|(\nabla\bar{u},\nabla\bar{B})\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})}+\|(\nabla\bar{u},\nabla\bar{B})\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\\&\leq C\exp\left\{\int_0^t\|\bar{u}(\tau)\|_{\dot{B}^{5/2}_{2,1}}+\|\bar{B}(\tau)\|_{\dot{B}^{5/2}_{2,1}}\,d\tau\right\}\Big\{\|(\nabla\bar{u}_0,\nabla\bar{B}_0)\|_{\dot{B}^{1/2}_{2,1}}\\&\quad+\|\nabla\bar{a}\|_{\tilde{L}^\infty_t(\dot{H}^1)}\left(\|\nabla\bar{\Pi}\|_{\tilde{L}^1_t(\dot{H}^1)}+\|\bar{u}\|_{L^1_t(\dot{B}^3_{2,1})}\right)\Big\},\end{aligned}$$

where we also used (5.5). Then thanks to (5.1) and (5.5), we conclude for every $\eta>0$ that

$$\begin{aligned}&\|(\nabla\bar{u},\nabla\bar{B})\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})}+\|(\nabla\bar{u},\nabla\bar{B})\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\\&\leq C+C_\eta\left(\|\Delta\bar{u}\|_{L^1_t(L^2)}+\|\nabla\bar{\Pi}\|_{L^1_t(L^2)}\right)+\eta\left(\|\bar{u}\|_{L^1_t(\dot{B}^{7/2}_{2,1})}+\|\nabla\bar{\Pi}\|_{\tilde{L}^1_t(\dot{H}^{3/2})}\right)\end{aligned}\tag{7.1}$$

Notice that $\operatorname{div}\bar{u}=\operatorname{div}\bar{B}=0$, we get by taking div to (5.9) that

$$\begin{aligned}\operatorname{div}\left((1+\bar{a})\nabla\partial_i\bar{\Pi}\right)=&-\operatorname{div}\partial_i\left[(\bar{u}\cdot\nabla)\bar{u}\right]+\operatorname{div}\partial_i\left[\bar{a}\,\Delta\bar{u}\right]-\operatorname{div}\left[\partial_i\bar{a}\,\nabla\bar{\Pi}\right]\\&+\operatorname{div}\partial_i\left(\bar{B}\cdot\nabla\bar{B}\right)+\operatorname{div}\left(\partial_i\bar{a}\,\bar{B}\cdot\nabla\bar{B}\right)\\&+\operatorname{div}\left(\bar{a}\,\partial_i\bar{B}\cdot\nabla\bar{B}\right)+\operatorname{div}\left(\bar{a}\,\bar{B}\cdot\nabla\partial_i\bar{B}\right).\end{aligned}$$

From this and Lemma 2.5, we deduce that

$$\begin{aligned}\|\nabla^2\bar{\Pi}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}\lesssim&\|\partial_i[\bar{u}\cdot\nabla\bar{u}]\|_{L^1_t(\dot{B}^{1/2}_{2,1})}+\|\partial_i[\bar{a}\,\Delta\bar{u}]\|_{L^1_t(\dot{B}^{1/2}_{2,1})}\\&+\|\partial_i\bar{a}\,\nabla\bar{\Pi}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}+\|\partial_i(\bar{B}\cdot\nabla\bar{B})\|_{L^1_t(\dot{B}^{1/2}_{2,1})}\\&+\|\partial\bar{a}\,\bar{B}\cdot\nabla\bar{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}+\|\bar{a}\,\partial_i\bar{B}\cdot\nabla\bar{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}\\&+\|\bar{a}\,\bar{B}\cdot\nabla\partial_i\bar{B}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}.\end{aligned}$$

Applying the product laws in Besov spaces yields that for any $\epsilon>0$

$$\begin{aligned}\|\nabla^2\bar{\Pi}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}\lesssim&\|\bar{u}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\bar{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\\&+\|\nabla\bar{a}\|_{\tilde{L}^\infty_t(\dot{B}^{3/2}_{2,1})}\|\Delta\bar{u}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}+\|\bar{a}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\bar{u}\|_{L^1_t(\dot{H}^{7/2})}\\&+\|\bar{B}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\bar{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}+\|\bar{a}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\bar{B}\|_{L^\infty_t(\dot{B}^{3/2}_{2,1})}\|\bar{B}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}\\&+\|\bar{a}\|_{\tilde{L}^\infty_t(\dot{H}^2)}\left(\epsilon\|\nabla^2\bar{\Pi}\|_{\tilde{L}^1_t(\dot{H}^{1/2})}+C_\eta\|\nabla\bar{\Pi}\|_{L^1_t(L^2)}\right).\end{aligned}$$

Thanks to Theorem 1.2, Proposition 5.1 and estimates (5.5), we get by taking ϵ sufficiently small in the above inequality that

$$\|\nabla^2\bar{\Pi}\|_{L^1_t(\dot{B}^{1/2}_{2,1})}\leq C\left(1+\|\nabla\bar{u}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})}+\|\nabla\bar{u}\|_{L^1_t(\dot{B}^{5/2}_{2,1})}+\|\nabla\bar{B}\|_{\tilde{L}^\infty_t(\dot{B}^{1/2}_{2,1})}\right).$$

Substituting the above inequality into (7.1) and then taking η sufficiently small, we arrive at

$$\|\nabla\bar{u}\|_{\tilde{L}_t^\infty(\dot{B}^{1/2}_{2,1})}+\|\nabla\bar{B}\|_{\tilde{L}_t^\infty(\dot{B}^{1/2}_{2,1})}+\|\nabla\bar{u}\|_{L_t^1(\dot{B}^{5/2}_{2,1})} + \|\nabla\bar{B}\|_{L_t^1(\dot{B}^{5/2}_{2,1})}+\|\nabla\bar{\Pi}\|_{L_t^1(\dot{B}^{3/2}_{2,1})}\leq C.$$

□

8. Acknowledgements

J. Peng and J. Jia's research is support partially by National Natural Science Foundation of China under the grant no.11131006, and by the National Basic Research Program of China under the grant no.2013CB329404. K. Li's research is support partially by National Natural Science Foundation of China under the grant no.11201366.

References

[1] P.A. Davidson, An Introduction to Magnetohydrodynamics, Cambridge University Press, Cambridge, 2001.
[2] X. Huang, Y. Wang, Global strong solution to the 2D nonhomogeneous incompressible MHD system Journal of Differential Equations, 254, 511-527 (2013).
[3] H. Abidi, M. Paicu, Global existence for the MHD system in critical spaces, Proceedings of the Royal Society of Edinburgh: Section A Mathematics, 138, 447-476 (2008).
[4] A. V. Kazhikov, Resolution of boundary value problems for nonhomogeneous viscous fluids, Dokl. Akad. Nauk, 216, 1008-1010 (1974).
[5] S. A. Antontesv, A. V. Kazhikov and V. N. Monakhov,Boundary Value Problems in Mechanics of Nonhomogeneous Fluids, North-Holland, Amsterdam, 1990.
[6] R. Danchin, P. B. Mucha, A lagrangian approach for the incompressible Navier-Stokes equations with variable density, Communications on Pure and Applied Mathematics, Vol. LXV, 1458-1480 (2012).
[7] H. Abidi, G. Gui, P. Zhang, On the wellposedness of three-dimensional inhomogeneous Navier-Stokes equations in the critical spaces, Arch. Rational Mech. Anal. 204, 189-230 (2012).
[8] J. Y. Chemin, M. Paicu, P. Zhang, Global large solutions to 3-D inhomogeneous Navier-Stokes system with one slow variable, arXiv:1301.6313
[9] R. Danchin, P. Zhang, Inhomogeneous Navier-Stokes equations in the half-space, with only bounded density, arXiv:1304.3235
[10] R. Danchin, P. B. Mucha, Incompressible flows with piecewise constant density, Archive for Rational Mechanics and Analysis, 207, 991-1023 (2013).
[11] G. Duraut, J. L. lions, Inquations en thermolasticit et magntohydrodynamique. (French) Arch. Rational Mech. Anal., 46, 241-279 (1972).
[12] C. Cao, J. Wu, Global regularity for the 2D MHD equations with mixed partial dissipation and magnetic diffusion, Advances in Mathematics, 226, 1803-1822 (2011).
[13] C. Cao, D. Regmi, J. Wu, The 2D MHD equations with horizontal dissipation and horizontal magnetic diffusion, Journal of Differential Equations, 254, 2661-2681 (2013).
[14] C. Cao, J. Wu, B. Yuan, The 2D Incompressible Magnetohydrodynamics Equations with only Magnetic Diffusion, arXiv:1306.3629.
[15] G. Ponce,R. Racke, T. C. Sideris, E. S. Titi,Global stability of large solutions to the 3D Navier-Stokes equations. Comm. Math. Phys., 159, 329C341 (1994).
[16] I. Gallagher, D. Iftimie, F. Planchon, Asymptotics and stability for global solutions to the Navier-Stokes equations. Ann. Inst. Fourier (Grenoble) 53 , 1387C1424 (2003).
[17] R. Danchin, Fourier analysis methods for PDE's, (2005).
[18] H. Bahouri, J.-Y. Chemin and R. Danchin, Fourier Analysis and Nonlinear Partial Differential Equations, Grundlehren der Mathematischen Wissenschaften, Vol. 343 (Springer, 2011).
[19] Chemin, J.-Y.; Lerner, N. Flot de champs de vecteurs non lipschitziens et équations de Navier-Stokes, Journal of Differential Equations, 121, 314-328 (1992).

[20] G. Gui, P. Zhang, Stability to the global large solutions of 3-D Navier-Stokes equations, Advances in Mathematics, 225, 1248-1284 (2010).
[21] H. Abidi, G. Gui, P. Zhang, On the Decay and Stability of Global Solutions to the 3D Inhomogeneous Navier-Stokes Equations, Communications on Pure and Applied Mahtematics,Vol. LXIV, 0832-0881 (2011).
[22] R. Danchin, Local and global well-posedness results for flows of inhomogeneous viscous fluids, Advances in Differential Equations, 9, 353-386 (2004).
[23] M. E. Schonbek, Large time behavior of solutions to the Navier-Stokes equation. Communications in Partial Differential Equations, 11, 733-763(1986).
[24] L. Grafakos. Classical Fourier analysis, Graduate Texts in Mathematics, Vol. 249 (Springer, New York, 2008).
[25] Hmidi,T., Keraani,S., Rousset,F. Global well-posedness for a Boussinesq-Navier-Stokes system with critical dissipation, Journal of Differential Equations, 249, 2147-2174 (2010).

School of Mathematics and Statistics, Xi'an Jiaotong University, Xi'an 710049, China;
Beijing Center for Mathematics and Information Interdisciplinary Sciences (BCMIIS);
E-mail address: jjx425@gmail.com

School of Mathematics and Statistics, Xi'an Jiaotong University, Xi'an 710049, China;
Beijing Center for Mathematics and Information Interdisciplinary Sciences (BCMIIS);
E-mail address: jgpeng@mail.xjtu.edu.cn

School of Mathematics and Statistics, Xi'an Jiaotong University, Xi'an 710049, China;
Beijing Center for Mathematics and Information Interdisciplinary Sciences (BCMIIS);
E-mail address: kexueli@gmail.com